



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2024 – 03:57 AM EST

PDB ID : 3GSZ
Title : Structure of the genotype 2B HCV polymerase
Authors : Rydberg, E.H.; Carfi, A.
Deposited on : 2009-03-27
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	563	 87% 10% ..
1	B	563	 89% 9% ..

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 9571 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA-directed RNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	558	4373	2759	778	809	27	0	0	0
1	B	558	4373	2759	778	809	27	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	5	THR	SER	conflict	UNP P26661
B	5	THR	SER	conflict	UNP P26661


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	452	Total	O	0	0
			452	452		
2	B	373	Total	O	0	0
			373	373		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: RNA-directed RNA polymerase



- Molecule 1: RNA-directed RNA polymerase



4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	153.25Å 64.50Å 135.58Å 90.00° 90.16° 90.00°	Depositor
Resolution (Å)	19.96 – 1.90 19.96 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.96-1.90) 96.8 (19.96-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.84 (at 1.90Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.166 , 0.210 0.177 , 0.218	Depositor DCC
R_{free} test set	5127 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	18.7	Xtrriage
Anisotropy	0.255	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 30.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	0.097 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9571	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.13	3/4471 (0.1%)	1.04	22/6066 (0.4%)
1	B	1.08	2/4471 (0.0%)	1.03	20/6066 (0.3%)
All	All	1.10	5/8942 (0.1%)	1.03	42/12132 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	187	MET	SD-CE	-6.73	1.40	1.77
1	B	470	GLU	CD-OE1	6.10	1.32	1.25
1	A	321	VAL	CB-CG2	-6.00	1.40	1.52
1	B	218	SER	CB-OG	-6.00	1.34	1.42
1	A	514	GLN	CB-CG	-5.15	1.38	1.52

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	544	ARG	NE-CZ-NH2	-8.59	116.00	120.30
1	B	234	ARG	NE-CZ-NH1	-8.04	116.28	120.30
1	B	498	ARG	NE-CZ-NH1	7.77	124.19	120.30
1	A	164	ASP	CB-CG-OD2	7.75	125.28	118.30
1	B	56	ARG	NE-CZ-NH2	-6.95	116.83	120.30
1	B	129	ASP	CB-CG-OD2	6.90	124.51	118.30
1	A	553	VAL	C-N-CA	-6.77	108.08	122.30
1	B	164	ASP	CB-CG-OD2	6.77	124.39	118.30
1	A	337	ARG	NE-CZ-NH2	6.61	123.60	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	310	ASP	CB-CG-OD2	6.48	124.13	118.30
1	A	278	ARG	NE-CZ-NH2	-6.30	117.15	120.30
1	A	277	ARG	NE-CZ-NH2	-6.10	117.25	120.30
1	B	544	ARG	CG-CD-NE	-6.08	99.03	111.80
1	B	125	ASP	CB-CG-OD2	6.03	123.72	118.30
1	A	508	ARG	NE-CZ-NH2	-5.88	117.36	120.30
1	B	319	ASP	CB-CG-OD2	5.84	123.56	118.30
1	B	355	ARG	NE-CZ-NH1	5.84	123.22	120.30
1	A	359	ASP	CB-CG-OD1	5.83	123.55	118.30
1	A	508	ARG	NE-CZ-NH1	5.79	123.20	120.30
1	B	65	ASP	CB-CG-OD2	5.79	123.51	118.30
1	B	536	LEU	CB-CG-CD1	5.77	120.81	111.00
1	A	332	ASP	CB-CG-OD2	5.74	123.47	118.30
1	A	386	ARG	NE-CZ-NH1	5.71	123.16	120.30
1	B	234	ARG	NE-CZ-NH2	5.71	123.15	120.30
1	A	65	ASP	CB-CG-OD1	5.70	123.43	118.30
1	B	377	ARG	NE-CZ-NH1	5.64	123.12	120.30
1	A	278	ARG	NE-CZ-NH1	5.54	123.07	120.30
1	A	73	ARG	NE-CZ-NH2	-5.52	117.54	120.30
1	B	81	ARG	NE-CZ-NH1	5.50	123.05	120.30
1	B	259	ARG	NE-CZ-NH2	-5.46	117.57	120.30
1	A	82	LEU	CB-CG-CD1	5.44	120.24	111.00
1	B	278	ARG	NE-CZ-NH1	5.38	122.99	120.30
1	A	32	ARG	NE-CZ-NH1	5.36	122.98	120.30
1	A	439	LEU	CA-CB-CG	5.34	127.59	115.30
1	A	32	ARG	NE-CZ-NH2	-5.28	117.66	120.30
1	B	387	ASP	CB-CG-OD2	5.25	123.03	118.30
1	A	182	LEU	CB-CG-CD2	5.16	119.77	111.00
1	A	220	ASP	CB-CG-OD2	5.15	122.94	118.30
1	B	377	ARG	NE-CZ-NH2	-5.12	117.74	120.30
1	A	280	ARG	NE-CZ-NH2	-5.07	117.77	120.30
1	A	168	ARG	NE-CZ-NH1	5.05	122.83	120.30
1	B	406	ASN	CB-CA-C	-5.04	100.33	110.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	465	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4373	0	4380	29	0
1	B	4373	0	4380	27	0
2	A	452	0	0	10	0
2	B	373	0	0	8	0
All	All	9571	0	8760	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:541:GLU:HG2	2:A:760:HOH:O	1.58	1.01
1:B:82:LEU:HD13	1:B:249:ALA:HB2	1.62	0.81
1:A:527:ASN:HD21	1:A:534:LEU:H	1.25	0.81
1:B:527:ASN:HD21	1:B:534:LEU:H	1.31	0.78
1:B:405:VAL:HG11	1:B:446:GLU:HG3	1.79	0.64
1:B:118:HIS:HE1	2:B:660:HOH:O	1.80	0.64
1:B:541:GLU:HG2	2:B:809:HOH:O	1.98	0.63
1:B:381:ARG:NE	1:B:470:GLU:HG2	2.15	0.62
1:B:461:ALA:HB2	1:B:541:GLU:HG3	1.83	0.61
1:B:83:LEU:HB2	1:B:173:MET:HA	1.82	0.61
1:B:452:TYR:OH	1:B:550:TRP:O	2.15	0.60
1:A:118:HIS:HE1	2:A:685:HOH:O	1.85	0.59
1:B:234:ARG:NH1	1:B:258:GLU:OE2	2.36	0.59
1:B:461:ALA:CB	1:B:541:GLU:HG3	2.34	0.58
1:A:458:ASP:OD1	1:A:517:ARG:NH1	2.36	0.57
1:A:337:ARG:HD3	2:A:842:HOH:O	2.06	0.55
1:A:20:LYS:HD2	2:A:781:HOH:O	2.07	0.54
1:A:531:LYS:HG2	2:A:910:HOH:O	2.07	0.54
1:B:523:ARG:HD2	2:B:851:HOH:O	2.08	0.54
1:B:380:ARG:CD	2:B:1024:HOH:O	2.57	0.53
1:A:546:ASP:OD1	1:A:546:ASP:N	2.37	0.52
1:A:34:HIS:HD2	2:A:1212:HOH:O	1.93	0.52
1:A:552:THR:HB	2:A:836:HOH:O	2.10	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:398:GLU:CD	2:A:935:HOH:O	2.47	0.51
1:B:234:ARG:HH12	1:B:258:GLU:CD	2.14	0.51
1:A:160:ILE:HD12	1:A:282:SER:OG	2.11	0.51
1:B:77:LYS:NZ	2:B:1090:HOH:O	2.44	0.51
1:A:289:MET:O	1:A:293:MET:HG3	2.12	0.50
1:B:562:HIS:O	1:B:563:SER:HB2	2.12	0.49
1:B:182:LEU:HB3	1:B:183:PRO:HD3	1.95	0.49
1:A:465:ARG:NH1	1:A:545:LEU:O	2.46	0.49
1:B:186:ILE:HG13	1:B:187:MET:HG3	1.95	0.48
1:A:205:LEU:HD11	1:A:382:TYR:CG	2.49	0.48
1:B:222:ARG:HH11	1:B:222:ARG:HG2	1.79	0.47
1:A:21:LEU:HD12	1:A:22:PRO:HD2	1.96	0.47
1:A:95:HIS:CD2	1:A:404:PRO:HB3	2.50	0.46
1:A:83:LEU:HB2	1:A:173:MET:HA	1.96	0.46
1:B:19:GLU:HG3	1:B:20:LYS:HG3	1.98	0.45
1:A:452:TYR:CE1	1:A:562:HIS:HB2	2.52	0.45
1:A:372:VAL:HG12	2:A:985:HOH:O	2.16	0.45
1:A:465:ARG:HA	1:A:465:ARG:HD2	1.79	0.44
1:A:405:VAL:HG11	1:A:446:GLU:HG3	1.99	0.44
1:B:172:LYS:HE3	1:B:560:ILE:HD12	1.99	0.44
1:A:461:ALA:HB2	1:A:541:GLU:HG3	2.00	0.43
1:A:334:ARG:HG2	1:A:337:ARG:NH1	2.33	0.43
1:A:37:VAL:HG22	1:A:393:THR:HG23	2.01	0.43
1:B:37:VAL:HG13	1:B:393:THR:CG2	2.49	0.43
1:B:218:SER:HB2	2:B:787:HOH:O	2.19	0.42
1:B:380:ARG:NE	2:B:1024:HOH:O	2.52	0.42
1:A:321:VAL:HG21	1:A:365:SER:HB2	2.01	0.42
1:B:234:ARG:HD2	2:B:975:HOH:O	2.19	0.42
1:A:444:ASN:HA	1:A:452:TYR:O	2.19	0.41
1:A:219:TYR:HB3	1:A:320:LEU:HD23	2.03	0.41
1:A:465:ARG:CZ	2:A:710:HOH:O	2.69	0.40
1:B:48:ARG:HG3	1:B:159:LEU:HD22	2.02	0.40
1:B:364:THR:CG2	1:B:367:SER:HA	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	554/563 (98%)	546 (99%)	8 (1%)	0	100	100
1	B	554/563 (98%)	542 (98%)	12 (2%)	0	100	100
All	All	1108/1126 (98%)	1088 (98%)	20 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	475/478 (99%)	461 (97%)	14 (3%)	42	35
1	B	475/478 (99%)	463 (98%)	12 (2%)	47	41
All	All	950/956 (99%)	924 (97%)	26 (3%)	44	38

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	ARG
1	A	82	LEU
1	A	83	LEU
1	A	100	ARG
1	A	159	LEU
1	A	182	LEU
1	A	184	LYS
1	A	439	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	527	ASN
1	A	531	LYS
1	A	536	LEU
1	A	546	ASP
1	A	548	SER
1	A	553	VAL
1	B	56	ARG
1	B	73	ARG
1	B	82	LEU
1	B	159	LEU
1	B	182	LEU
1	B	402	HIS
1	B	403	SER
1	B	439	LEU
1	B	527	ASN
1	B	535	LYS
1	B	536	LEU
1	B	560	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	HIS
1	A	95	HIS
1	A	118	HIS
1	A	329	ASN
1	A	335	ASN
1	A	527	ASN
1	B	63	HIS
1	B	118	HIS
1	B	254	HIS
1	B	335	ASN
1	B	514	GLN
1	B	527	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	558/563 (99%)	-0.28	4 (0%) 87 88	11, 20, 34, 45	0
1	B	558/563 (99%)	-0.19	10 (1%) 68 71	12, 23, 39, 52	0
All	All	1116/1126 (99%)	-0.23	14 (1%) 77 79	11, 21, 37, 52	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	553	VAL	4.4
1	A	14	CYS	4.0
1	B	14	CYS	4.0
1	A	563	SER	4.0
1	B	563	SER	3.3
1	B	402	HIS	3.3
1	B	16	PRO	3.1
1	B	552	THR	2.9
1	B	334	ARG	2.5
1	B	548	SER	2.5
1	A	334	ARG	2.4
1	B	448	TYR	2.2
1	A	546	ASP	2.1
1	B	379	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.