



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 27, 2023 – 10:33 AM EDT

PDB ID : 3HVA  
Title : Crystal structure of FimX GGDEF domain from *Pseudomonas aeruginosa*  
Authors : Navarro, M.V.A.S.; De, N.; Bae, N.; Sondermann, H.  
Deposited on : 2009-06-15  
Resolution : 2.04 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.04 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	177	 4% 53% 37% 8%
1	B	177	 6% 59% 32% 8%

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 2599 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein FimX.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	162	1228	763	226	235	4	0	0	0
1	B	162	1228	763	226	235	4	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	255	GLY	-	expression tag	UNP Q9HUK6
A	256	ALA	-	expression tag	UNP Q9HUK6
A	257	MET	-	expression tag	UNP Q9HUK6
A	258	GLY	-	expression tag	UNP Q9HUK6
A	259	SER	-	expression tag	UNP Q9HUK6
B	255	GLY	-	expression tag	UNP Q9HUK6
B	256	ALA	-	expression tag	UNP Q9HUK6
B	257	MET	-	expression tag	UNP Q9HUK6
B	258	GLY	-	expression tag	UNP Q9HUK6
B	259	SER	-	expression tag	UNP Q9HUK6


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	78	Total	O	0	0
			78	78		
2	B	65	Total	O	0	0
			65	65		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Protein FimX



- Molecule 1: Protein FimX



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	33.18Å 94.11Å 45.50Å 90.00° 90.61° 90.00°	Depositor
Resolution (Å)	27.12 – 2.04 27.12 – 2.04	Depositor EDS
% Data completeness (in resolution range)	94.0 (27.12-2.04) 93.4 (27.12-2.04)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.78 (at 2.04Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.213 , 0.246 0.217 , 0.237	Depositor DCC
$R_{free}$ test set	859 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.3	Xtrriage
Anisotropy	0.153	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 28.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.197 for h,-k,-l	Xtrriage
Reported twinning fraction	0.191 for h,-k,-l	Depositor
Outliers	0 of 16942 reflections	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	2599	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.62% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.22	0/1245	0.41	0/1676
1	B	0.24	0/1245	0.43	0/1676
All	All	0.23	0/2490	0.42	0/3352

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1228	0	1227	46	0
1	B	1228	0	1227	46	0
2	A	78	0	0	5	0
2	B	65	0	0	5	0
All	All	2599	0	2454	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

All (92) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:273:THR:HG23	1:A:275:LEU:H	1.30	0.94

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:355:LYS:HD3	1:B:355:LYS:H	1.48	0.77
1:A:297:LYS:HE2	1:A:356:GLY:HA2	1.66	0.77
1:A:296:ARG:HG2	1:A:398:GLU:HG2	1.71	0.72
1:A:317:GLY:O	1:A:321:ILE:HG22	1.89	0.72
1:A:363:GLN:NE2	1:A:367:GLN:HE21	1.89	0.70
1:B:309:TYR:N	1:B:310:PRO:HD2	2.10	0.66
1:A:370:LEU:HD22	1:A:425:ILE:HG13	1.77	0.65
1:B:268:SER:HB2	1:B:280:HIS:HB2	1.79	0.65
1:B:296:ARG:HD2	1:B:398:GLU:OE2	1.98	0.64
1:B:373:VAL:HG21	1:B:389:LEU:HD11	1.79	0.64
1:B:360:GLU:HG3	2:B:85:HOH:O	1.97	0.64
1:A:297:LYS:HE2	1:A:356:GLY:CA	2.28	0.63
1:B:355:LYS:HD3	1:B:355:LYS:N	2.14	0.63
1:A:373:VAL:HG21	1:A:389:LEU:HD11	1.82	0.61
1:B:333:ARG:HG3	1:B:334:GLU:N	2.17	0.60
1:A:316:HIS:HB3	1:A:320:GLY:HA3	1.84	0.60
1:A:386:GLN:HG2	2:A:2:HOH:O	2.01	0.59
1:B:345:PHE:HB3	1:B:349:ILE:HG22	1.85	0.58
1:B:397:ASP:OD1	1:B:399:LYS:N	2.35	0.58
1:A:396:LEU:HD21	1:A:406:VAL:HG21	1.86	0.58
1:B:330:GLY:HA2	1:B:333:ARG:HG2	1.85	0.58
1:B:340:ALA:HB2	1:B:354:PHE:CE1	2.38	0.57
1:A:396:LEU:CD2	1:A:406:VAL:HG21	2.36	0.56
1:A:397:ASP:OD1	1:A:399:LYS:N	2.37	0.55
1:A:397:ASP:OD1	1:A:399:LYS:HB2	2.06	0.55
1:A:391:ILE:HB	1:A:425:ILE:HG12	1.89	0.55
1:B:309:TYR:CG	1:B:348:SER:HB3	2.43	0.53
1:A:371:LYS:HE3	2:A:87:HOH:O	2.08	0.53
1:A:309:TYR:CG	1:A:348:SER:HB3	2.44	0.52
1:A:313:GLN:HE22	1:A:318:LEU:HD23	1.73	0.52
1:A:360:GLU:H	1:A:360:GLU:CD	2.13	0.52
1:A:335:GLN:OE1	1:A:372:LYS:HG2	2.09	0.51
1:A:376:HIS:HD2	2:A:123:HOH:O	1.92	0.51
1:B:380:LEU:O	1:B:383:ARG:HB2	2.10	0.51
1:A:299:SER:OG	1:A:355:LYS:HA	2.10	0.51
1:A:344:ARG:HH21	1:A:347:ASP:HA	1.75	0.51
1:B:367:GLN:HG3	1:B:371:LYS:HE3	1.92	0.51
1:B:324:LEU:HG	2:B:162:HOH:O	2.11	0.50
1:A:289:VAL:HG22	1:A:396:LEU:CD2	2.40	0.50
1:B:331:LEU:O	1:B:335:GLN:HB2	2.12	0.50
1:A:363:GLN:HE21	1:A:367:GLN:HE21	1.60	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:345:PHE:HB3	1:B:349:ILE:CG2	2.41	0.50
1:B:418:ALA:HB1	2:B:75:HOH:O	2.11	0.49
1:B:305:HIS:HD2	1:B:414:ALA:HB1	1.76	0.49
1:B:318:LEU:O	1:B:321:ILE:HG22	2.12	0.49
1:B:325:LEU:HD23	2:B:162:HOH:O	2.12	0.49
1:A:332:MET:HE2	1:A:342:LEU:HD22	1.95	0.49
1:B:289:VAL:HG22	1:B:396:LEU:HD21	1.95	0.48
1:A:370:LEU:HG	1:A:423:SER:HB3	1.94	0.48
1:B:335:GLN:HE21	1:B:368:ARG:HE	1.61	0.48
1:A:289:VAL:HG22	1:A:396:LEU:HD21	1.94	0.48
1:B:293:VAL:HG23	1:B:294:THR:HG23	1.95	0.48
1:B:325:LEU:HA	2:B:162:HOH:O	2.14	0.47
1:B:289:VAL:HG21	1:B:403:ALA:CB	2.44	0.47
1:A:336:PHE:CZ	1:A:369:LEU:HD22	2.50	0.47
1:A:325:LEU:HD22	1:A:350:PHE:CZ	2.50	0.47
1:B:360:GLU:O	1:B:363:GLN:HG2	2.15	0.47
1:A:405:ASP:O	1:A:409:ARG:HG3	2.15	0.47
1:B:303:TYR:OH	1:B:349:ILE:HD13	2.15	0.46
1:A:340:ALA:HB2	1:A:354:PHE:CE1	2.50	0.46
1:B:341:ASP:OD2	1:B:355:LYS:HD2	2.15	0.46
1:B:332:MET:CE	1:B:342:LEU:HD22	2.47	0.45
1:B:416:ASP:HA	1:B:419:ARG:CZ	2.46	0.45
1:A:302:ALA:HB3	1:A:352:ALA:HB3	1.98	0.45
1:B:299:SER:OG	1:B:355:LYS:HA	2.16	0.45
1:A:298:PRO:HA	1:A:396:LEU:O	2.16	0.44
1:B:268:SER:HA	1:B:277:ASN:ND2	2.32	0.44
1:B:383:ARG:HE	1:B:383:ARG:HB3	1.62	0.44
1:A:307:ASN:HB2	1:A:388:THR:O	2.18	0.44
1:B:416:ASP:HA	1:B:419:ARG:NH2	2.32	0.44
1:A:309:TYR:CD2	1:A:348:SER:HB3	2.53	0.44
1:A:270:ASP:HB3	1:A:273:THR:HG22	1.99	0.44
1:B:313:GLN:HE22	1:B:318:LEU:N	2.15	0.44
1:B:336:PHE:CZ	1:B:369:LEU:HD22	2.53	0.44
1:A:401:ALA:O	1:A:402:LYS:HG3	2.19	0.43
1:A:300:THR:HB	1:A:354:PHE:HB2	1.99	0.42
1:A:288:ALA:HB2	1:A:355:LYS:HE2	2.01	0.42
1:B:313:GLN:NE2	1:B:317:GLY:C	2.73	0.42
1:B:289:VAL:HG21	1:B:403:ALA:HB2	2.00	0.42
1:A:321:ILE:HB	2:A:49:HOH:O	2.19	0.42
1:A:353:LEU:HD21	1:A:355:LYS:HE2	2.02	0.42
1:A:309:TYR:N	1:A:310:PRO:CD	2.83	0.41

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:340:ALA:HB2	1:B:354:PHE:CD1	2.55	0.41
1:A:296:ARG:HG2	1:A:398:GLU:CG	2.44	0.41
2:A:41:HOH:O	1:B:266:VAL:HA	2.21	0.41
1:A:270:ASP:HA	1:A:271:PRO:HD3	1.86	0.41
1:A:313:GLN:NE2	1:A:318:LEU:HD23	2.34	0.41
1:B:268:SER:HA	1:B:277:ASN:HD21	1.86	0.41
1:B:296:ARG:HB3	1:B:296:ARG:NH2	2.35	0.40
1:B:313:GLN:HE22	1:B:317:GLY:C	2.25	0.40
1:B:312:LEU:HD13	1:B:324:LEU:CD2	2.51	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	160/177 (90%)	159 (99%)	1 (1%)	0	100	100
1	B	160/177 (90%)	160 (100%)	0	0	100	100
All	All	320/354 (90%)	319 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	125/136 (92%)	114 (91%)	11 (9%)	10	4
1	B	125/136 (92%)	119 (95%)	6 (5%)	25	18
All	All	250/272 (92%)	233 (93%)	17 (7%)	16	8

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	269	GLN
1	A	290	GLN
1	A	291	GLN
1	A	312	LEU
1	A	321	ILE
1	A	333	ARG
1	A	338	GLU
1	A	379	GLU
1	A	383	ARG
1	A	386	GLN
1	A	398	GLU
1	B	290	GLN
1	B	311	SER
1	B	355	LYS
1	B	390	SER
1	B	398	GLU
1	B	427	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	269	GLN
1	A	290	GLN
1	A	307	ASN
1	A	313	GLN
1	A	363	GLN
1	A	376	HIS
1	A	386	GLN
1	B	280	HIS
1	B	290	GLN
1	B	307	ASN
1	B	313	GLN
1	B	335	GLN
1	B	363	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	411	HIS
1	B	427	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	162/177 (91%)	0.52	7 (4%) 35 38	15, 23, 41, 57	0
1	B	162/177 (91%)	0.59	11 (6%) 17 18	14, 24, 40, 53	0
All	All	324/354 (91%)	0.56	18 (5%) 24 26	14, 24, 43, 57	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	267	SER	5.1
1	A	401	ALA	4.9
1	B	401	ALA	4.2
1	A	266	VAL	4.2
1	A	419	ARG	2.8
1	B	267	SER	2.6
1	B	333	ARG	2.5
1	A	296	ARG	2.5
1	A	400	THR	2.4
1	A	399	LYS	2.3
1	B	266	VAL	2.3
1	B	400	THR	2.3
1	B	381	ASN	2.2
1	B	416	ASP	2.2
1	B	345	PHE	2.1
1	B	268	SER	2.1
1	B	402	LYS	2.1
1	B	296	ARG	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.