



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 31, 2023 – 12:37 PM EDT

PDB ID : 3LTN  
Title : Inhibitor-stabilized topoisomerase IV-DNA cleavage complex (*S. pneumoniae*)  
Authors : Laponogov, I.; Pan, X.-S.; Veselkov, D.A.; McAuley, K.E.; Fisher, L.M.; Sanderson, M.R.  
Deposited on : 2010-02-16  
Resolution : 3.10 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RSRZ outliers	127900	1067 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	496	 55% 37% 5% •
1	B	496	 54% 38% 5% •
2	C	268	 40% 37% 7% 17%
2	D	268	 39% 35% 8% 17%
3	E	15	 20% 20% 7% 53%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
4	F	19	
5	G	15	
6	H	19	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
8	PDQ	F	0	-	-	X	-

## 2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 11119 atoms, of which 48 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA topoisomerase 4 subunit A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	481	Total 3529	C 2224	N 619	O 672	S 14	0	0	0
1	B	481	Total 3523	C 2218	N 619	O 672	S 14	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	489	LEU	-	EXPRESSION TAG	UNP P72525
A	490	GLU	-	EXPRESSION TAG	UNP P72525
A	491	HIS	-	EXPRESSION TAG	UNP P72525
A	492	HIS	-	EXPRESSION TAG	UNP P72525
A	493	HIS	-	EXPRESSION TAG	UNP P72525
A	494	HIS	-	EXPRESSION TAG	UNP P72525
A	495	HIS	-	EXPRESSION TAG	UNP P72525
A	496	HIS	-	EXPRESSION TAG	UNP P72525
B	489	LEU	-	EXPRESSION TAG	UNP P72525
B	490	GLU	-	EXPRESSION TAG	UNP P72525
B	491	HIS	-	EXPRESSION TAG	UNP P72525
B	492	HIS	-	EXPRESSION TAG	UNP P72525
B	493	HIS	-	EXPRESSION TAG	UNP P72525
B	494	HIS	-	EXPRESSION TAG	UNP P72525
B	495	HIS	-	EXPRESSION TAG	UNP P72525
B	496	HIS	-	EXPRESSION TAG	UNP P72525

- Molecule 2 is a protein called DNA topoisomerase 4 subunit B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	222	Total 1615	C 1018	N 286	O 303	S 8	0	0	0
2	D	222	Total 1615	C 1018	N 286	O 303	S 8	0	0	0

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	380	MET	-	INITIATING METHIONINE	UNP Q59961
C	381	GLY	-	EXPRESSION TAG	UNP Q59961
C	382	HIS	-	EXPRESSION TAG	UNP Q59961
C	383	HIS	-	EXPRESSION TAG	UNP Q59961
C	384	HIS	-	EXPRESSION TAG	UNP Q59961
C	385	HIS	-	EXPRESSION TAG	UNP Q59961
C	386	HIS	-	EXPRESSION TAG	UNP Q59961
C	387	HIS	-	EXPRESSION TAG	UNP Q59961
C	388	HIS	-	EXPRESSION TAG	UNP Q59961
C	389	HIS	-	EXPRESSION TAG	UNP Q59961
C	390	HIS	-	EXPRESSION TAG	UNP Q59961
C	391	HIS	-	EXPRESSION TAG	UNP Q59961
C	392	SER	-	EXPRESSION TAG	UNP Q59961
C	393	SER	-	EXPRESSION TAG	UNP Q59961
C	394	GLY	-	EXPRESSION TAG	UNP Q59961
C	395	HIS	-	EXPRESSION TAG	UNP Q59961
C	396	ILE	-	EXPRESSION TAG	UNP Q59961
C	397	ASP	-	EXPRESSION TAG	UNP Q59961
C	398	ASP	-	EXPRESSION TAG	UNP Q59961
C	399	ASP	-	EXPRESSION TAG	UNP Q59961
C	400	ASP	-	EXPRESSION TAG	UNP Q59961
C	401	LYS	-	EXPRESSION TAG	UNP Q59961
C	402	HIS	-	EXPRESSION TAG	UNP Q59961
C	403	MET	-	EXPRESSION TAG	UNP Q59961
D	380	MET	-	INITIATING METHIONINE	UNP Q59961
D	381	GLY	-	EXPRESSION TAG	UNP Q59961
D	382	HIS	-	EXPRESSION TAG	UNP Q59961
D	383	HIS	-	EXPRESSION TAG	UNP Q59961
D	384	HIS	-	EXPRESSION TAG	UNP Q59961
D	385	HIS	-	EXPRESSION TAG	UNP Q59961
D	386	HIS	-	EXPRESSION TAG	UNP Q59961
D	387	HIS	-	EXPRESSION TAG	UNP Q59961
D	388	HIS	-	EXPRESSION TAG	UNP Q59961
D	389	HIS	-	EXPRESSION TAG	UNP Q59961
D	390	HIS	-	EXPRESSION TAG	UNP Q59961
D	391	HIS	-	EXPRESSION TAG	UNP Q59961
D	392	SER	-	EXPRESSION TAG	UNP Q59961
D	393	SER	-	EXPRESSION TAG	UNP Q59961
D	394	GLY	-	EXPRESSION TAG	UNP Q59961
D	395	HIS	-	EXPRESSION TAG	UNP Q59961
D	396	ILE	-	EXPRESSION TAG	UNP Q59961
D	397	ASP	-	EXPRESSION TAG	UNP Q59961

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	398	ASP	-	EXPRESSION TAG	UNP Q59961
D	399	ASP	-	EXPRESSION TAG	UNP Q59961
D	400	ASP	-	EXPRESSION TAG	UNP Q59961
D	401	LYS	-	EXPRESSION TAG	UNP Q59961
D	402	HIS	-	EXPRESSION TAG	UNP Q59961
D	403	MET	-	EXPRESSION TAG	UNP Q59961

- Molecule 3 is a DNA chain called 5'-D(\*AP\*CP\*CP\*AP\*AP\*GP\*GP\*T\*CP\*AP\*TP\*GP\*AP\*AP\*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	E	7	140	69	27	38	6	0	0	0

- Molecule 4 is a DNA chain called 5'-D(P\*AP\*GP\*TP\*CP\*AP\*TP\*TP\*CP\*AP\*TP\*GP\*AP\*CP\*CP\*TP\*TP\*GP\*GP\*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
4	F	11	225	108	39	67	11	0	0	0

- Molecule 5 is a DNA chain called 5'-D(\*CP\*TP\*GP\*TP\*TP\*TP\*TP\*A\*CP\*GP\*TP\*GP\*CP\*AP\*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
5	G	7	139	68	25	40	6	0	0	0

- Molecule 6 is a DNA chain called 5'-D(P\*GP\*AP\*CP\*TP\*AP\*TP\*GP\*CP\*AP\*CP\*GP\*TP\*AP\*AP\*AP\*AP\*CP\*AP\*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
6	H	11	226	107	43	65	11	0	0	0

- Molecule 7 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	C	1	Total	Mg	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	D	1	Total	Mg	0	0
			1	1		

- Molecule 8 is 3-amino-7-((3R)-3-((1S)-1-aminoethyl)pyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-8-methylquinazoline-2,4(1H,3H)-dione (three-letter code: PDQ) (formula: C<sub>18</sub>H<sub>24</sub>FN<sub>5</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	F	H	N			O
8	F	1	Total	C	F	H	N	O	0	0
			50	18	1	24	5	2		
8	H	1	Total	C	F	H	N	O	0	0
			50	18	1	24	5	2		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	2	Total	O	0	0
			2	2		
9	B	2	Total	O	0	0
			2	2		
9	D	1	Total	O	0	0
			1	1		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA topoisomerase 4 subunit A




- Molecule 1: DNA topoisomerase 4 subunit A









• Molecule 2: DNA topoisomerase 4 subunit B




• Molecule 2: DNA topoisomerase 4 subunit B




• Molecule 3: 5'-D(\*AP\*CP\*CP\*AP\*AP\*GP\*GP\*T\*CP\*AP\*TP\*GP\*AP\*AP\*T)-3'




- Molecule 4: 5'-D(P\*AP\*GP\*TP\*CP\*AP\*TP\*TP\*CP\*AP\*TP\*GP\*AP\*CP\*CP\*TP\*TP\*GP\*GP\*T)-3'



- Molecule 5: 5'-D(\*CP\*TP\*GP\*TP\*TP\*TP\*TP\*A\*CP\*GP\*TP\*GP\*CP\*AP\*T)-3'



- Molecule 6: 5'-D(P\*GP\*AP\*CP\*TP\*AP\*TP\*GP\*CP\*AP\*CP\*GP\*TP\*AP\*AP\*AP\*AP\*CP\*AP\*G)-3'



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	122.51Å 122.51Å 178.30Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	34.13 – 3.10 34.13 – 3.10	Depositor EDS
% Data completeness (in resolution range)	99.2 (34.13-3.10) 99.1 (34.13-3.10)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.98 (at 3.12Å)	Xtriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, $R_{free}$	0.186 , 0.228 0.181 , 0.222	Depositor DCC
$R_{free}$ test set	5393 reflections (10.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	95.4	Xtriage
Anisotropy	0.096	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.26 , 73.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.479 for -h,-k,l 0.043 for h,-h-k,-l 0.041 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	11119	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	112.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.55% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, PDQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	0/3590	0.62	0/4885
1	B	0.38	0/3583	0.63	1/4876 (0.0%)
2	C	0.37	0/1641	0.61	0/2230
2	D	0.37	0/1641	0.61	0/2230
3	E	0.96	0/157	1.68	4/241 (1.7%)
4	F	0.86	0/251	1.72	9/385 (2.3%)
5	G	0.94	0/155	1.50	3/238 (1.3%)
6	H	0.75	0/253	1.66	8/388 (2.1%)
All	All	0.43	0/11271	0.76	25/15473 (0.2%)

There are no bond length outliers.

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	H	1	DG	O4'-C4'-C3'	-7.53	101.48	106.00
3	E	11	DT	N3-C4-O4	7.45	124.37	119.90
4	F	9	DA	O4'-C1'-N9	-7.16	102.99	108.00
3	E	11	DT	C5-C4-O4	-7.00	120.00	124.90
6	H	9	DA	O4'-C1'-N9	-6.97	103.12	108.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3529	0	3246	227	0
1	B	3523	0	3239	235	0
2	C	1615	0	1463	151	0
2	D	1615	0	1463	154	0
3	E	140	0	78	4	0
4	F	225	0	126	12	0
5	G	139	0	78	4	0
6	H	226	0	124	12	0
7	C	1	0	0	0	0
7	D	1	0	0	0	0
8	F	26	24	24	9	0
8	H	26	24	24	8	0
9	A	2	0	0	1	0
9	B	2	0	0	0	0
9	D	1	0	0	1	0
All	All	11071	48	9865	761	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 36.

The worst 5 of 761 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:70:MET:HE1	1:B:78:ASP:HB3	1.18	1.18
1:B:28:ARG:HG3	1:B:28:ARG:HH11	1.04	1.15
2:D:520:THR:HG21	2:D:622:MET:HG3	1.28	1.15
2:D:493:ILE:HG22	2:D:494:GLU:H	0.99	1.14
1:A:285:ALA:HB3	1:A:301:GLU:O	1.50	1.12

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	479/496 (97%)	412 (86%)	52 (11%)	15 (3%)	4	23
1	B	479/496 (97%)	412 (86%)	52 (11%)	15 (3%)	4	23
2	C	214/268 (80%)	187 (87%)	20 (9%)	7 (3%)	4	21
2	D	214/268 (80%)	188 (88%)	18 (8%)	8 (4%)	3	19
All	All	1386/1528 (91%)	1199 (86%)	142 (10%)	45 (3%)	4	22

5 of 45 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	52	SER
1	A	395	SER
1	B	395	SER
2	C	493	ILE
2	C	572	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	327/431 (76%)	300 (92%)	27 (8%)	11	38
1	B	326/431 (76%)	297 (91%)	29 (9%)	9	34
2	C	141/224 (63%)	122 (86%)	19 (14%)	4	16
2	D	141/224 (63%)	121 (86%)	20 (14%)	3	14
All	All	935/1310 (71%)	840 (90%)	95 (10%)	7	27

5 of 95 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	429	LEU
2	C	608	ILE
2	C	475	GLU
2	C	552	LYS
2	D	416	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
2	C	450	GLN
1	B	461	ASN
1	B	91	ASN
1	B	352	HIS
1	B	68	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
8	PDQ	H	0	-	26,29,29	5.35	17 (65%)	30,45,45	1.82	5 (16%)
8	PDQ	F	0	-	26,29,29	5.33	16 (61%)	30,45,45	1.86	6 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	PDQ	H	0	-	-	2/11/23/23	0/4/4/4
8	PDQ	F	0	-	-	2/11/23/23	0/4/4/4

The worst 5 of 33 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	H	0	PDQ	C12-C10	14.64	1.64	1.40
8	F	0	PDQ	C12-C10	14.34	1.63	1.40
8	F	0	PDQ	C4-C3	13.35	1.60	1.39
8	H	0	PDQ	C4-C3	13.26	1.60	1.39
8	F	0	PDQ	C12-N13	11.62	1.62	1.42

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	F	0	PDQ	C20-C18-N13	-5.02	113.23	118.29
8	H	0	PDQ	C18-N13-C15	4.82	124.48	117.33
8	F	0	PDQ	C18-N13-C15	4.40	123.86	117.33
8	H	0	PDQ	C20-C18-N13	-4.20	114.05	118.29
8	H	0	PDQ	C1-N17-C15	-2.98	121.35	126.61

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	F	0	PDQ	C20-C18-N13-C15
8	F	0	PDQ	C23-C18-N13-C15
8	H	0	PDQ	C20-C18-N13-C15
8	H	0	PDQ	C23-C18-N13-C15

There are no ring outliers.


2 monomers are involved in 17 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	H	0	PDQ	8	0
8	F	0	PDQ	9	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is



within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	481/496 (96%)	-0.44	4 (0%) 86 72	67, 106, 160, 242	0
1	B	481/496 (96%)	-0.46	1 (0%) 95 90	68, 106, 158, 241	0
2	C	222/268 (82%)	-0.53	0 100 100	75, 116, 152, 182	0
2	D	222/268 (82%)	-0.52	0 100 100	75, 115, 153, 181	0
3	E	7/15 (46%)	-0.36	0 100 100	88, 91, 134, 170	0
4	F	11/19 (57%)	-0.68	0 100 100	96, 115, 135, 152	0
5	G	7/15 (46%)	-0.43	0 100 100	84, 90, 145, 171	0
6	H	11/19 (57%)	-0.62	0 100 100	95, 116, 124, 155	0
All	All	1442/1596 (90%)	-0.48	5 (0%) 94 88	67, 110, 158, 242	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	252	GLU	5.8
1	B	252	GLU	3.7
1	A	283	GLY	3.5
1	A	282	ALA	2.9
1	A	302	LEU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
8	PDQ	H	0	26/26	0.94	0.21	84,145,196,201	0
7	MG	D	742	1/1	0.95	0.26	81,81,81,81	0
8	PDQ	F	0	26/26	0.95	0.20	86,126,182,203	0
7	MG	C	742	1/1	0.95	0.32	81,81,81,81	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.