

Full wwPDB X-ray Structure Validation Report (i)

Aug 30, 2023 – 12:05 AM EDT

PDB ID	:	3N95
Title	:	Crystal structure of human CRFR2 alpha extracellular domain in complex
		with Urocortin 2
Authors	:	Pal, K.; Swaminathan, K.; Pioszak, A.A.; Xu, H.E.
Deposited on	:	2010-05-28
Resolution	:	2.72 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\;DIFFRACTION$

The reported resolution of this entry is 2.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motric	Whole archive	Similar resolution
IVIETIC	$(\# { m Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
R _{free}	130704	3359(2.74-2.70)
Clashscore	141614	3686 (2.74-2.70)
Ramachandran outliers	138981	3622(2.74-2.70)
Sidechain outliers	138945	3623 (2.74-2.70)
RSRZ outliers	127900	3276 (2.74-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	٨	400	.% 		
	A	482	86%	11%	•
			2%		
1	В	482	85%	12%	••
	-		2%		
1	С	482	84%	11%	•••
			11%		
1	D	482	84%	11%	·
			18%		
2	E	17	94%		6%



Mol	Chain	Length	Quality of chain				
2	F	17	6% 	12%			
3	G	2	100%				
3	Н	2	10	00%			
3	Ι	2	50%	50%			
3	J	2	50%	50%			



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 15086 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Δ	470	Total	С	Ν	Ο	S	0	0	0
	A	470	3643	2335	594	702	12	0	0	0
1	1 B	472	Total	С	Ν	0	S	0	0	0
			3659	2347	596	704	12	0		U
1	C	462	Total	С	Ν	0	S	0	0	0
			3584	2297	585	690	12	0		0
1 D	462	Total	С	Ν	0	S	0	0	0	
	403	3592	2301	586	693	12		0	0	

• Molecule 1 is a protein called Maltose binding protein-CRFR2 alpha extracellular domain.

• Molecule 2 is a protein called Urocortin-2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	Е	17	Total 122	С 72	N 28	O 22	0	0	1
2	F	17	Total 122	С 72	N 28	0 22	0	0	1

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Е	42	NH2	-	amidation	UNP Q96RP3
F	42	NH2	-	amidation	UNP Q96RP3

• Molecule 3 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace	
3	G	2	Total 23	C 12	0 11	0	0	0



0 0										
Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace				
3	Н	2	Total C O 23 12 11	0	0	0				
3	Ι	2	Total C O 23 12 11	0	0	0				
3	J	2	Total C O 23 12 11	0	0	0				

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	97	Total O 97 97	0	0
4	В	65	Total O 65 65	0	0
4	С	61	$\begin{array}{cc} \text{Total} & \text{O} \\ 61 & 61 \end{array}$	0	0
4	D	47	$\begin{array}{cc} \text{Total} & \text{O} \\ 47 & 47 \end{array}$	0	0
4	Е	1	Total O 1 1	0	0
4	F	1	Total O 1 1	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



 \bullet Molecule 1: Maltose binding protein-CRFR2 alpha extracellular domain







 \bullet Molecule 1: Maltose binding protein-CRFR2 alpha extracellular domain



Chain H:

100%



GLC1 GLC2

• Molecule 3: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain I: 50% 50%

GLC1 GLC2

• Molecule 3: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain J:	50%	50%
6103 6102		



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	54.31Å 212.07Å 107.32Å	Deperitor
a, b, c, α , β , γ	90.00° 104.54° 90.00°	Depositor
D ecolution (\hat{A})	50.00 - 2.72	Depositor
Resolution (A)	47.22 - 2.72	EDS
% Data completeness	98.9 (50.00-2.72)	Depositor
(in resolution range)	98.4 (47.22-2.72)	EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.70 (at 2.73Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
D D	0.224 , 0.266	Depositor
$\mathbf{n}, \mathbf{n}_{free}$	0.223 , 0.265	DCC
R_{free} test set	3148 reflections $(5.07%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	48.4	Xtriage
Anisotropy	0.316	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.33, 9.6	EDS
L-test for $twinning^2$	$< L > = 0.49, < L^2 > = 0.32$	Xtriage
Estimated twinning fraction	0.358 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	15086	wwPDB-VP
Average B, all atoms $(Å^2)$	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.81% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, $\rm NH2$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bo	nd lengths	Bond angles		
	Unain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.55	0/3735	0.59	1/5082~(0.0%)	
1	В	0.54	0/3751	0.60	0/5104	
1	С	0.53	1/3669~(0.0%)	0.61	3/4986~(0.1%)	
1	D	0.51	0/3677	0.59	0/4997	
2	Е	0.40	0/120	0.53	0/161	
2	F	0.44	0/120	0.56	0/161	
All	All	0.53	1/15072~(0.0%)	0.60	4/20491~(0.0%)	

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
1	С	-92	GLU	CB-CG	-6.32	1.40	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	С	-92	GLU	N-CA-CB	-5.56	100.59	110.60
1	С	-92	GLU	CA-CB-CG	5.56	125.63	113.40
1	С	38	SER	N-CA-C	5.27	125.24	111.00
1	А	31	LEU	CA-CB-CG	5.26	127.40	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within



3N95

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	3643	0	3559	27	1
1	В	3659	0	3581	29	0
1	С	3584	0	3515	39	0
1	D	3592	0	3519	38	1
2	Е	122	0	128	0	0
2	F	122	0	128	4	0
3	G	23	0	21	0	0
3	Н	23	0	21	0	0
3	Ι	23	0	21	0	0
3	J	23	0	21	0	0
4	А	97	0	0	5	0
4	В	65	0	0	1	0
4	С	61	0	0	11	0
4	D	47	0	0	13	0
4	Е	1	0	0	0	0
4	F	1	0	0	0	0
All	All	15086	0	14514	126	1

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (126) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
Atom-1	A00111-2	distance $(Å)$	overlap (Å)
4:A:608:HOH:O	1:C:37:TYR:HB3	1.39	1.21
1:D:-318:ALA:HB1	2:F:36:ARG:HD2	1.47	0.95
1:A:-192:ILE:HD12	1:C:38:SER:HB3	1.54	0.87
1:A:-165:ASN:HB2	4:A:215:HOH:O	1.75	0.86
1:D:-38:ASN:HB2	4:D:314:HOH:O	1.77	0.83
1:C:-298:GLN:HE21	1:C:-298:GLN:HA	1.44	0.83
1:B:-192:ILE:HD12	1:D:38:SER:HB3	1.63	0.80
1:C:-350:LEU:HA	4:C:640:HOH:O	1.81	0.79
1:D:-318:ALA:CB	2:F:36:ARG:HD2	2.12	0.79
1:D:-203:TYR:CE1	4:D:532:HOH:O	2.36	0.77
1:C:37:TYR:O	1:C:37:TYR:CD2	2.39	0.76
1:A:-302:GLY:HA3	1:A:-38:ASN:O	1.91	0.70
1:D:-298:GLN:HE21	1:D:-298:GLN:HA	1.56	0.70
1:D:-199:TYR:OH	1:D:-196:GLY:HA2	1.92	0.69
1:B:-302:GLY:HA3	1:B:-38:ASN:O	1.93	0.68
1:C:101:ILE:HG23	4:C:202:HOH:O	1.94	0.68



		Interatomic	Clash		
Atom-1	Atom-2	distance (\AA)	overlap (Å)		
1:C:37:TYR:O	4:C:536:HOH:O	2.11	0.67		
1:C:-298:GLN:HE21	1:C:-298:GLN:CA	2.08	0.66		
1:B:86:ASN:HB2	1:D:-198:GLU:CD	2.16	0.66		
1:C:-302:GLY:HA3	1:C:-38:ASN:O	1.97	0.65		
1:C:-38:ASN:HB2	4:C:425:HOH:O	1.96	0.65		
1:D:-277:THR:OG1	4:D:729:HOH:O	2.13	0.65		
1:D:-285:PHE:HD1	4:D:535:HOH:O	1.80	0.64		
1:D:-302:GLY:HA3	1:D:-38:ASN:O	1.98	0.64		
1:A:-242:THR:OG1	1:A:-239:GLU:HG2	1.99	0.62		
1:D:-242:THR:OG1	1:D:-239:GLU:HG3	1.99	0.62		
1:C:-152:ASN:HD21	1:C:-135:ILE:HG12	1.65	0.62		
1:C:-199:TYR:OH	1:C:-196:GLY:HA2	2.00	0.61		
1:D:-203:TYR:CD1	4:D:532:HOH:O	2.53	0.60		
1:B:86:ASN:HB2	1:D:-198:GLU:OE2	2.01	0.59		
1:D:-152:ASN:HD21	1:D:-135:ILE:HG12	1.67	0.59		
1:B:-199:TYR:OH	1:B:-196:GLY:HA2	2.04	0.58		
1:D:-190:ASP:O	4:D:532:HOH:O	2.17	0.58		
1:D:52:PRO:HD3	1:D:62:ARG:HD3	1.86	0.57		
1:D:-281:LEU:HD12	1:D:-276:TRP:CZ2	2.38	0.57		
1:B:-242:THR:OG1	1:B:-239:GLU:HG2	2.05	0.57		
1:C:37:TYR:HA	4:C:536:HOH:O	2.03	0.56		
1:A:51:TRP:CZ3	1:A:62:ARG:HG2	2.40	0.56		
1:C:91:SER:HB3	4:C:406:HOH:O	2.05	0.56		
1:A:-199:TYR:OH	1:A:-196:GLY:HA2	2.07	0.55		
1:A:52:PRO:HD3	1:A:62:ARG:HD3	1.88	0.54		
1:A:-3:ASN:O	1:A:1:GLU:HG3	2.07	0.54		
1:B:51:TRP:CZ3	1:B:62:ARG:HG2	2.43	0.54		
1:A:-368:ILE:HG12	1:A:-314:GLY:O	2.08	0.54		
1:B:-152:ASN:HD21	1:B:-135:ILE:HG12	1.73	0.53		
1:B:52:PRO:HD3	1:B:62:ARG:HD3	1.90	0.53		
1:D:51:TRP:CZ3	1:D:62:ARG:HG2	2.44	0.53		
1:C:51:TRP:CZ3	1:C:62:ARG:HG2	2.44	0.52		
1:B:86:ASN:HB2	1:D:-198:GLU:OE1	2.09	0.52		
1:C:-289:PRO:O	4:C:438:HOH:O	2.19	0.52		
1:A:31:LEU:O	1:A:33:PRO:HD3	2.10	0.52		
2:F:36:ARG:HB3	2:F:40:ARG:HH12	1.75	0.52		
1:C:-298:GLN:HA	1:C:-298:GLN:NE2	2.21	0.51		
1:C:-248:LEU:HD21	1:C:-235:LEU:HD11	1.92	0.51		
1:C:-242:THR:OG1	1:C:-239:GLU:HG3	2.11	0.51		
1:A:-150:GLY:HA2	4:A:130:HOH:O	2.10	0.51		
1:A:7:HIS:NE2	1:A:11:GLU:OE1	2.45	0.50		



		Interatomic	Clash		
Atom-1	Atom-2	distance (Å)	overlap (Å)		
1:B:-248:LEU:HD21	1:B:-235:LEU:HD11	1.93	0.50		
1:C:-350:LEU:HD12	4:C:640:HOH:O	2.10	0.50		
1:A:-336:LYS:HE2	4:A:282:HOH:O	2.10	0.50		
1:D:-358:ASN:ND2	1:D:-356:ASP:OD1	2.45	0.50		
1:D:-318:ALA:HB1	2:F:36:ARG:CD	2.33	0.49		
1:B:-155:ALA:O	1:B:-151:LYS:HB2	2.12	0.49		
1:A:-192:ILE:HD12	1:C:38:SER:CB	2.34	0.49		
1:D:-302:GLY:N	4:D:314:HOH:O	2.46	0.49		
1:A:-326:GLU:OE1	1:A:-326:GLU:N	2.44	0.48		
1:C:86:ASN:ND2	4:C:113:HOH:O	2.46	0.48		
1:A:-248:LEU:HD21	1:A:-235:LEU:HD11	1.96	0.48		
1:B:-355:LYS:HE3	1:B:-259:GLU:OE2	2.14	0.48		
1:B:38:SER:HA	4:B:305:HOH:O	2.14	0.48		
1:D:-302:GLY:CA	4:D:314:HOH:O	2.62	0.48		
1:A:-152:ASN:HD21	1:A:-135:ILE:HG12	1.79	0.47		
1:B:-57:LYS:NZ	1:C:-337:ILE:HD11	2.29	0.47		
1:C:52:PRO:HD3	1:C:62:ARG:HD3	1.97	0.47		
1:B:-298:GLN:OE1	1:D:88:THR:HG21	2.14	0.47		
1:A:32:ASP:HB2	4:A:117:HOH:O	2.14	0.47		
1:B:-326:GLU:OE1	1:B:-326:GLU:N	2.45	0.47		
1:B:7:HIS:NE2	1:B:11:GLU:OE1	2.49	0.46		
1:C:-358:ASN:ND2	1:C:-356:ASP:OD1	2.49	0.46		
1:D:-248:LEU:HD21	1:D:-235:LEU:HD11	1.97	0.46		
1:C:-116:PRO:HB3	1:C:-44:LYS:HD3	1.98	0.46		
1:B:-298:GLN:NE2	1:B:-298:GLN:HA	2.31	0.46		
1:C:-355:LYS:HE3	1:C:-259:GLU:OE2	2.16	0.45		
1:D:-289:PRO:HB2	4:D:535:HOH:O	2.16	0.45		
1:D:40:CYS:HB3	1:D:89:TRP:CE2	2.51	0.45		
1:C:-155:ALA:O	1:C:-151:LYS:HB2	2.17	0.45		
1:D:-308:TRP:HB3	1:D:-303:PHE:HE1	1.81	0.45		
1:D:-56:ASP:HA	1:D:-55:PRO:HD2	1.88	0.45		
1:A:-355:LYS:HE3	1:A:-259:GLU:OE2	2.17	0.45		
1:C:-298:GLN:CA	1:C:-298:GLN:NE2	2.78	0.45		
1:D:-26:ARG:CZ	4:D:394:HOH:O	2.64	0.45		
1:B:21:GLU:HG3	1:B:53:ARG:HB2	1.98	0.45		
1:C:-281:LEU:HD12	1:C:-276:TRP:CZ2	2.51	0.45		
1:A:-241:TRP:HA	1:A:-238:ILE:HD12	1.98	0.44		
1:D:-203:TYR:HE1	4:D:532:HOH:O	1.86	0.44		
1:C:62:ARG:NH2	4:C:452:HOH:O	2.51	0.44		
1:D:-328:LYS:HE2	4:D:385:HOH:O	2.16	0.44		
1:D:-273:VAL:O	1:D:-266:ILE:HG13	2.16	0.43		



		Interatomic	Clash
Atom-1	Atom-2	distance (\AA)	overlap (Å)
1:B:-358:ASN:ND2	1:B:-356:ASP:OD1	2.48	0.43
1:A:-141:PRO:HA	1:A:-138:TRP:CE2	2.53	0.43
1:C:-298:GLN:HG2	1:C:-271:TYR:OH	2.19	0.43
1:B:-368:ILE:HG12	1:B:-314:GLY:O	2.19	0.43
1:A:-116:PRO:HB3	1:A:-44:LYS:HD3	2.00	0.43
1:D:-116:PRO:HB3	1:D:-44:LYS:HD3	2.01	0.43
1:B:-116:PRO:HB3	1:B:-44:LYS:HD3	2.00	0.43
1:A:61:GLU:O	1:A:62:ARG:HD2	2.20	0.42
1:C:-273:VAL:O	1:C:-266:ILE:HG13	2.20	0.42
1:B:64:CYS:O	1:B:75:THR:HG22	2.20	0.42
1:C:-56:ASP:HA	1:C:-55:PRO:HD2	1.92	0.42
1:B:-241:TRP:HA	1:B:-238:ILE:HD12	2.00	0.42
1:C:37:TYR:C	4:C:536:HOH:O	2.55	0.42
1:A:64:CYS:O	1:A:75:THR:HG22	2.20	0.41
1:D:-189:VAL:HA	4:D:532:HOH:O	2.19	0.41
1:D:-298:GLN:HA	1:D:-298:GLN:NE2	2.32	0.41
1:A:-358:ASN:ND2	1:A:-356:ASP:OD1	2.50	0.41
1:B:-309:PHE:HA	1:B:-107:SER:O	2.20	0.41
1:C:-141:PRO:HA	1:C:-138:TRP:CE2	2.55	0.41
1:A:51:TRP:CE3	1:A:62:ARG:HG2	2.56	0.41
1:B:-141:PRO:HA	1:B:-138:TRP:CE2	2.56	0.41
1:C:-266:ILE:HG13	1:C:-266:ILE:H	1.78	0.41
1:D:-355:LYS:HE3	1:D:-259:GLU:OE2	2.21	0.41
1:C:37:TYR:O	1:C:37:TYR:HD2	1.97	0.40
1:B:-266:ILE:H	1:B:-266:ILE:HG13	1.77	0.40
1:B:-49:MET:HA	1:B:-49:MET:CE	2.51	0.40
1:A:-266:ILE:H	1:A:-266:ILE:HG13	1.74	0.40
1:C:18:LEU:HG	1:C:22:LEU:HD22	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:-55:PRO:CD	1:D:-340:ASP:O[1_455]	2.12	0.08



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	468/482~(97%)	460 (98%)	7 (2%)	1 (0%)	47	72
1	В	470/482~(98%)	461 (98%)	8 (2%)	1 (0%)	47	72
1	С	458/482~(95%)	449 (98%)	8 (2%)	1 (0%)	47	72
1	D	459/482~(95%)	448 (98%)	10 (2%)	1 (0%)	47	72
2	Е	15/17~(88%)	14 (93%)	1 (7%)	0	100	100
2	F	15/17~(88%)	15 (100%)	0	0	100	100
All	All	1885/1962~(96%)	1847 (98%)	34 (2%)	4 (0%)	47	72

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	С	-202	ALA
1	А	-202	ALA
1	D	-202	ALA
1	В	-202	ALA

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	nalysed Rotameric Outliers		Percentiles		
1	А	377/388~(97%)	363~(96%)	14 (4%)	34 61		
1	В	379/388~(98%)	357 (94%)	22~(6%)	20 42		
1	С	371/388~(96%)	354 (95%)	17 (5%)	27 52		

Continued on next page...



Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	D	372/388~(96%)	357~(96%)	15~(4%)	31	58
2	Е	11/11 (100%)	10 (91%)	1 (9%)	9	21
2	F	11/11~(100%)	11 (100%)	0	100	100
All	All	1521/1574~(97%)	1452~(96%)	69~(4%)	27	53

Continued from previous page...

All (69) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	-345	LYS
1	А	-327	LEU
1	А	-310	ILE
1	А	-287	LYS
1	А	-270	ASN
1	А	-225	SER
1	А	-187	VAL
1	А	-137	SER
1	А	-112	PHE
1	А	-41	ILE
1	А	-35	GLN
1	А	-8	LYS
1	А	31	LEU
1	А	43	THR
1	В	-364	LYS
1	В	-345	LYS
1	В	-327	LEU
1	В	-310	ILE
1	В	-298	GLN
1	В	-287	LYS
1	В	-270	ASN
1	В	-266	ILE
1	В	-239	GLU
1	В	-225	SER
1	В	-187	VAL
1	В	-175	LEU
1	В	-159	SER
1	В	-137	SER
1	В	-112	PHE
1	В	-41	ILE
1	В	-35	GLN
1	В	-8	LYS
1	В	34	GLU



Mol	Chain	Res	Type
1	В	43	THR
1	В	61	GLU
1	В	101	ILE
1	С	-327	LEU
1	C	-310	ILE
1	С	-298	GLN
1	С	-287	LYS
1	С	-270	ASN
1	С	-266	ILE
1	С	-187	VAL
1	С	-175	LEU
1	С	-131	LYS
1	С	-112	PHE
1	С	-85	LEU
1	С	-41	ILE
1	С	-35	GLN
1	С	22	LEU
1	С	24	LEU
1	С	43	THR
1	С	61	GLU
1	D	-327	LEU
1	D	-310	ILE
1	D	-298	GLN
1	D	-287	LYS
1	D	-270	ASN
1	D	-266	ILE
1	D	-187	VAL
1	D	-131	LYS
1	D	-112	PHE
1	D	-41	ILE
1	D	-35	GLN
1	D	-8	LYS
1	D	24	LEU
1	D	43	THR
1	D	61	GLU
2	Е	36	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	-352	ASN
1	А	-321	GLN



Mol	Chain	\mathbf{Res}	Type
1	А	-152	ASN
1	А	-35	GLN
1	В	-352	ASN
1	В	-169	ASN
1	В	-152	ASN
1	В	-35	GLN
1	С	-352	ASN
1	С	-298	GLN
1	С	-169	ASN
1	С	-152	ASN
1	С	-35	GLN
1	D	-352	ASN
1	D	-298	GLN
1	D	-169	ASN
1	D	-152	ASN
1	D	-35	GLN
1	D	13	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Dog	Tink	Bo	ond leng	ths	B	ond ang	les
WIOI	Type	Ullalli	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GLC	G	1	3	12,12,12	0.78	0	17,17,17	1.02	1 (5%)



Mal	Turne	Chain	Dec	Tink	Bo	ond leng	$_{\rm ths}$	В	ond ang	gles
WIOI	туре	Unain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GLC	G	2	3	11,11,12	0.65	0	$15,\!15,\!17$	1.27	2 (13%)
3	GLC	Н	1	3	12,12,12	0.55	0	$17,\!17,\!17$	1.30	3 (17%)
3	GLC	Н	2	3	11,11,12	0.77	0	$15,\!15,\!17$	1.06	1 (6%)
3	GLC	Ι	1	3	12,12,12	0.56	0	$17,\!17,\!17$	0.80	0
3	GLC	Ι	2	3	11,11,12	0.82	0	$15,\!15,\!17$	1.09	1 (6%)
3	GLC	J	1	3	12,12,12	0.53	0	$17,\!17,\!17$	0.85	0
3	GLC	J	2	3	11,11,12	0.75	0	$15,\!15,\!17$	0.97	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLC	G	1	3	-	2/2/22/22	0/1/1/1
3	GLC	G	2	3	-	0/2/19/22	0/1/1/1
3	GLC	Н	1	3	-	0/2/22/22	0/1/1/1
3	GLC	Н	2	3	-	0/2/19/22	0/1/1/1
3	GLC	Ι	1	3	-	2/2/22/22	0/1/1/1
3	GLC	Ι	2	3	-	2/2/19/22	0/1/1/1
3	GLC	J	1	3	-	2/2/22/22	0/1/1/1
3	GLC	J	2	3	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

Mol	Chain	Res	Type	Atoms		$Observed(^{o})$	$Ideal(^{o})$
3	G	2	GLC	C1-O5-C5	3.05	116.33	112.19
3	Ι	2	GLC	C1-O5-C5	3.05	116.32	112.19
3	Н	1	GLC	O5-C1-C2	3.04	115.72	110.28
3	Н	2	GLC	C1-O5-C5	2.35	115.38	112.19
3	G	2	GLC	C2-C3-C4	-2.09	107.27	110.89
3	Н	1	GLC	C1-C2-C3	2.05	114.56	110.31
3	J	2	GLC	C1-O5-C5	2.03	114.94	112.19
3	G	1	GLC	O5-C5-C6	-2.02	101.42	106.44
3	Н	1	GLC	O2-C2-C3	-2.01	105.69	110.35

All (9) bond angle outliers are listed below:

There are no chirality outliers.

All (8) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	J	1	GLC	C4-C5-C6-O6
3	J	1	GLC	O5-C5-C6-O6
3	G	1	GLC	O5-C5-C6-O6
3	Ι	1	GLC	C4-C5-C6-O6
3	G	1	GLC	C4-C5-C6-O6
3	Ι	1	GLC	O5-C5-C6-O6
3	Ι	2	GLC	C4-C5-C6-O6
3	Ι	2	GLC	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









3N95





5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	470/482~(97%)	0.06	4 (0%) 84 85	16, 25, 26, 35	0
1	В	472/482~(97%)	0.15	10 (2%) 63 65	14, 25, 26, 33	0
1	С	462/482~(95%)	0.30	9 (1%) 66 69	9, 25, 26, 31	0
1	D	463/482~(96%)	0.53	51 (11%) 5 4	12, 25, 26, 37	0
2	Ε	16/17~(94%)	0.83	3(18%) 1 1	2, 14, 16, 18	0
2	F	16/17~(94%)	0.39	1 (6%) 20 19	9, 16, 27, 27	0
All	All	1899/1962~(96%)	0.27	78 (4%) 37 36	2, 25, 26, 37	0

All (78) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	-337	ILE	6.0
1	D	-365	GLY	5.6
1	D	-88	ASN	5.5
1	D	-87	TYR	5.4
1	В	34	GLU	5.1
1	D	-94	ALA	5.0
1	D	-91	PHE	4.7
1	В	102	LEU	4.7
1	D	-95	LEU	4.6
1	D	-343	PHE	4.1
1	D	-368	ILE	3.9
1	D	-363	LEU	3.8
1	D	-92	GLU	3.7
1	D	-85	LEU	3.7
1	D	-339	THR	3.7
1	D	-67	ALA	3.7
1	D	-285	PHE	3.6
1	D	-90	LEU	3.6
1	D	-318	ALA	3.6



Continued from previous page...MolChainResTypeRSRZ

1	D	-367	GLU	3.6
1	D	-86	LEU	3.6
1	D	-364	LYS	3.5
1	D	102	LEU	3.5
1	В	67	TYR	3.4
1	D	-96	GLU	3.4
1	С	-318	ALA	3.3
1	В	-197	ASN	3.3
1	D	-104	ILE	3.3
1	С	-365	GLY	3.2
1	D	-278	PHE	3.1
1	D	-291	ILE	3.1
1	D	-341	LYS	3.1
2	Е	27	ALA	3.1
1	D	-89	GLU	3.1
1	D	-265	ALA	3.0
1	D	-338	GLY	3.0
1	D	-100	SER	3.0
2	Е	29	GLU	2.9
1	D	-63	TYR	2.9
1	D	37	TYR	2.9
1	В	70	GLY	2.8
1	D	-264	TYR	2.8
1	D	-362	VAL	2.8
1	D	-289	PRO	2.8
1	А	67	TYR	2.8
1	В	100	PRO	2.7
1	С	-61	GLU	2.7
1	D	-61	GLU	2.7
1	D	-320	VAL	2.7
1	D	-292	GLU	2.7
1	D	-286	ALA	2.6
1	С	37	TYR	2.6
1	D	-279	PRO	2.6
1	С	-102	ALA	2.6
1	D	-66	LEU	2.5
1	D	-366	GLU	2.5
1	D	-336	LYS	2.5
1	D	-361	ILE	2.4
2	Е	26	ALA	2.4
1	D	-266	ILE	2.4
1	В	85	GLU	2.4



Mol	Mol Chain		Type	RSRZ	
1	D	-346	GLY	2.3	
1	С	-279	PRO	2.3	
1	А	34	GLU	2.2	
1	С	-100	SER	2.2	
1	D	-280	TYR	2.2	
1	В	73	TYR	2.2	
1	В	32	ASP	2.2	
1	С	-45	GLN	2.1	
1	А	100	PRO	2.1	
1	D	-283	ASP	2.1	
1	В	-369	LYS	2.1	
1	D	26	GLY	2.1	
2	F	40	ARG	2.0	
1	D	-316	GLY	2.0	
1	D	85	GLU	2.0	
1	A	59	LEU	2.0	
1	С	-289	PRO	2.0	

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} extsf{-}\mathbf{B} extsf{-}\mathbf{factors}(\mathbf{A}^2)$	Q<0.9
3	GLC	Ι	2	11/12	0.94	0.16	38,40,41,44	0
3	GLC	Н	1	12/12	0.95	0.14	31,32,33,36	0
3	GLC	G	1	12/12	0.95	0.18	30,32,33,36	0
3	GLC	J	2	11/12	0.95	0.18	44,45,46,47	0
3	GLC	J	1	12/12	0.96	0.17	35,44,45,47	0
3	GLC	Ι	1	12/12	0.96	0.16	38,41,43,44	0
3	GLC	Н	2	11/12	0.97	0.15	27,28,30,30	0
3	GLC	G	2	11/12	0.98	0.15	26,27,29,29	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

















6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.

