



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2024 – 12:15 AM EST

PDB ID : 3NM3
Title : The Crystal Structure of *Candida glabrata* THI6, a Bifunctional Enzyme involved in Thiamin Biosynthesis of Eukaryotes
Authors : Paul, D.; Chatterjee, A.; Begley, T.P.; Ealick, S.E.
Deposited on : 2010-06-21
Resolution : 3.10 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RSRZ outliers	127900	1067 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	540	 57% 32% 5% 6%
1	B	540	 55% 32% 7% 6%
1	C	540	 57% 31% 6% 6%
1	D	540	 57% 34% 5% 4%
1	E	540	 56% 32% 5% 6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	540	 53% 35% 6% 6%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 22991 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Thiamine biosynthetic bifunctional enzyme.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	510	3809	2408	643	736	22	0	0	0
1	B	508	3799	2402	641	734	22	0	0	0
1	C	510	3803	2403	642	736	22	0	0	0
1	D	511	3812	2409	644	737	22	0	0	0
1	E	507	3792	2397	640	733	22	0	0	0
1	F	507	3784	2391	638	733	22	0	0	0

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Mg 1	0	0
2	B	1	Total 1	Mg 1	0	0
2	C	1	Total 1	Mg 1	0	0
2	D	1	Total 1	Mg 1	0	0
2	E	1	Total 1	Mg 1	0	0
2	F	1	Total 1	Mg 1	0	0

- Molecule 3 is THIAMIN PHOSPHATE (three-letter code: TPS) (formula: C₁₂H₁₈N₄O₄PS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			S
3	A	1	Total	C	N	O	P	S	0	0
			22	12	4	4	1	1		
3	B	1	Total	C	N	O	P	S	0	0
			22	12	4	4	1	1		
3	C	1	Total	C	N	O	P	S	0	0
			22	12	4	4	1	1		
3	D	1	Total	C	N	O	P	S	0	0
			22	12	4	4	1	1		
3	E	1	Total	C	N	O	P	S	0	0
			22	12	4	4	1	1		
3	F	1	Total	C	N	O	P	S	0	0
			22	12	4	4	1	1		

- Molecule 4 is PYROPHOSPHATE 2- (three-letter code: POP) (formula: H₂O₇P₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			9	7	2		
4	B	1	Total	O	P	0	0
			9	7	2		
4	C	1	Total	O	P	1	0
			9	7	2		
4	D	1	Total	O	P	0	0
			9	7	2		
4	E	1	Total	O	P	1	0
			9	7	2		
4	F	1	Total	O	P	0	0
			9	7	2		


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Thiamine biosynthetic bifunctional enzyme




- Molecule 1: Thiamine biosynthetic bifunctional enzyme







• Molecule 1: Thiamine biosynthetic bifunctional enzyme




• Molecule 1: Thiamine biosynthetic bifunctional enzyme






• Molecule 1: Thiamine biosynthetic bifunctional enzyme


Chain E: 56% 32% 5% 6%



• Molecule 1: Thiamine biosynthetic bifunctional enzyme

Chain F: 53% 35% 6% 6%





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	105.19Å 154.60Å 148.65Å 90.00° 102.10° 90.00°	Depositor
Resolution (Å)	46.00 – 3.10 47.87 – 3.10	Depositor EDS
% Data completeness (in resolution range)	80.6 (46.00-3.10) 59.4 (47.87-3.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.62 (at 3.12Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.197 , 0.251 0.194 , 0.194	Depositor DCC
R_{free} test set	2678 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	69.7	Xtrriage
Anisotropy	0.534	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 52.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	22991	wwPDB-VP
Average B, all atoms (Å ²)	84.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, POP, TPS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.37	0/3869	0.57	1/5246 (0.0%)
1	B	0.37	0/3859	0.57	1/5232 (0.0%)
1	C	0.37	0/3861	0.55	0/5235
1	D	0.38	0/3870	0.56	0/5246
1	E	0.37	0/3851	0.55	0/5221
1	F	0.36	0/3843	0.57	0/5213
All	All	0.37	0/23153	0.56	2/31393 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	153	PRO	CA-N-CD	-8.58	99.49	111.50
1	B	153	PRO	CA-N-CD	-8.57	99.51	111.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3809	0	3814	163	0
1	B	3799	0	3810	166	0
1	C	3803	0	3796	152	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	3812	0	3809	156	0
1	E	3792	0	3802	162	0
1	F	3784	0	3780	173	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	22	0	16	4	0
3	B	22	0	16	3	0
3	C	22	0	16	0	0
3	D	22	0	16	1	0
3	E	22	0	16	4	0
3	F	22	0	16	4	0
4	A	9	0	0	2	0
4	B	9	0	0	1	0
4	C	9	0	0	2	0
4	D	9	0	0	1	0
4	E	9	0	0	0	0
4	F	9	0	0	1	0
All	All	22991	0	22907	962	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 962 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:209:VAL:HG11	1:F:213:ILE:HG12	1.27	1.13
1:A:167:GLU:HG2	1:A:198:ASN:HD21	1.14	1.12
1:F:208:VAL:HB	1:F:209:VAL:HA	1.30	1.08
1:C:266:LEU:HD11	1:C:291:ILE:HG13	1.43	1.01
1:B:78:ILE:HG13	1:B:89:ILE:HD12	1.45	0.96

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	500/540 (93%)	438 (88%)	53 (11%)	9 (2%)	8	34
1	B	498/540 (92%)	446 (90%)	39 (8%)	13 (3%)	5	26
1	C	499/540 (92%)	446 (89%)	44 (9%)	9 (2%)	8	34
1	D	499/540 (92%)	448 (90%)	41 (8%)	10 (2%)	7	31
1	E	497/540 (92%)	440 (88%)	47 (10%)	10 (2%)	7	31
1	F	497/540 (92%)	427 (86%)	55 (11%)	15 (3%)	4	23
All	All	2990/3240 (92%)	2645 (88%)	279 (9%)	66 (2%)	6	29

5 of 66 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	11	SER
1	A	138	GLY
1	B	20	MET
1	B	434	SER
1	D	39	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	413/449 (92%)	374 (91%)	39 (9%)	8	32
1	B	413/449 (92%)	368 (89%)	45 (11%)	6	25
1	C	411/449 (92%)	371 (90%)	40 (10%)	8	30

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	412/449 (92%)	369 (90%)	43 (10%)	7	27
1	E	412/449 (92%)	371 (90%)	41 (10%)	7	28
1	F	410/449 (91%)	362 (88%)	48 (12%)	5	22
All	All	2471/2694 (92%)	2215 (90%)	256 (10%)	7	27

5 of 256 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	272	ASN
1	F	299	VAL
1	C	187	ILE
1	C	167	GLU
1	F	322	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 72 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	275	HIS
1	F	484	ASN
1	E	281	ASN
1	F	272	ASN
1	B	355	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 18 ligands modelled in this entry, 6 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	POP	F	4006	2	6,8,8	0.61	0	13,13,13	1.88	4 (30%)
4	POP	D	4004	2	6,8,8	0.75	0	13,13,13	1.51	2 (15%)
4	POP	A	4001	2	6,8,8	0.53	0	13,13,13	1.64	2 (15%)
3	TPS	A	2001	-	20,23,23	6.43	11 (55%)	25,33,33	2.19	11 (44%)
3	TPS	E	2005	-	20,23,23	6.33	12 (60%)	25,33,33	1.99	7 (28%)
3	TPS	F	2006	-	20,23,23	6.56	11 (55%)	25,33,33	2.01	8 (32%)
4	POP	C	4003	2	6,8,8	0.69	0	13,13,13	1.69	1 (7%)
3	TPS	C	2003	-	20,23,23	6.79	12 (60%)	25,33,33	2.20	11 (44%)
3	TPS	D	2004	-	20,23,23	6.52	10 (50%)	25,33,33	2.16	9 (36%)
4	POP	B	4002	2	6,8,8	0.62	0	13,13,13	1.58	2 (15%)
4	POP	E	4005	2	6,8,8	0.79	0	13,13,13	1.49	2 (15%)
3	TPS	B	2002	-	20,23,23	6.06	11 (55%)	25,33,33	1.63	5 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	POP	F	4006	2	-	4/6/6/6	-
4	POP	D	4004	2	-	0/6/6/6	-
4	POP	A	4001	2	-	2/6/6/6	-
3	TPS	A	2001	-	-	5/10/11/11	0/2/2/2
3	TPS	E	2005	-	-	5/10/11/11	0/2/2/2
3	TPS	F	2006	-	-	7/10/11/11	0/2/2/2
4	POP	C	4003	2	-	1/6/6/6	-
3	TPS	C	2003	-	-	2/10/11/11	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TPS	D	2004	-	-	3/10/11/11	0/2/2/2
4	POP	B	4002	2	-	3/6/6/6	-
4	POP	E	4005	2	-	2/6/6/6	-
3	TPS	B	2002	-	-	3/10/11/11	0/2/2/2

The worst 5 of 67 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	2003	TPS	C4-N3	26.19	1.62	1.39
3	F	2006	TPS	C4-N3	24.04	1.60	1.39
3	D	2004	TPS	C4-N3	23.50	1.60	1.39
3	A	2001	TPS	C4-N3	23.27	1.59	1.39
3	E	2005	TPS	C4-N3	23.00	1.59	1.39

The worst 5 of 64 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	2001	TPS	C6-C5-C4	-5.33	123.16	127.43
4	C	4003	POP	P2-O-P1	-5.16	115.12	132.83
3	D	2004	TPS	C6-C5-C4	-4.92	123.48	127.43
4	A	4001	POP	P2-O-P1	-4.72	116.62	132.83
3	D	2004	TPS	C6A-C5A-C4A	4.56	121.93	115.72

There are no chirality outliers.

5 of 37 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	2001	TPS	C5A-C7A-N3-C4
3	A	2001	TPS	C4-C5-C6-C7
3	A	2001	TPS	C6A-C5A-C7A-N3
3	B	2002	TPS	C5-C6-C7-O7
3	B	2002	TPS	C6A-C5A-C7A-N3

There are no ring outliers.

10 monomers are involved in 23 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	F	4006	POP	1	0
4	D	4004	POP	1	0
4	A	4001	POP	2	0



Continued on next page...



Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	2001	TPS	4	0
3	E	2005	TPS	4	0
3	F	2006	TPS	4	0
4	C	4003	POP	2	0
3	D	2004	TPS	1	0
4	B	4002	POP	1	0
3	B	2002	TPS	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	510/540 (94%)	-0.51	0 100 100	26, 86, 109, 133	0
1	B	508/540 (94%)	-0.57	1 (0%) 95 90	26, 80, 103, 129	0
1	C	510/540 (94%)	-0.42	2 (0%) 92 84	42, 87, 113, 135	0
1	D	511/540 (94%)	-0.56	0 100 100	40, 79, 107, 134	0
1	E	507/540 (93%)	-0.52	0 100 100	40, 80, 105, 129	0
1	F	507/540 (93%)	-0.42	1 (0%) 95 90	43, 87, 113, 131	0
All	All	3053/3240 (94%)	-0.50	4 (0%) 95 92	26, 83, 109, 135	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	465	GLY	2.7
1	B	455	ILE	2.2
1	F	216	LEU	2.1
1	C	247	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MG	B	3002	1/1	0.91	0.28	61,61,61,61	0
2	MG	A	3001	1/1	0.97	0.15	70,70,70,70	0
2	MG	D	3004	1/1	0.97	0.23	61,61,61,61	0
2	MG	F	3006	1/1	0.97	0.28	63,63,63,63	0
3	TPS	C	2003	22/22	0.97	0.21	66,73,76,79	7
3	TPS	E	2005	22/22	0.97	0.19	54,60,66,68	8
3	TPS	F	2006	22/22	0.97	0.19	68,73,77,83	8
4	POP	B	4002	9/9	0.97	0.20	48,57,67,70	2
4	POP	D	4004	9/9	0.97	0.17	53,60,65,68	5
3	TPS	D	2004	22/22	0.98	0.19	57,63,69,74	6
2	MG	C	3003	1/1	0.98	0.28	55,55,55,55	0
3	TPS	A	2001	22/22	0.98	0.18	60,69,76,77	7
4	POP	A	4001	9/9	0.98	0.17	56,67,80,80	3
3	TPS	B	2002	22/22	0.98	0.18	55,62,70,73	8
4	POP	C	4003	9/9	0.98	0.22	55,68,75,82	3
2	MG	E	3005	1/1	0.98	0.21	48,48,48,48	0
4	POP	E	4005	9/9	0.98	0.15	57,61,63,64	6
4	POP	F	4006	9/9	0.98	0.16	55,63,69,69	4




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around TPS C 2003:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around TPS E 2005:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around TPS F 2006:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around TPS D 2004:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around TPS A 2001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around TPS B 2002:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.