



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 17, 2023 – 02:47 AM EST

PDB ID : 3NSC  
Title : C500S MUTANT OF CueO BOUND TO Cu(II)  
Authors : Roberts, S.A.; Montfort, W.R.; Singh, S.K.  
Deposited on : 2010-07-01  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	505	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4121 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Blue copper oxidase cueO.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	470	3648	2319	631	672	26	0	9	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	500	SER	CYS	engineered mutation	UNP P36649
A	517	LEU	-	expression tag	UNP P36649
A	518	GLN	-	expression tag	UNP P36649
A	519	GLY	-	expression tag	UNP P36649
A	520	ASP	-	expression tag	UNP P36649
A	521	HIS	-	expression tag	UNP P36649
A	522	GLY	-	expression tag	UNP P36649
A	523	LEU	-	expression tag	UNP P36649
A	524	SER	-	expression tag	UNP P36649
A	525	ALA	-	expression tag	UNP P36649
A	526	TRP	-	expression tag	UNP P36649
A	527	SER	-	expression tag	UNP P36649
A	528	HIS	-	expression tag	UNP P36649
A	529	PRO	-	expression tag	UNP P36649
A	530	GLN	-	expression tag	UNP P36649
A	531	PHE	-	expression tag	UNP P36649
A	532	GLU	-	expression tag	UNP P36649
A	533	LYS	-	expression tag	UNP P36649

- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Cu	0	0
			3	3		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	1
			4	2	2		


- Molecule 5 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	A	461	Total 461	O 461	0	3

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Blue copper oxidase cueO



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	50.33Å 91.57Å 54.25Å 90.00° 102.48° 90.00°	Depositor
Resolution (Å)	20.30 – 1.50 20.34 – 1.50	Depositor EDS
% Data completeness (in resolution range)	94.9 (20.30-1.50) 94.9 (20.34-1.50)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.60 (at 1.50Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.181 , 0.213 0.184 , 0.216	Depositor DCC
$R_{free}$ test set	3647 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.5	Xtrriage
Anisotropy	0.281	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 56.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4121	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.04% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, CU, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.79	1/3764 (0.0%)	0.85	4/5113 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	181	VAL	CB-CG2	-5.36	1.41	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	431	ARG	NE-CZ-NH2	5.67	123.14	120.30
1	A	40	LEU	CB-CG-CD2	-5.18	102.20	111.00
1	A	431	ARG	NE-CZ-NH1	-5.17	117.71	120.30
1	A	234	ARG	NE-CZ-NH2	-5.14	117.73	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3648	0	3641	32	0
2	A	3	0	0	0	0
3	A	5	0	0	0	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	4	0	3	0	0
5	A	461	0	0	6	0
All	All	4121	0	3644	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:106[B]:GLU:H	1:A:133:GLN:HE22	1.24	0.83
1:A:106[A]:GLU:H	1:A:133:GLN:HE22	1.24	0.83
1:A:428:GLN:HE22	1:A:520:ASP:H	1.38	0.69
1:A:450:THR:HG21	1:A:484[A]:VAL:HG23	1.80	0.64
1:A:106[A]:GLU:H	1:A:133:GLN:NE2	1.95	0.64
1:A:106[B]:GLU:H	1:A:133:GLN:NE2	1.95	0.64
1:A:355:MET:CE	1:A:408:ASN:HD21	2.14	0.61
1:A:488:HIS:HD2	5:A:681:HOH:O	1.83	0.60
1:A:355:MET:HE2	5:A:932:HOH:O	2.03	0.59
1:A:314:HIS:HE1	5:A:684:HOH:O	1.85	0.59
1:A:355:MET:HE2	1:A:408:ASN:HD21	1.69	0.57
1:A:203:MET:O	1:A:207:VAL:HG12	2.04	0.57
1:A:364:MET:O	1:A:368:MET:HG2	2.08	0.52
1:A:337:LEU:HD13	5:A:729:HOH:O	2.10	0.52
1:A:454:ILE:HG21	1:A:457:GLU:HB2	1.92	0.51
1:A:314:HIS:HD2	1:A:315:PRO:O	1.94	0.51
1:A:355:MET:CE	5:A:932:HOH:O	2.57	0.50
1:A:488:HIS:HE1	5:A:563:HOH:O	1.93	0.49
1:A:450:THR:HG21	1:A:484[A]:VAL:CG2	2.42	0.48
1:A:265:PRO:HD3	1:A:337:LEU:HD22	1.96	0.48
1:A:370:LYS:HE3	1:A:371:TYR:CZ	2.51	0.46
1:A:141:HIS:HB2	1:A:142:PRO:HD2	1.97	0.45
1:A:492:LYS:HB2	1:A:492:LYS:HE2	1.69	0.45
1:A:204[B]:THR:HG22	1:A:504:GLU:HG2	1.99	0.44
1:A:56[A]:GLN:OE1	1:A:63[A]:THR:CG2	2.65	0.43
1:A:74:LEU:HD23	1:A:74:LEU:HA	1.90	0.43
1:A:102:TRP:HB3	1:A:105:LEU:HG	1.99	0.43
1:A:56[A]:GLN:OE1	1:A:63[A]:THR:HG22	2.19	0.43
1:A:99:THR:O	1:A:142:PRO:HA	2.20	0.42
1:A:520:ASP:O	1:A:521:HIS:HD2	2.02	0.42
1:A:341:PRO:CG	1:A:483:LEU:HD21	2.50	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:181:VAL:HG22	1:A:233:LEU:HD23	2.01	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	475/505 (94%)	463 (98%)	12 (2%)	0	<a href="#">100</a> <a href="#">100</a>

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	395/414 (95%)	389 (98%)	6 (2%)	<a href="#">65</a> <a href="#">39</a>

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	143	HIS
1	A	171	MET
1	A	202	VAL
1	A	280	ARG
1	A	343	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	492	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	133	GLN
1	A	186	GLN
1	A	314	HIS
1	A	374	GLN
1	A	408	ASN
1	A	418	ASN
1	A	428	GLN
1	A	488	HIS
1	A	521	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	ACT	A	1[A]	2	3,3,3	0.81	0	3,3,3	1.67	1 (33%)
3	SO4	A	800	-	4,4,4	0.28	0	6,6,6	0.46	0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
4	A	1[A]	ACT	OXT-C-CH3	2.20	124.26	115.18

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	470/505 (93%)	0.37	34 (7%) <b>15</b> <b>16</b>	15, 23, 39, 51	1 (0%)

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	522	GLY	5.6
1	A	361	MET	4.9
1	A	520	ASP	4.5
1	A	171	MET	4.2
1	A	326	ALA	4.0
1	A	403	ASP	4.0
1	A	358	MET	3.9
1	A	343	LEU	3.7
1	A	342	SER	3.6
1	A	344	GLU	3.6
1	A	339	ALA	3.6
1	A	378	GLY	3.5
1	A	327	SER	3.5
1	A	337	LEU	3.3
1	A	345	GLY	3.3
1	A	406	HIS	3.2
1	A	30	GLU	2.8
1	A	379	MET	2.8
1	A	199	GLN	2.7
1	A	419	LYS	2.7
1	A	341	PRO	2.7
1	A	340	LEU	2.5
1	A	165	ASP	2.5
1	A	517	LEU	2.4
1	A	460	LYS	2.4
1	A	484[A]	VAL	2.4
1	A	357	PRO	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	482	VAL	2.3
1	A	405	HIS	2.2
1	A	362	MET	2.2
1	A	338	PRO	2.1
1	A	202	VAL	2.1
1	A	192	ALA	2.1
1	A	521	HIS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	SO4	A	800	5/5	0.92	0.12	23,25,29,29	5
4	ACT	A	1[A]	4/4	0.97	0.24	14,15,15,16	4
2	CU	A	604	1/1	0.98	0.05	22,22,22,22	0
2	CU	A	602	1/1	0.99	0.05	19,19,19,19	0
2	CU	A	603	1/1	1.00	0.04	19,19,19,19	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.