



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 29, 2023 – 03:12 PM EDT

PDB ID : 3O1W  
Title : Crystal structure of dimeric KIHxk1 in crystal form III  
Authors : Kuettner, E.B.; Kettner, K.; Keim, A.; Kriegel, T.M.; Strater, N.  
Deposited on : 2010-07-22  
Resolution : 1.66 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	485	
1	B	485	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NHE	A	488	-	-	X	-
4	PO4	B	490	-	X	-	-

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8575 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hexokinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	484	Total 3900	C 2497	N 636	O 748	S 19	0	26	0
1	B	484	Total 3832	C 2439	N 629	O 748	S 16	0	15	0

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 6	C 3	O 3	0	0
2	B	1	Total 6	C 3	O 3	0	0
2	B	1	Total 6	C 3	O 3	0	0

- Molecule 3 is 2-[N-CYCLOHEXYLAMINO]ETHANE SULFONIC ACID (three-letter code: NHE) (formula: C<sub>8</sub>H<sub>17</sub>NO<sub>3</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
3	A	1	Total	8	1	3	1	0	0
3	A	1	Total	8	1	3	1	0	0
3	B	1	Total	8	1	3	1	0	0
3	B	1	Total	8	1	3	1	0	0

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O P 5 4 1	0	0
4	A	1	Total O P 5 4 1	0	0
4	A	1	Total O P 5 4 1	0	0
4	B	1	Total O P 5 4 1	0	0
4	B	1	Total O P 5 4 1	0	0
4	B	1	Total O P 5 4 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	379	Total O 379 379	0	0
5	B	364	Total O 364 364	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Hexokinase



- Molecule 1: Hexokinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	60.31Å 135.84Å 72.59Å 90.00° 92.06° 90.00°	Depositor
Resolution (Å)	29.85 – 1.66 29.42 – 1.66	Depositor EDS
% Data completeness (in resolution range)	94.5 (29.85-1.66) 94.5 (29.42-1.66)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.75 (at 1.66Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.161 , 0.201 0.176 , 0.214	Depositor DCC
$R_{free}$ test set	1313 reflections (1.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.5	Xtrriage
Anisotropy	0.227	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 42.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.026 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	8575	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.41% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NHE, PO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.21	5/4046 (0.1%)	1.09	10/5473 (0.2%)
1	B	1.23	10/3954 (0.3%)	1.07	9/5353 (0.2%)
All	All	1.22	15/8000 (0.2%)	1.08	19/10826 (0.2%)

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	220	TYR	CE1-CZ	6.90	1.47	1.38
1	B	268[A]	GLU	CD-OE1	6.88	1.33	1.25
1	B	268[B]	GLU	CD-OE1	6.88	1.33	1.25
1	B	358	GLU	CB-CG	6.52	1.64	1.52
1	B	408	TYR	CE2-CZ	5.93	1.46	1.38
1	A	396	VAL	CB-CG2	5.83	1.65	1.52
1	A	158	TYR	CE2-CZ	5.83	1.46	1.38
1	B	268[A]	GLU	CB-CG	5.33	1.62	1.52
1	B	268[B]	GLU	CB-CG	5.33	1.62	1.52
1	B	158	TYR	CB-CG	-5.32	1.43	1.51
1	B	440	TRP	CZ3-CH2	5.22	1.48	1.40
1	A	480	VAL	CB-CG1	5.21	1.63	1.52
1	A	278	VAL	CB-CG1	-5.11	1.42	1.52
1	A	206	ALA	CA-CB	5.10	1.63	1.52
1	B	177	PHE	CE2-CZ	5.00	1.46	1.37

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	154	PHE	CB-CG-CD2	-8.16	115.08	120.80
1	A	416	ASP	CB-CG-OD1	7.84	125.36	118.30
1	A	359	ASP	CB-CG-OD2	-7.35	111.68	118.30
1	A	154	PHE	CB-CG-CD2	-6.86	116.00	120.80

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	422	ARG	NE-CZ-NH2	6.74	123.67	120.30
1	B	406	ARG	NE-CZ-NH1	6.56	123.58	120.30
1	A	314	ARG	NE-CZ-NH2	-6.29	117.16	120.30
1	A	277	LEU	CB-CG-CD1	-6.22	100.42	111.00
1	B	93	ARG	NE-CZ-NH1	6.21	123.40	120.30
1	A	73	TYR	CD1-CE1-CZ	-6.18	114.24	119.80
1	A	473	ARG	NE-CZ-NH2	6.04	123.32	120.30
1	B	416	ASP	CB-CG-OD1	5.92	123.63	118.30
1	B	154	PHE	CB-CG-CD1	5.82	124.87	120.80
1	A	390	ARG	NE-CZ-NH2	-5.71	117.44	120.30
1	B	408	TYR	CZ-CE2-CD2	-5.51	114.84	119.80
1	B	359	ASP	CB-CG-OD2	-5.49	113.36	118.30
1	A	436	ASP	CB-CG-OD1	5.45	123.21	118.30
1	A	48	PHE	CB-CG-CD1	-5.25	117.12	120.80
1	B	406	ARG	NE-CZ-NH2	-5.09	117.75	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3900	0	3971	50	0
1	B	3832	0	3844	19	0
2	A	6	0	8	0	0
2	B	12	0	16	1	0
3	A	26	0	33	15	0
3	B	26	0	33	0	0
4	A	15	0	0	0	0
4	B	15	0	0	0	0
5	A	379	0	0	10	0
5	B	364	0	0	4	0
All	All	8575	0	7905	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:445[A]:MET:HE1	3:A:488:NHE:C4'	1.29	1.59
1:A:445[A]:MET:CE	3:A:488:NHE:H4'1	0.96	1.44
1:A:445[A]:MET:SD	3:A:488:NHE:H3'1	1.64	1.36
1:A:445[A]:MET:CE	3:A:488:NHE:C4'	1.90	1.30
1:A:445[A]:MET:SD	3:A:488:NHE:C3'	2.28	1.20
1:A:445[A]:MET:HE3	3:A:488:NHE:H4'1	1.27	1.14
1:B:90[A]:THR:HG23	1:B:91[A]:ASN:ND2	1.69	1.07
1:A:85:LEU:HD11	1:A:92[B]:LEU:CD1	1.89	1.02
1:A:85:LEU:HD11	1:A:92[B]:LEU:HD11	1.03	1.02
1:A:85:LEU:CD1	1:A:92[B]:LEU:HD11	1.91	1.00
1:A:445[A]:MET:HE1	3:A:488:NHE:H4'2	1.50	0.94
1:A:445[A]:MET:SD	3:A:488:NHE:C4'	2.57	0.91
1:A:445[A]:MET:SD	3:A:488:NHE:H4'1	2.16	0.85
1:A:328:LYS:O	1:A:329:ASP:HB2	1.80	0.82
1:A:472:LYS:HD3	5:A:637:HOH:O	1.79	0.81
1:B:90[A]:THR:HG22	5:B:771:HOH:O	1.82	0.80
1:B:90[A]:THR:CG2	1:B:91[A]:ASN:ND2	2.47	0.77
1:A:47[B]:HIS:CE1	5:A:867:HOH:O	2.46	0.69
1:A:31:GLU:OE2	1:A:382:LYS:HE2	1.93	0.69
1:A:47[B]:HIS:HD2	5:A:626:HOH:O	1.75	0.68
1:A:59[B]:LYS:HG3	5:A:769:HOH:O	1.98	0.62
1:A:445[A]:MET:SD	3:A:488:NHE:C2'	2.88	0.61
1:B:158:TYR:CD1	1:B:158:TYR:N	2.71	0.59
1:A:445[A]:MET:HE1	3:A:488:NHE:H4'1	0.60	0.59
1:B:281:ARG:HD2	5:B:801:HOH:O	2.04	0.58
1:A:192:GLN:O	1:A:196[A]:GLU:HG2	2.04	0.58
1:A:445[A]:MET:CE	3:A:488:NHE:C3'	2.66	0.56
1:B:363:LEU:HD12	1:B:367:ASN:HD22	1.71	0.56
1:B:90[A]:THR:HG23	1:B:91[A]:ASN:CG	2.26	0.55
1:A:328:LYS:O	1:A:329:ASP:CB	2.45	0.54
1:A:136[B]:GLU:HG2	5:A:764:HOH:O	2.09	0.53
1:A:445[B]:MET:HG2	3:A:488:NHE:H2'2	1.91	0.53
1:A:90:THR:OG1	1:A:91[B]:ASN:ND2	2.42	0.52
1:A:2:VAL:N	5:A:540:HOH:O	2.42	0.51
1:A:31:GLU:HG2	1:A:382:LYS:HE2	1.93	0.51
1:A:66:ILE:HD12	1:A:268[B]:GLU:HG3	1.92	0.51
1:B:117:HIS:H	1:B:117:HIS:CD2	2.30	0.49
1:A:119:ARG:HD2	5:A:571:HOH:O	2.12	0.49
2:B:487:GOL:H11	5:B:592:HOH:O	2.13	0.49
1:B:445:MET:SD	5:B:813:HOH:O	2.60	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:23[A]:LEU:CD1	1:A:379:LEU:HD12	2.44	0.48
1:A:23[A]:LEU:HD11	1:A:379:LEU:CD1	2.44	0.47
1:B:90[A]:THR:HG23	1:B:91[A]:ASN:HD22	1.72	0.47
1:A:116:ASP:OD1	1:A:116:ASP:N	2.46	0.46
1:B:217:ALA:HB2	1:B:462:GLY:CA	2.45	0.46
1:A:427:LYS:HD3	3:A:488:NHE:HC12	1.97	0.46
1:A:195:ILE:HG23	1:A:200:ILE:HB	1.97	0.45
1:B:53:ASP:OD1	1:B:405:LYS:HE2	2.17	0.45
1:A:192:GLN:HE21	1:A:196[A]:GLU:CD	2.21	0.44
1:B:363:LEU:HD12	1:B:367:ASN:ND2	2.32	0.44
1:A:47[A]:HIS:CD2	1:A:278:VAL:HG21	2.52	0.44
1:A:31:GLU:CG	1:A:382:LYS:HE2	2.48	0.44
1:A:472:LYS:HD2	5:A:740:HOH:O	2.19	0.43
1:A:66:ILE:CD1	1:A:268[B]:GLU:HG3	2.49	0.43
1:A:335:LEU:HD12	1:A:335:LEU:HA	1.73	0.43
1:A:59[B]:LYS:HB2	5:A:658:HOH:O	2.19	0.43
1:A:445[A]:MET:SD	3:A:488:NHE:H2'2	2.58	0.43
1:A:23[A]:LEU:HD11	1:A:379:LEU:HD13	2.01	0.42
1:B:166:ASN:HB3	1:B:205:VAL:O	2.20	0.42
1:B:217:ALA:HB2	1:B:462:GLY:HA2	2.01	0.42
1:B:31:GLU:OE2	1:B:382:LYS:HE2	2.20	0.42
1:B:188:VAL:HB	1:B:189:PRO:HD3	2.02	0.42
1:B:158:TYR:CE2	1:B:175:LYS:HD2	2.55	0.41
1:A:85:LEU:CD1	1:A:92[B]:LEU:CD1	2.72	0.41
1:B:74:PRO:HD2	1:B:219:LEU:HD23	2.02	0.41
1:A:23[A]:LEU:HD22	1:A:23[A]:LEU:HA	1.94	0.40
1:A:47[B]:HIS:CD2	5:A:626:HOH:O	2.61	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	508/485 (105%)	494 (97%)	14 (3%)	0	100	100
1	B	497/485 (102%)	486 (98%)	11 (2%)	0	100	100
All	All	1005/970 (104%)	980 (98%)	25 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	437/412 (106%)	427 (98%)	10 (2%)	50	25
1	B	426/412 (103%)	420 (99%)	6 (1%)	67	46
All	All	863/824 (105%)	847 (98%)	16 (2%)	60	34

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ARG
1	A	23[A]	LEU
1	A	23[B]	LEU
1	A	48	PHE
1	A	116	ASP
1	A	162[A]	GLN
1	A	162[B]	GLN
1	A	269	TYR
1	A	333	SER
1	A	335	LEU
1	B	2	VAL
1	B	13	LYS
1	B	48	PHE
1	B	124	GLU
1	B	269	TYR
1	B	351	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	185	HIS
1	B	117	HIS
1	B	171	GLN
1	B	185	HIS
1	B	367	ASN
1	B	439	ASN
1	B	471	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GOL	B	486	-	5,5,5	0.66	0	5,5,5	0.85	0
4	PO4	A	489	-	4,4,4	1.84	1 (25%)	6,6,6	1.20	1 (16%)
2	GOL	B	487	-	5,5,5	0.56	0	5,5,5	1.02	0
4	PO4	A	490	-	4,4,4	1.13	0	6,6,6	0.82	0
3	NHE	B	489	-	13,13,13	1.78	1 (7%)	16,17,17	1.84	5 (31%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	PO4	B	490	-	4,4,4	1.51	1 (25%)	6,6,6	2.41	3 (50%)
3	NHE	A	487	-	13,13,13	1.99	1 (7%)	16,17,17	1.61	4 (25%)
4	PO4	B	492	-	4,4,4	1.19	0	6,6,6	1.69	2 (33%)
4	PO4	B	491	-	4,4,4	1.33	1 (25%)	6,6,6	1.77	2 (33%)
4	PO4	A	491	-	4,4,4	1.26	1 (25%)	6,6,6	1.68	2 (33%)
2	GOL	A	486	-	5,5,5	1.19	0	5,5,5	1.00	1 (20%)
3	NHE	B	488	-	13,13,13	1.36	1 (7%)	16,17,17	2.21	4 (25%)
3	NHE	A	488	-	13,13,13	2.10	1 (7%)	16,17,17	1.62	3 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	486	-	-	0/4/4/4	-
2	GOL	B	487	-	-	4/4/4/4	-
3	NHE	B	489	-	-	1/7/15/15	0/1/1/1
3	NHE	A	487	-	-	0/7/15/15	0/1/1/1
2	GOL	A	486	-	-	0/4/4/4	-
3	NHE	B	488	-	-	3/7/15/15	0/1/1/1
3	NHE	A	488	-	-	0/7/15/15	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	488	NHE	C2-S	-7.13	1.67	1.77
3	A	487	NHE	C2-S	-7.01	1.67	1.77
3	B	489	NHE	C2-S	-6.15	1.68	1.77
3	B	488	NHE	C2-S	-3.57	1.72	1.77
4	A	489	PO4	P-O2	3.01	1.63	1.54
4	B	490	PO4	P-O4	2.86	1.63	1.54
4	A	491	PO4	P-O4	-2.19	1.48	1.54
4	B	491	PO4	P-O3	-2.11	1.48	1.54

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	488	NHE	O1-S-C2	-6.34	99.28	106.92
3	A	488	NHE	O3-S-O2	-4.24	100.92	111.27

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	490	PO4	O4-P-O1	-4.01	96.20	110.89
3	B	488	NHE	O3-S-C2	3.95	112.16	105.77
3	B	489	NHE	O1-S-C2	-3.53	102.67	106.92
3	B	489	NHE	C6'-C1'-C2'	-3.35	105.01	110.82
4	B	491	PO4	O2-P-O1	-3.31	98.78	110.89
4	B	490	PO4	O3-P-O2	3.15	118.06	107.97
3	A	487	NHE	C1-N-C1'	3.13	120.29	114.14
4	A	491	PO4	O4-P-O1	3.01	121.93	110.89
3	B	488	NHE	C5'-C6'-C1'	-2.89	105.67	111.11
4	B	492	PO4	O3-P-O2	-2.82	98.90	107.97
4	B	492	PO4	O4-P-O1	2.79	121.12	110.89
3	A	487	NHE	O2-S-C2	2.79	110.27	106.92
3	B	489	NHE	O3-S-O1	2.78	118.07	111.27
4	B	491	PO4	O4-P-O3	2.74	116.77	107.97
3	A	487	NHE	O3-S-O2	-2.67	104.76	111.27
3	B	488	NHE	C6'-C1'-C2'	2.63	115.38	110.82
4	A	489	PO4	O3-P-O2	-2.49	99.96	107.97
3	B	489	NHE	C1-N-C1'	2.48	119.00	114.14
4	B	490	PO4	O4-P-O3	2.30	115.34	107.97
3	A	487	NHE	O1-S-C2	2.23	109.60	106.92
3	B	489	NHE	O2-S-O1	-2.23	106.25	113.95
3	A	488	NHE	C4'-C3'-C2'	-2.18	106.97	111.42
3	A	488	NHE	C4'-C5'-C6'	2.11	115.71	111.42
4	A	491	PO4	O3-P-O2	-2.07	101.32	107.97
2	A	486	GOL	C3-C2-C1	-2.06	103.68	111.70

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	487	GOL	C1-C2-C3-O3
2	B	487	GOL	O2-C2-C3-O3
3	B	488	NHE	C2-C1-N-C1'
2	B	487	GOL	O1-C1-C2-O2
3	B	489	NHE	C2-C1-N-C1'
3	B	488	NHE	C2'-C1'-N-C1
3	B	488	NHE	C6'-C1'-N-C1
2	B	487	GOL	O1-C1-C2-C3

There are no ring outliers.

2 monomers are involved in 16 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	487	GOL	1	0
3	A	488	NHE	15	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	484/485 (99%)	0.06	20 (4%) 37 37	13, 20, 38, 56	0
1	B	484/485 (99%)	0.08	16 (3%) 46 47	13, 20, 38, 52	1 (0%)
All	All	968/970 (99%)	0.07	36 (3%) 41 41	13, 20, 39, 56	1 (0%)

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	6	PRO	5.2
1	A	2	VAL	4.9
1	A	3	ARG	4.7
1	A	5	GLY	4.0
1	A	147	SER	3.5
1	B	2	VAL	3.2
1	A	4	LEU	3.2
1	A	8	LYS	3.1
1	A	59[A]	LYS	3.1
1	A	329	ASP	3.0
1	A	9	PRO	2.8
1	A	445[A]	MET	2.7
1	B	158	TYR	2.7
1	A	441	ASP	2.7
1	B	441	ASP	2.7
1	B	442	VAL	2.6
1	A	443	GLU	2.6
1	B	145	GLY	2.6
1	B	172	ARG	2.6
1	B	178	ASP	2.5
1	B	147	SER	2.5
1	B	294	ARG	2.4
1	A	152	LEU	2.4
1	A	144	ASP	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	59	LYS	2.2
1	A	22	ASN	2.2
1	A	208	ILE	2.2
1	A	85	LEU	2.1
1	B	215	LEU	2.1
1	B	144	ASP	2.1
1	B	216	VAL	2.1
1	B	177	PHE	2.1
1	A	312	ILE	2.1
1	B	6	PRO	2.0
1	B	180	GLU	2.0
1	A	143	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GOL	B	487	6/6	0.87	0.20	23,35,39,42	0
4	PO4	B	492	5/5	0.92	0.17	12,19,24,26	5
3	NHE	B	489	13/13	0.93	0.30	23,25,26,28	13
2	GOL	A	486	6/6	0.94	0.13	16,20,23,28	0
4	PO4	A	490	5/5	0.94	0.12	28,29,33,34	5
3	NHE	A	487	13/13	0.94	0.30	26,28,29,30	13
3	NHE	A	488	13/13	0.95	0.14	12,21,25,25	13
4	PO4	B	490	5/5	0.95	0.12	24,29,31,34	0
2	GOL	B	486	6/6	0.95	0.08	19,21,26,28	0
4	PO4	A	489	5/5	0.97	0.08	22,25,29,31	0
4	PO4	B	491	5/5	0.97	0.09	26,28,34,35	5

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	PO4	A	491	5/5	0.97	0.13	19,29,33,36	5
3	NHE	B	488	13/13	0.98	0.06	17,21,24,26	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.