



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 17, 2024 – 10:18 PM EDT

PDB ID : 3ON4  
Title : Crystal structure of TetR transcriptional regulator from *Legionella pneumophila*  
Authors : Michalska, K.; Li, H.; Gu, M.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)  
Deposited on : 2010-08-27  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	191	 2% 92% 7% ..
1	B	191	 % 93% 6% .
1	C	191	 2% 87% 10% ..
1	D	191	 % 89% 8% .
1	E	191	 % 93% 5% .

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	191	 <p>91% 6% •</p>
1	G	191	 <p>88% 9% •</p>
1	H	191	 <p>88% 9% ••</p>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 12866 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcriptional regulator, TetR family.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	190	1506	965	248	288	1	4	0	2	0
1	B	189	1524	977	249	293	1	4	0	5	0
1	C	187	1485	952	245	284	1	3	0	2	0
1	D	186	1498	959	247	288	1	3	0	5	0
1	E	188	1519	974	249	291	1	4	0	6	0
1	F	186	1503	961	246	292	1	3	0	5	0
1	G	187	1499	960	246	288	1	4	0	4	0
1	H	186	1477	947	241	285	1	3	0	2	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	EXPRESSION TAG	UNP Q5ZU41
A	-1	ASN	-	EXPRESSION TAG	UNP Q5ZU41
A	0	ALA	-	EXPRESSION TAG	UNP Q5ZU41
B	-2	SER	-	EXPRESSION TAG	UNP Q5ZU41
B	-1	ASN	-	EXPRESSION TAG	UNP Q5ZU41
B	0	ALA	-	EXPRESSION TAG	UNP Q5ZU41
C	-2	SER	-	EXPRESSION TAG	UNP Q5ZU41
C	-1	ASN	-	EXPRESSION TAG	UNP Q5ZU41
C	0	ALA	-	EXPRESSION TAG	UNP Q5ZU41
D	-2	SER	-	EXPRESSION TAG	UNP Q5ZU41
D	-1	ASN	-	EXPRESSION TAG	UNP Q5ZU41
D	0	ALA	-	EXPRESSION TAG	UNP Q5ZU41
E	-2	SER	-	EXPRESSION TAG	UNP Q5ZU41

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	-1	ASN	-	EXPRESSION TAG	UNP Q5ZU41
E	0	ALA	-	EXPRESSION TAG	UNP Q5ZU41
F	-2	SER	-	EXPRESSION TAG	UNP Q5ZU41
F	-1	ASN	-	EXPRESSION TAG	UNP Q5ZU41
F	0	ALA	-	EXPRESSION TAG	UNP Q5ZU41
G	-2	SER	-	EXPRESSION TAG	UNP Q5ZU41
G	-1	ASN	-	EXPRESSION TAG	UNP Q5ZU41
G	0	ALA	-	EXPRESSION TAG	UNP Q5ZU41
H	-2	SER	-	EXPRESSION TAG	UNP Q5ZU41
H	-1	ASN	-	EXPRESSION TAG	UNP Q5ZU41
H	0	ALA	-	EXPRESSION TAG	UNP Q5ZU41

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Na 1 1	0	0
2	C	1	Total Na 1 1	0	0
2	D	1	Total Na 1 1	0	0
2	E	1	Total Na 1 1	0	0

- Molecule 3 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	C O	0	0
			7	4 3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	127	Total	O	0	0
			127	127		
4	B	126	Total	O	0	0
			126	126		
4	C	100	Total	O	0	0
			100	100		
4	D	108	Total	O	0	0
			108	108		
4	E	140	Total	O	0	0
			140	140		
4	F	117	Total	O	0	0
			117	117		
4	G	73	Total	O	0	0
			73	73		
4	H	53	Total	O	0	0
			53	53		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transcriptional regulator, TetR family

Chain A: 




- Molecule 1: Transcriptional regulator, TetR family

Chain B: 




- Molecule 1: Transcriptional regulator, TetR family

Chain C: 



- Molecule 1: Transcriptional regulator, TetR family

Chain D: 



- Molecule 1: Transcriptional regulator, TetR family


Chain E: 





- Molecule 1: Transcriptional regulator, TetR family



- Molecule 1: Transcriptional regulator, TetR family



- Molecule 1: Transcriptional regulator, TetR family





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.38Å 124.99Å 151.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.19 – 1.85 48.19 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.7 (48.19-1.85) 99.7 (48.19-1.85)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.60 (at 1.86Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.179 , 0.218 0.194 , 0.230	Depositor DCC
$R_{free}$ test set	1310 reflections (1.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.1	Xtrriage
Anisotropy	0.191	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 48.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	12866	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 62.58 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.1029e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.55	0/1528	0.60	0/2054
1	B	0.57	0/1547	0.59	0/2081
1	C	0.55	0/1508	0.59	0/2029
1	D	0.56	0/1521	0.59	0/2048
1	E	0.59	0/1542	0.61	0/2076
1	F	0.59	0/1526	0.59	0/2055
1	G	0.52	0/1522	0.57	0/2047
1	H	0.50	0/1500	0.56	0/2020
All	All	0.55	0/12194	0.59	0/16410

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1506	0	1518	13	1
1	B	1524	0	1528	14	0
1	C	1485	0	1493	18	0
1	D	1498	0	1503	10	0
1	E	1519	0	1532	14	1

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	1503	0	1500	14	0
1	G	1499	0	1504	16	0
1	H	1477	0	1480	17	0
2	A	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
3	B	7	0	10	0	0
4	A	127	0	0	0	0
4	B	126	0	0	6	0
4	C	100	0	0	0	0
4	D	108	0	0	2	0
4	E	140	0	0	2	0
4	F	117	0	0	1	0
4	G	73	0	0	3	0
4	H	53	0	0	0	0
All	All	12866	0	12068	107	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (107) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:3:MSE:HE2	1:B:5:ILE:HD11	1.21	1.17
1:F:3:MSE:CE	1:F:5:ILE:HD11	1.75	1.15
1:F:3:MSE:HE2	1:F:5:ILE:HD11	1.09	1.06
1:F:3:MSE:HE2	1:F:5:ILE:CD1	1.90	0.99
1:C:3:MSE:CE	1:C:5:ILE:HD11	1.95	0.96
1:B:3:MSE:HE2	1:B:5:ILE:CD1	1.97	0.95
1:B:170[A]:TYR:HB2	4:B:441:HOH:O	1.72	0.87
1:B:3:MSE:CE	1:B:5:ILE:HD11	2.04	0.85
1:B:170[A]:TYR:HD1	4:B:441:HOH:O	1.60	0.83
1:D:11[B]:ARG:HD2	4:D:209:HOH:O	1.79	0.81
1:E:172:ASP:OD1	1:E:174[B]:THR:HG23	1.82	0.80
1:B:170[A]:TYR:CD1	4:B:441:HOH:O	2.35	0.78
1:H:3:MSE:HE2	1:H:5:ILE:HD11	1.67	0.77
1:E:172:ASP:OD1	1:E:174[B]:THR:CG2	2.33	0.76
1:E:174[B]:THR:HG21	4:E:762:HOH:O	1.84	0.75
1:G:152:LEU:HD11	1:G:186:THR:HG21	1.68	0.73
1:C:3:MSE:HE2	1:C:5:ILE:HD11	1.73	0.70

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:5[A]:ILE:O	1:E:5[A]:ILE:HG23	1.92	0.69
4:B:318:HOH:O	1:E:11[A]:ARG:HD2	1.91	0.69
1:G:28:PHE:HB3	4:G:780:HOH:O	1.93	0.69
1:C:169:LEU:HD12	1:H:113:PHE:CE1	2.28	0.69
1:C:3:MSE:HE1	1:C:5:ILE:HD11	1.75	0.66
1:C:169:LEU:CD1	1:H:113:PHE:CE1	2.78	0.66
1:G:183:ILE:HG23	1:G:187:ILE:HD12	1.79	0.64
1:H:187:ILE:O	1:H:188:LYS:HB2	1.98	0.62
1:B:176:LEU:HD12	1:E:5[A]:ILE:HD11	1.82	0.62
1:E:3:MSE:CE	1:E:5[B]:ILE:HD11	2.30	0.61
1:G:15:VAL:HG13	4:G:382:HOH:O	2.00	0.61
1:A:11[A]:ARG:HD2	4:D:394:HOH:O	2.02	0.60
1:H:3:MSE:HE2	1:H:5:ILE:CD1	2.33	0.59
1:E:5[A]:ILE:O	1:E:5[A]:ILE:CG2	2.51	0.58
1:G:15:VAL:HG11	1:G:37:ILE:CG2	2.34	0.58
1:A:-1:ASN:OD1	1:A:-1:ASN:N	2.35	0.58
1:F:3:MSE:HE2	1:F:5:ILE:CG1	2.34	0.57
1:G:15:VAL:HG21	1:G:33:ILE:HG23	1.88	0.56
1:C:66:ILE:HD11	1:C:95:LEU:CD1	2.35	0.56
1:C:169:LEU:HD13	1:H:113:PHE:CD1	2.41	0.56
1:B:62:HIS:O	1:B:66:ILE:HD13	2.05	0.56
1:H:85:ILE:HD12	1:H:187:ILE:HD13	1.87	0.56
1:A:66:ILE:HD11	1:A:95:LEU:HD13	1.87	0.55
1:C:7:ASN:O	1:C:11[A]:ARG:HG3	2.06	0.55
1:F:133[A]:GLU:HG2	4:F:734:HOH:O	2.06	0.55
1:A:11[A]:ARG:O	1:A:15[A]:VAL:HG23	2.06	0.55
1:C:85:ILE:HD12	1:C:187:ILE:HD13	1.89	0.54
1:D:169:LEU:HD22	1:G:113:PHE:CD1	2.43	0.54
1:C:113:PHE:HA	1:C:116:LEU:HD12	1.90	0.53
1:H:3:MSE:CE	1:H:5:ILE:HD11	2.36	0.53
1:E:66:ILE:HD11	1:E:95:LEU:HD13	1.91	0.53
1:B:3:MSE:HE2	1:B:5:ILE:CG1	2.39	0.53
1:E:172:ASP:OD1	1:E:174[B]:THR:HG22	2.08	0.53
1:B:170[A]:TYR:CB	4:B:441:HOH:O	2.45	0.53
1:D:113:PHE:HA	1:D:116:LEU:HD12	1.90	0.52
1:C:159:LEU:HG	1:C:179:VAL:HG22	1.91	0.52
1:G:15:VAL:CG2	1:G:33:ILE:HG23	2.40	0.52
1:D:137:GLY:O	1:D:141:THR:HG23	2.10	0.52
1:E:3:MSE:HE1	1:E:5[B]:ILE:HD11	1.90	0.52
1:H:34:ALA:HB1	1:H:39:ILE:O	2.10	0.51
1:E:133:GLU:HG2	4:E:684:HOH:O	2.10	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:3:MSE:HE2	1:C:5:ILE:CD1	2.40	0.51
1:F:85:ILE:CD1	1:F:187:ILE:HD13	2.41	0.50
1:H:56:VAL:HG13	1:H:123:GLN:HG3	1.93	0.50
1:F:85:ILE:HD13	1:F:187:ILE:CD1	2.41	0.50
1:C:81:ALA:CB	1:C:142:ASN:HB2	2.41	0.50
1:H:85:ILE:HD12	1:H:187:ILE:CD1	2.42	0.49
1:B:137:GLY:O	1:B:141:THR:HG23	2.11	0.48
1:G:5:ILE:HG23	1:G:5:ILE:O	2.13	0.48
1:A:11[B]:ARG:O	1:A:15[B]:VAL:HG22	2.13	0.48
1:F:152:LEU:HD11	1:F:186:THR:HG21	1.95	0.48
1:H:66:ILE:O	1:H:69:VAL:HG22	2.12	0.48
1:A:15[A]:VAL:HG21	1:A:37:ILE:HG21	1.96	0.48
1:H:3:MSE:HE2	1:H:5:ILE:CG1	2.44	0.48
1:B:176:LEU:HD12	1:E:5[A]:ILE:CD1	2.42	0.47
1:F:137:GLY:O	1:F:141:THR:HG23	2.14	0.47
1:A:59:ILE:CD1	1:A:120:ILE:HG23	2.45	0.47
1:A:15[A]:VAL:HG21	1:A:37:ILE:CG2	2.45	0.46
1:B:116:LEU:HD13	1:B:120:ILE:HG22	1.96	0.46
1:C:3:MSE:HE2	1:C:5:ILE:CG1	2.45	0.46
1:A:136:LYS:HD2	1:A:150:LEU:HD13	1.97	0.46
1:A:66:ILE:HD11	1:A:95:LEU:CD1	2.46	0.45
1:F:159:LEU:HG	1:F:179:VAL:HG22	1.99	0.45
1:G:15:VAL:HG11	1:G:37:ILE:HG21	1.98	0.45
1:A:183:ILE:HG23	1:A:187:ILE:HD12	1.98	0.45
1:A:5:ILE:HD11	1:D:176:LEU:HD12	1.99	0.44
1:F:152:LEU:HD11	1:F:186:THR:CG2	2.47	0.44
1:G:11[A]:ARG:HD2	4:G:190:HOH:O	2.16	0.44
1:H:136:LYS:O	1:H:140:GLU:HG3	2.18	0.44
1:F:85:ILE:HD12	1:F:187:ILE:HD13	1.99	0.44
1:C:183:ILE:HG23	1:C:187:ILE:HD12	2.00	0.43
1:G:15:VAL:HG11	1:G:37:ILE:HG23	1.99	0.43
1:A:176:LEU:HD12	1:D:5:ILE:HD11	1.99	0.43
1:B:11[A]:ARG:HD3	4:B:305:HOH:O	2.17	0.43
1:E:1:MSE:HA	1:E:1:MSE:HE2	2.01	0.43
1:H:113:PHE:HA	1:H:116:LEU:HD12	2.01	0.42
1:C:3:MSE:SE	1:C:5:ILE:HD11	2.68	0.42
1:G:3[B]:MSE:HE3	1:G:5:ILE:HG21	2.02	0.42
1:G:159:LEU:HG	1:G:179:VAL:HG22	2.01	0.42
1:D:34:ALA:HB1	1:D:39:ILE:O	2.21	0.41
1:C:93:LEU:CD2	1:C:176:LEU:HD13	2.50	0.41
1:H:66:ILE:O	1:H:70:LEU:HG	2.20	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:136:LYS:NZ	1:D:146:ASN:OD1	2.47	0.41
1:H:21:GLN:HB3	1:H:102:LYS:HB3	2.02	0.41
1:D:162:GLY:HA3	1:G:158:SER:O	2.21	0.41
1:F:85:ILE:CD1	1:F:187:ILE:CD1	2.98	0.41
1:G:3[B]:MSE:HE3	1:G:5:ILE:CG2	2.50	0.41
1:C:164:LEU:HD23	1:C:176:LEU:HD21	2.02	0.40
1:D:66[A]:ILE:HD11	1:D:95:LEU:CD1	2.51	0.40
1:F:139:LEU:HD13	1:F:152:LEU:HB3	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:46:TYR:OH	1:E:46:TYR:OH[4_465]	2.15	0.05

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	190/191 (100%)	189 (100%)	1 (0%)	0	100	100
1	B	192/191 (100%)	189 (98%)	3 (2%)	0	100	100
1	C	187/191 (98%)	186 (100%)	1 (0%)	0	100	100
1	D	189/191 (99%)	188 (100%)	1 (0%)	0	100	100
1	E	192/191 (100%)	191 (100%)	1 (0%)	0	100	100
1	F	189/191 (99%)	188 (100%)	1 (0%)	0	100	100
1	G	189/191 (99%)	188 (100%)	1 (0%)	0	100	100
1	H	186/191 (97%)	185 (100%)	1 (0%)	0	100	100
All	All	1514/1528 (99%)	1504 (99%)	10 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	165/160 (103%)	164 (99%)	1 (1%)	86	83
1	B	167/160 (104%)	167 (100%)	0	100	100
1	C	163/160 (102%)	161 (99%)	2 (1%)	71	62
1	D	165/160 (103%)	164 (99%)	1 (1%)	86	83
1	E	168/160 (105%)	168 (100%)	0	100	100
1	F	165/160 (103%)	165 (100%)	0	100	100
1	G	165/160 (103%)	162 (98%)	3 (2%)	59	45
1	H	162/160 (101%)	161 (99%)	1 (1%)	86	83
All	All	1320/1280 (103%)	1312 (99%)	8 (1%)	86	83

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-1	ASN
1	C	31	LYS
1	C	169	LEU
1	D	170	TYR
1	G	130	LEU
1	G	180	ARG
1	G	188	LYS
1	H	188	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	G	181	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	PEG	B	189	-	6,6,6	0.47	0	5,5,5	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEG	B	189	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.



No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	186/191 (97%)	-0.06	3 (1%) 72 72	13, 22, 42, 62	0
1	B	185/191 (96%)	-0.08	2 (1%) 80 81	14, 21, 49, 72	0
1	C	184/191 (96%)	-0.06	3 (1%) 72 72	14, 24, 45, 63	0
1	D	183/191 (95%)	-0.03	2 (1%) 80 81	16, 25, 46, 61	0
1	E	184/191 (96%)	-0.14	2 (1%) 80 81	13, 21, 40, 75	0
1	F	183/191 (95%)	-0.11	1 (0%) 91 91	13, 20, 35, 54	0
1	G	184/191 (96%)	0.21	5 (2%) 54 53	19, 29, 53, 82	0
1	H	183/191 (95%)	0.39	14 (7%) 13 13	18, 33, 72, 94	0
All	All	1472/1528 (96%)	0.02	32 (2%) 62 61	13, 25, 48, 94	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	0	ALA	4.1
1	D	76	ASN	3.4
1	G	170	TYR	3.3
1	B	0	ALA	3.3
1	E	188	LYS	3.3
1	A	-1	ASN	3.2
1	G	188	LYS	3.1
1	H	79	LEU	2.9
1	B	188	LYS	2.9
1	A	77	SER	2.7
1	H	40	LYS	2.7
1	H	4	ASN	2.7
1	H	31	LYS	2.7
1	F	77	SER	2.6
1	H	78	SER	2.6
1	G	169	LEU	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	H	170	TYR	2.5
1	H	138	VAL	2.4
1	H	143	GLY	2.4
1	G	171	GLY	2.3
1	H	27	ALA	2.3
1	H	28	PHE	2.2
1	H	181	HIS	2.2
1	H	144	TYR	2.2
1	C	2	PRO	2.1
1	H	185	GLN	2.1
1	D	171	GLY	2.1
1	C	170	TYR	2.1
1	H	30	PHE	2.1
1	E	142	ASN	2.1
1	C	72	ASP	2.0
1	G	181	HIS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	PEG	B	189	7/7	0.83	0.16	44,45,46,47	0
2	NA	E	189	1/1	0.98	0.09	16,16,16,16	0
2	NA	D	189	1/1	0.99	0.05	21,21,21,21	0
2	NA	A	189	1/1	0.99	0.07	16,16,16,16	0
2	NA	C	189	1/1	0.99	0.05	26,26,26,26	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.