



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 15, 2024 – 07:32 AM EST

PDB ID : 3P4G
Title : X-ray crystal structure of a hyperactive, Ca²⁺-dependent, beta-helical antifreeze protein from an Antarctic bacterium
Authors : Garnham, C.P.; Campbell, R.L.; Davies, P.L.
Deposited on : 2010-10-06
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	323	 2% 83% 9% 7%
1	B	323	 6% 89% 8%
1	C	323	 2% 85% 7% 7%
1	D	323	 3% 89% 7%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 10219 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Antifreeze protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	301	2222	1356	376	488	2	0	11	0
1	B	313	2303	1406	389	506	2	0	9	0
1	C	301	2205	1342	374	487	2	0	7	0
1	D	312	2292	1398	389	503	2	0	7	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP A1YIY3
B	1	MET	-	initiating methionine	UNP A1YIY3
C	1	MET	-	initiating methionine	UNP A1YIY3
D	1	MET	-	initiating methionine	UNP A1YIY3

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	13	Total 13	Ca 13	0	0
2	B	13	Total 13	Ca 13	0	0
2	C	13	Total 13	Ca 13	0	0
2	D	13	Total 13	Ca 13	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	3	Total Mg 3 3	0	0
3	B	1	Total Mg 1 1	0	0

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0



- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	292	Total O 292 292	0	0
5	B	232	Total O 232 232	0	0
5	C	282	Total O 282 282	0	0
5	D	327	Total O 327 327	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Antifreeze protein




- Molecule 1: Antifreeze protein



- Molecule 1: Antifreeze protein



- Molecule 1: Antifreeze protein





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	57.50Å 70.78Å 84.25Å 99.06° 92.30° 97.47°	Depositor
Resolution (Å)	34.60 – 1.70 34.61 – 1.70	Depositor EDS
% Data completeness (in resolution range)	94.9 (34.60-1.70) 94.9 (34.61-1.70)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.44 (at 1.70Å)	Xtrriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.164 , 0.200 0.181 , 0.210	Depositor DCC
R_{free} test set	6803 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	12.1	Xtrriage
Anisotropy	0.197	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 34.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10219	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, CA, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.88	0/2277	0.87	4/3098 (0.1%)
1	B	0.79	0/2355	0.83	3/3205 (0.1%)
1	C	0.76	0/2244	0.80	4/3052 (0.1%)
1	D	0.73	0/2338	0.80	7/3181 (0.2%)
All	All	0.79	0/9214	0.82	18/12536 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	258	ASP	O-C-N	-7.55	110.61	122.70
1	C	50	LEU	CA-CB-CG	7.28	132.04	115.30
1	C	50	LEU	CB-CG-CD1	6.02	121.23	111.00
1	D	50	LEU	CA-CB-CG	5.94	128.96	115.30
1	D	56	ASP	CB-CG-OD1	5.88	123.60	118.30
1	D	116[A]	ASP	CB-CG-OD1	5.85	123.57	118.30
1	D	116[B]	ASP	CB-CG-OD1	5.85	123.57	118.30
1	A	319	ASP	CB-CG-OD1	5.69	123.42	118.30
1	C	199	ASP	CB-CG-OD2	5.24	123.02	118.30
1	B	199	ASP	CB-CG-OD2	5.22	123.00	118.30
1	D	258	ASP	CB-CG-OD2	5.22	123.00	118.30
1	B	258	ASP	CB-CG-OD2	5.21	122.99	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	199	ASP	CB-CG-OD2	5.20	122.98	118.30
1	C	258	ASP	CB-CG-OD2	5.19	122.97	118.30
1	A	258	ASP	CB-CG-OD2	5.18	122.96	118.30
1	A	199	ASP	CB-CG-OD2	5.16	122.95	118.30
1	D	97	ASP	CB-CG-OD1	5.12	122.91	118.30
1	B	68	VAL	CG1-CB-CG2	5.04	118.97	110.90

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	258	ASP	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2222	0	2109	25	0
1	B	2303	0	2173	17	0
1	C	2205	0	2082	20	0
1	D	2292	0	2155	15	0
2	A	13	0	0	0	0
2	B	13	0	0	0	0
2	C	13	0	0	0	0
2	D	13	0	0	0	0
3	A	3	0	0	0	0
3	B	1	0	0	0	0
4	B	4	0	3	0	0
4	C	4	0	3	0	0
5	A	292	0	0	8	0
5	B	232	0	0	2	0
5	C	282	0	0	4	0
5	D	327	0	0	1	0
All	All	10219	0	8525	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:104[B]:GLU:OE1	5:C:1659:HOH:O	1.72	1.08
1:A:283:LYS:HE3	1:A:323:TYR:C	1.80	1.01
1:C:232:GLN:HE21	1:C:234:GLY:H	1.07	0.93
1:B:129:ASN:HD22	1:B:148:ASP:H	1.20	0.90
1:C:129:ASN:HD22	1:C:148:ASP:H	1.18	0.87
1:B:27:HIS:HD2	1:B:45:ASN:H	1.23	0.86
1:D:232[B]:GLN:HE21	1:D:234:GLY:H	1.25	0.84
1:D:129:ASN:HD22	1:D:148:ASP:H	1.28	0.81
1:B:249:GLU:HB2	5:B:347:HOH:O	1.80	0.80
1:B:232:GLN:HE21	1:B:234:GLY:H	1.29	0.80
1:A:286:THR:HG23	5:A:1537:HOH:O	1.83	0.79
1:D:27:HIS:HE1	1:D:47:SER:OG	1.69	0.75
1:C:23:HIS:HE1	1:C:38:ASN:HD22	1.32	0.75
1:C:286:THR:HG22	1:C:309:SER:OG	1.90	0.71
1:A:129:ASN:HD22	1:A:148:ASP:H	1.38	0.71
1:B:129:ASN:ND2	1:B:148:ASP:H	1.89	0.71
1:A:173:ASN:HB2	5:A:378:HOH:O	1.92	0.68
1:A:286:THR:HG22	1:A:309:SER:OG	1.96	0.66
1:A:283:LYS:HE3	1:A:323:TYR:O	1.96	0.65
1:C:273:GLY:H	1:C:302:ASN:ND2	1.95	0.65
1:D:286:THR:HG22	1:D:309:SER:OG	1.98	0.64
1:D:129:ASN:ND2	1:D:148:ASP:H	1.97	0.63
1:B:230[B]:ASN:ND2	5:B:1647:HOH:O	2.32	0.61
1:C:286:THR:HG22	1:C:309:SER:HG	1.63	0.61
1:B:286:THR:H	1:B:289:GLN:HE21	1.48	0.60
1:D:272:ALA:HA	1:D:302:ASN:HD21	1.67	0.60
1:A:232[B]:GLN:HG3	5:A:1634:HOH:O	2.02	0.59
1:A:27:HIS:HE1	1:A:47:SER:OG	1.85	0.59
1:A:263:VAL:HG23	5:A:1507:HOH:O	2.01	0.58
1:B:286:THR:H	1:B:289:GLN:NE2	2.02	0.58
1:D:33:PHE:CE1	1:D:50:LEU:HD13	2.40	0.57
1:C:273:GLY:H	1:C:302:ASN:HD21	1.52	0.57
1:A:163:ILE:HG12	5:A:375:HOH:O	2.05	0.56
1:B:27:HIS:CD2	1:B:45:ASN:H	2.12	0.56
1:B:272:ALA:HA	1:B:302:ASN:HD21	1.72	0.54
1:A:273:GLY:H	1:A:302:ASN:ND2	2.05	0.54
1:C:129:ASN:ND2	1:C:148:ASP:H	1.98	0.53
1:A:230:ASN:HD21	1:A:251[B]:ASN:HD22	1.55	0.52
1:D:146:GLY:O	1:D:166:HIS:HD2	1.93	0.52
1:D:232[B]:GLN:NE2	5:D:799:HOH:O	2.43	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:27:HIS:HE1	1:C:47:SER:OG	1.95	0.49
1:B:273:GLY:H	1:B:302:ASN:ND2	2.10	0.49
1:D:286:THR:H	1:D:289:GLN:NE2	2.11	0.48
1:D:273:GLY:H	1:D:302:ASN:ND2	2.11	0.48
1:D:50:LEU:HG	1:D:77:LEU:HD13	1.96	0.48
1:A:151:ALA:HB3	5:A:592:HOH:O	2.14	0.47
1:D:27:HIS:CE1	1:D:47:SER:OG	2.60	0.47
1:A:269:GLY:O	1:A:299:ARG:HD3	2.16	0.46
1:A:286:THR:HG22	1:A:309:SER:HG	1.81	0.46
1:B:12:TRP:CE2	1:B:58:LYS:HE2	2.51	0.45
1:A:129:ASN:ND2	1:A:148:ASP:H	2.10	0.45
1:A:286:THR:H	1:A:289:GLN:NE2	2.16	0.44
1:B:232:GLN:HE21	1:B:234:GLY:N	2.07	0.44
1:B:251:ASN:HD22	1:B:270:THR:HB	1.83	0.44
1:C:77:LEU:HD21	1:C:86:LEU:CD1	2.49	0.43
1:D:167:ASP:OD1	1:D:185:ARG:HB3	2.19	0.43
1:B:225:GLY:C	1:B:244:SER:HB2	2.39	0.43
1:C:286:THR:CG2	5:C:1552:HOH:O	2.67	0.43
1:C:77:LEU:HD21	1:C:86:LEU:HD11	2.00	0.43
1:C:286:THR:HG23	5:C:1552:HOH:O	2.18	0.43
1:D:273:GLY:H	1:D:302:ASN:HD21	1.65	0.43
1:C:166:HIS:HE1	5:C:377:HOH:O	2.02	0.42
1:A:232[B]:GLN:CG	5:A:1634:HOH:O	2.62	0.42
1:A:254:LEU:HD11	1:A:261:LEU:HB2	2.01	0.42
1:A:318:ALA:HB2	1:B:55:GLY:HA2	2.02	0.42
1:A:98:ALA:HB2	1:A:105:ILE:HD11	2.02	0.42
1:C:286:THR:H	1:C:289:GLN:NE2	2.17	0.42
1:C:104[A]:GLU:HG3	1:C:123[A]:GLU:HB3	2.02	0.42
1:C:254:LEU:HB3	1:C:259:ASP:CB	2.50	0.41
1:A:273:GLY:H	1:A:302:ASN:HD21	1.67	0.41
1:C:68:VAL:HG22	1:C:76:VAL:HG22	2.02	0.41
1:A:254:LEU:HB3	1:A:259:ASP:CB	2.50	0.41
1:B:232:GLN:NE2	1:B:234:GLY:H	2.08	0.41
1:C:269:GLY:O	1:C:299:ARG:HD3	2.19	0.41
1:A:285:TYR:O	1:A:308:VAL:HB	2.21	0.41
1:A:161:ASN:ND2	5:A:1373:HOH:O	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	310/323 (96%)	294 (95%)	16 (5%)	0	100	100
1	B	320/323 (99%)	307 (96%)	12 (4%)	1 (0%)	41	24
1	C	306/323 (95%)	290 (95%)	16 (5%)	0	100	100
1	D	317/323 (98%)	302 (95%)	15 (5%)	0	100	100
All	All	1253/1292 (97%)	1193 (95%)	59 (5%)	1 (0%)	51	33

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	26	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	243/251 (97%)	241 (99%)	2 (1%)	81	74
1	B	251/251 (100%)	246 (98%)	5 (2%)	55	38
1	C	239/251 (95%)	236 (99%)	3 (1%)	69	56
1	D	248/251 (99%)	244 (98%)	4 (2%)	62	48
All	All	981/1004 (98%)	967 (99%)	14 (1%)	67	53

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	263	VAL
1	A	289	GLN
1	B	8	SER
1	B	249	GLU
1	B	288[A]	GLU
1	B	288[B]	GLU
1	B	319	ASP
1	C	50	LEU
1	C	68	VAL
1	C	289	GLN
1	D	24	SER
1	D	50	LEU
1	D	289	GLN
1	D	319	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (26) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	27	HIS
1	A	129	ASN
1	A	230	ASN
1	A	289	GLN
1	A	302	ASN
1	B	27	HIS
1	B	129	ASN
1	B	232	GLN
1	B	251	ASN
1	B	289	GLN
1	B	302	ASN
1	C	23	HIS
1	C	27	HIS
1	C	129	ASN
1	C	166	HIS
1	C	230	ASN
1	C	232	GLN
1	C	251	ASN
1	C	289	GLN
1	C	302	ASN
1	D	27	HIS
1	D	129	ASN
1	D	166	HIS
1	D	230	ASN
1	D	289	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	302	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 58 ligands modelled in this entry, 56 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ACT	B	325	-	3,3,3	0.58	0	3,3,3	1.36	0
4	ACT	C	324	-	3,3,3	0.61	0	3,3,3	0.96	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	301/323 (93%)	0.10	5 (1%) 70 74	5, 11, 23, 44	0
1	B	313/323 (96%)	0.10	20 (6%) 19 21	6, 14, 35, 49	0
1	C	301/323 (93%)	0.05	8 (2%) 54 58	8, 18, 29, 44	0
1	D	312/323 (96%)	-0.05	10 (3%) 47 52	10, 17, 34, 47	0
All	All	1227/1292 (94%)	0.05	43 (3%) 44 49	5, 15, 32, 49	0

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	321	ALA	7.6
1	B	8	SER	5.2
1	B	290	TYR	4.8
1	B	286	THR	4.6
1	D	19	ALA	3.9
1	D	18	GLN	3.9
1	D	17	THR	3.9
1	D	9	PHE	3.9
1	C	71	ALA	3.5
1	B	319	ASP	3.4
1	B	316[A]	SER	3.4
1	B	294[A]	SER	3.3
1	C	286	THR	3.2
1	C	72	ALA	3.0
1	B	295	ASP	2.9
1	D	286	THR	2.7
1	B	313	VAL	2.7
1	B	291	ASN	2.7
1	D	292[A]	ASN	2.6
1	B	312	VAL	2.6
1	A	322	ASP	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	69	LEU	2.6
1	B	284	PHE	2.5
1	B	308	VAL	2.5
1	A	286	THR	2.4
1	B	311	GLY	2.4
1	D	10	GLY	2.4
1	A	26	ASP	2.3
1	A	321	ALA	2.3
1	C	247	TRP	2.3
1	B	310	ASP	2.3
1	B	315	GLY	2.3
1	D	288	GLU	2.2
1	C	295	ASP	2.2
1	B	292	ASN	2.2
1	B	318	ALA	2.2
1	C	143	ILE	2.1
1	D	295	ASP	2.1
1	B	287	LYS	2.0
1	A	120	GLY	2.0
1	B	9	PHE	2.0
1	B	263[A]	VAL	2.0
1	D	319	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MG	A	326	1/1	0.93	0.24	23,23,23,23	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	MG	B	324	1/1	0.95	0.20	18,18,18,18	0
3	MG	A	324	1/1	0.96	0.15	13,13,13,13	0
4	ACT	C	324	4/4	0.97	0.07	18,18,18,19	0
4	ACT	B	325	4/4	0.98	0.10	13,16,16,17	0
3	MG	A	325	1/1	0.98	0.14	13,13,13,13	0
2	CA	C	413	1/1	0.99	0.03	7,7,7,7	0
2	CA	D	401	1/1	0.99	0.08	4,4,4,4	0
2	CA	D	402	1/1	0.99	0.03	4,4,4,4	0
2	CA	D	403	1/1	0.99	0.03	3,3,3,3	0
2	CA	D	405	1/1	0.99	0.04	4,4,4,4	0
2	CA	D	406	1/1	0.99	0.03	5,5,5,5	0
2	CA	D	407	1/1	0.99	0.03	6,6,6,6	0
2	CA	D	413	1/1	0.99	0.14	9,9,9,9	0
2	CA	B	402	1/1	0.99	0.04	3,3,3,3	0
2	CA	B	413	1/1	0.99	0.09	5,5,5,5	0
2	CA	C	401	1/1	0.99	0.06	8,8,8,8	0
2	CA	C	402	1/1	0.99	0.04	5,5,5,5	0
2	CA	C	403	1/1	0.99	0.03	5,5,5,5	0
2	CA	C	404	1/1	0.99	0.03	5,5,5,5	0
2	CA	B	408	1/1	1.00	0.05	4,4,4,4	0
2	CA	B	409	1/1	1.00	0.04	4,4,4,4	0
2	CA	B	410	1/1	1.00	0.04	4,4,4,4	0
2	CA	B	411	1/1	1.00	0.02	4,4,4,4	0
2	CA	B	412	1/1	1.00	0.04	5,5,5,5	0
2	CA	A	402	1/1	1.00	0.04	3,3,3,3	0
2	CA	A	403	1/1	1.00	0.06	3,3,3,3	0
2	CA	A	404	1/1	1.00	0.09	3,3,3,3	0
2	CA	A	405	1/1	1.00	0.09	4,4,4,4	0
2	CA	A	406	1/1	1.00	0.09	4,4,4,4	0
2	CA	C	405	1/1	1.00	0.04	4,4,4,4	0
2	CA	C	406	1/1	1.00	0.04	5,5,5,5	0
2	CA	C	407	1/1	1.00	0.07	5,5,5,5	0
2	CA	C	408	1/1	1.00	0.05	5,5,5,5	0
2	CA	C	409	1/1	1.00	0.04	5,5,5,5	0
2	CA	C	410	1/1	1.00	0.03	6,6,6,6	0
2	CA	C	411	1/1	1.00	0.02	7,7,7,7	0
2	CA	C	412	1/1	1.00	0.02	6,6,6,6	0
2	CA	A	407	1/1	1.00	0.10	4,4,4,4	0
2	CA	A	408	1/1	1.00	0.09	4,4,4,4	0
2	CA	A	409	1/1	1.00	0.07	4,4,4,4	0
2	CA	A	410	1/1	1.00	0.09	4,4,4,4	0
2	CA	D	404	1/1	1.00	0.02	5,5,5,5	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CA	A	411	1/1	1.00	0.05	3,3,3,3	0
2	CA	A	412	1/1	1.00	0.04	3,3,3,3	0
2	CA	A	413	1/1	1.00	0.01	3,3,3,3	0
2	CA	D	408	1/1	1.00	0.02	6,6,6,6	0
2	CA	D	409	1/1	1.00	0.03	6,6,6,6	0
2	CA	D	410	1/1	1.00	0.05	6,6,6,6	0
2	CA	D	411	1/1	1.00	0.07	5,5,5,5	0
2	CA	D	412	1/1	1.00	0.12	7,7,7,7	0
2	CA	B	401	1/1	1.00	0.03	3,3,3,3	0
2	CA	A	401	1/1	1.00	0.02	3,3,3,3	0
2	CA	B	403	1/1	1.00	0.05	3,3,3,3	0
2	CA	B	404	1/1	1.00	0.05	4,4,4,4	0
2	CA	B	405	1/1	1.00	0.07	4,4,4,4	0
2	CA	B	406	1/1	1.00	0.06	3,3,3,3	0
2	CA	B	407	1/1	1.00	0.05	3,3,3,3	0

6.5 Other polymers [i](#)

There are no such residues in this entry.