



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 5, 2023 – 09:27 PM EST

PDB ID : 3UT5
Title : Tubulin-Colchicine-Ustiloxin: Stathmin-like domain complex
Authors : Ranaivoson, F.M.; Gigant, B.; Knossow, M.
Deposited on : 2011-11-25
Resolution : 2.73 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1271 (2.76-2.72)
Clashscore	141614	1322 (2.76-2.72)
Ramachandran outliers	138981	1297 (2.76-2.72)
Sidechain outliers	138945	1298 (2.76-2.72)
RSRZ outliers	127900	1243 (2.76-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	451	 7% 77% 16% . .
1	C	451	 4% 74% 20% . .
2	B	445	 11% 76% 18% . .
2	D	445	 5% 76% 16% . .
3	E	142	 13% 73% 18% . 7%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	F	4	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	0E5	F	3	X	X	-	-

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 15107 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tubulin alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	433	3399	2153	577	647	22	0	1	0
1	C	432	3404	2158	575	648	23	0	4	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	232	SER	GLY	SEE REMARK 999	UNP D0VWZ0
A	340	SER	THR	SEE REMARK 999	UNP D0VWZ0
C	232	SER	GLY	SEE REMARK 999	UNP D0VWZ0
C	340	SER	THR	SEE REMARK 999	UNP D0VWZ0

- Molecule 2 is a protein called Tubulin beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	432	3417	2141	584	665	27	0	3	0
2	D	432	3433	2153	584	670	26	0	5	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	317	THR	ALA	SEE REMARK 999	UNP D0VWY9
B	318	ILE	VAL	SEE REMARK 999	UNP D0VWY9
B	335	ILE	VAL	SEE REMARK 999	UNP D0VWY9
B	375	SER	ALA	SEE REMARK 999	UNP D0VWY9
D	317	THR	ALA	SEE REMARK 999	UNP D0VWY9
D	318	ILE	VAL	SEE REMARK 999	UNP D0VWY9
D	335	ILE	VAL	SEE REMARK 999	UNP D0VWY9
D	375	SER	ALA	SEE REMARK 999	UNP D0VWY9

- Molecule 3 is a protein called Stathmin-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	E	132	1077	666	195	212	4	0	1	0


There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	4	ALA	-	expression tag	UNP P63043
E	14	ALA	CYS	engineered mutation	UNP P63043
E	20	TRP	PHE	engineered mutation	UNP P63043

- Molecule 4 is a protein called Vinca tetrapeptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	F	4	35	23	4	8	0	0	0

- Molecule 5 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
5	A	1	32	10	5	14	3	0	0
5	C	1	32	10	5	14	3	0	0

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Mg	0	0
			1	1		
6	C	1	Total	Mg	0	0
			1	1		

- Molecule 7 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	O	S	0	0
			5	4	1		
7	A	1	Total	O	S	0	0
			5	4	1		
7	B	1	Total	O	S	0	0
			5	4	1		
7	D	1	Total	O	S	0	0
			5	4	1		
7	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 8 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: C₁₀H₁₅N₅O₁₁P₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
8	B	1	28	10	5	11	2	0	0
8	D	1	28	10	5	11	2	0	0

- Molecule 9 is N-[(7S)-1,2,3,10-tetramethoxy-9-oxo-6,7-dihydro-5H-benzo[d]heptalen-7-yl]ethanamide (three-letter code: LOC) (formula: C₂₂H₂₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
9	B	1	29	22	1	6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	D	1	Total	C	N	O	0	0
			29	22	1	6		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	27	Total	O	0	0
			27	27		
10	B	28	Total	O	0	0
			28	28		
10	C	43	Total	O	0	0
			43	43		
10	D	36	Total	O	0	0
			36	36		
10	E	3	Total	O	0	0
			3	3		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tubulin alpha chain




- Molecule 1: Tubulin alpha chain





- Molecule 2: Tubulin beta chain






• Molecule 2: Tubulin beta chain



• Molecule 3: Stathmin-4



• Molecule 4: Vinca tetrapeptide



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	64.81Å 128.87Å 254.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.26 – 2.73 38.26 – 2.73	Depositor EDS
% Data completeness (in resolution range)	(Not available) (38.26-2.73) 97.6 (38.26-2.73)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.73 (at 2.72Å)	Xtrriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.184 , 0.222 0.201 , 0.241	Depositor DCC
R_{free} test set	2872 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	66.6	Xtrriage
Anisotropy	0.638	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 60.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	15107	wwPDB-VP
Average B, all atoms (Å ²)	93.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, 0EA, GTP, MG, SO4, 0E5, LOC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/3479	0.68	0/4723
1	C	0.45	0/3491	0.70	0/4741
2	B	0.42	0/3503	0.69	0/4744
2	D	0.43	0/3516	0.68	0/4762
3	E	0.46	0/1092	0.68	0/1455
4	F	1.32	0/10	0.38	0/9
All	All	0.44	0/15091	0.69	0/20434

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
4	F	1	0

There are no bond length outliers.

There are no bond angle outliers.

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	F	3	0E5	C3

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3399	0	3312	31	0
1	C	3404	0	3311	39	0
2	B	3417	0	3283	47	0
2	D	3433	0	3301	53	0
3	E	1077	0	1068	7	0
4	F	35	0	24	1	0
5	A	32	0	12	1	0
5	C	32	0	12	1	0
6	A	1	0	0	0	0
6	C	1	0	0	0	0
7	A	10	0	0	0	0
7	B	5	0	0	0	0
7	D	10	0	0	0	0
8	B	28	0	12	3	0
8	D	28	0	12	2	0
9	B	29	0	25	4	0
9	D	29	0	25	3	0
10	A	27	0	0	0	0
10	B	28	0	0	1	0
10	C	43	0	0	0	0
10	D	36	0	0	1	0
10	E	3	0	0	0	0
All	All	15107	0	14397	168	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 168 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:206:ASN:HD21	8:B:501:GDP:HN22	1.24	0.86
2:D:237:GLY:HA3	2:D:376:THR:HG21	1.59	0.85
1:C:206:ASN:HD21	5:C:600:GTP:HN22	1.24	0.83
1:A:206:ASN:HD21	5:A:501:GTP:HN22	1.26	0.82
2:D:206:ASN:HD21	8:D:501:GDP:HN22	1.25	0.82

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	430/451 (95%)	404 (94%)	21 (5%)	5 (1%)	13	24
1	C	432/451 (96%)	406 (94%)	23 (5%)	3 (1%)	22	40
2	B	433/445 (97%)	411 (95%)	18 (4%)	4 (1%)	17	32
2	D	435/445 (98%)	411 (94%)	17 (4%)	7 (2%)	9	17
3	E	129/142 (91%)	118 (92%)	7 (5%)	4 (3%)	4	6
4	F	1/4 (25%)	0	1 (100%)	0	100	100
All	All	1860/1938 (96%)	1750 (94%)	87 (5%)	23 (1%)	13	24

5 of 23 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	178[A]	SER
2	D	178[B]	SER
2	D	248	LEU
2	B	178	SER
2	B	248	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	368/379 (97%)	326 (89%)	42 (11%)	5	9
1	C	369/379 (97%)	322 (87%)	47 (13%)	4	6
2	B	378/385 (98%)	331 (88%)	47 (12%)	4	7

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	D	380/385 (99%)	336 (88%)	44 (12%)	5	9
3	E	113/125 (90%)	93 (82%)	20 (18%)	2	2
4	F	1/1 (100%)	1 (100%)	0	100	100
All	All	1609/1654 (97%)	1409 (88%)	200 (12%)	4	7

5 of 200 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	282	TYR
2	D	85	GLN
3	E	139	LEU
1	C	302	MET
1	C	384	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 62 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	8	HIS
2	D	385	GLN
1	C	176	GLN
2	D	380	ASN
3	E	108	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	0E5	F	3	4	6,7,9	1.26	0	5,8,13	4.69	4 (80%)
4	0EA	F	1	4	13,15,16	2.71	8 (61%)	17,20,22	2.90	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	0E5	F	3	4	1/1/2/3	6/7/8/13	-
4	0EA	F	1	4	-	0/9/12/14	0/1/1/1

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	F	1	0EA	O5-C7	-4.89	1.26	1.36
4	F	1	0EA	C6-C5	3.98	1.44	1.38
4	F	1	0EA	O1-C5	3.92	1.44	1.36
4	F	1	0EA	C6-C10	3.64	1.44	1.39
4	F	1	0EA	O8-C11	2.72	1.48	1.42

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	F	3	0E5	C3-CA-C	9.21	126.91	112.83
4	F	1	0EA	C10-C11-CA	-7.28	102.82	112.16
4	F	1	0EA	C6-C5-C7	-6.11	114.46	119.86
4	F	1	0EA	C8-C7-C5	5.56	125.77	119.67
4	F	3	0E5	O-C-CA	-3.52	115.54	124.78

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	F	3	0E5	C3

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	F	3	0E5	C2-C3-CA-C
4	F	3	0E5	C4-C3-CA-C
4	F	3	0E5	C4-C3-CA-N

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	F	3	0E5	C1-C2-C3-CA
4	F	3	0E5	C1-C2-C3-C4

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	F	3	0E5	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 2 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
9	LOC	D	502	-	31,31,31	2.60	8 (25%)	44,44,44	3.21	20 (45%)
7	SO4	B	503	-	4,4,4	0.17	0	6,6,6	0.09	0
5	GTP	C	600	6	26,34,34	1.27	2 (7%)	32,54,54	1.57	7 (21%)
7	SO4	A	504	-	4,4,4	0.14	0	6,6,6	0.12	0
9	LOC	B	502	-	31,31,31	2.41	8 (25%)	44,44,44	2.74	20 (45%)
7	SO4	D	503	-	4,4,4	0.30	0	6,6,6	0.08	0
8	GDP	D	501	-	24,30,30	0.98	2 (8%)	30,47,47	1.12	3 (10%)
7	SO4	A	503	-	4,4,4	0.29	0	6,6,6	0.10	0
8	GDP	B	501	-	24,30,30	0.99	1 (4%)	30,47,47	1.30	4 (13%)
5	GTP	A	501	6	26,34,34	1.19	2 (7%)	32,54,54	1.55	7 (21%)
7	SO4	D	504	-	4,4,4	0.17	0	6,6,6	0.13	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	LOC	D	502	-	-	2/12/25/25	0/3/3/3
5	GTP	C	600	6	-	5/18/38/38	0/3/3/3
9	LOC	B	502	-	-	2/12/25/25	0/3/3/3
8	GDP	D	501	-	-	4/12/32/32	0/3/3/3
8	GDP	B	501	-	-	4/12/32/32	0/3/3/3
5	GTP	A	501	6	-	5/18/38/38	0/3/3/3

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	D	502	LOC	C11-C14	-9.13	1.38	1.53
9	B	502	LOC	C22-C21	-6.77	1.41	1.50
9	B	502	LOC	C11-C14	-6.43	1.42	1.53
9	D	502	LOC	C9-C8	-5.30	1.39	1.51
9	B	502	LOC	C9-C8	-5.28	1.39	1.51

The worst 5 of 61 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	D	502	LOC	C1-C22-C21	7.19	128.59	121.09
9	D	502	LOC	C22-C21-C14	6.61	123.83	118.51
9	D	502	LOC	O6-C17-C16	6.61	116.01	109.56
9	D	502	LOC	C11-C14-C15	-6.60	110.88	117.12
9	B	502	LOC	C9-C8-C22	6.28	129.77	119.97

There are no chirality outliers.

5 of 22 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	501	GTP	PB-O3B-PG-O2G
5	A	501	GTP	C5'-O5'-PA-O1A
5	A	501	GTP	C5'-O5'-PA-O2A
5	C	600	GTP	PB-O3B-PG-O2G
5	C	600	GTP	C5'-O5'-PA-O1A


There are no ring outliers.


6 monomers are involved in 14 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	D	502	LOC	3	0
5	C	600	GTP	1	0
9	B	502	LOC	4	0
8	D	501	GDP	2	0
8	B	501	GDP	3	0
5	A	501	GTP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	433/451 (96%)	0.39	31 (7%) 15 17	60, 97, 153, 198	0
1	C	432/451 (95%)	0.25	17 (3%) 39 43	42, 81, 133, 180	0
2	B	432/445 (97%)	0.54	51 (11%) 4 4	59, 90, 155, 215	2 (0%)
2	D	432/445 (97%)	0.24	24 (5%) 24 27	44, 77, 122, 162	2 (0%)
3	E	132/142 (92%)	0.98	19 (14%) 2 2	62, 107, 176, 214	0
4	F	2/4 (50%)	1.33	1 (50%) 0 0	117, 117, 117, 129	0
All	All	1863/1938 (96%)	0.40	143 (7%) 13 15	42, 89, 145, 215	4 (0%)

The worst 5 of 143 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	E	34	GLU	8.4
3	E	141	GLU	7.8
2	B	283	TYR	7.5
3	E	140	LYS	7.4
3	E	32	VAL	7.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	0EA	F	1	15/16	0.93	0.26	113,115,118,119	0
4	0E5	F	3	8/10	0.93	0.27	110,113,121,122	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	SO4	D	504	5/5	0.88	0.18	141,143,147,149	0
7	SO4	A	504	5/5	0.90	0.21	140,143,147,147	0
7	SO4	A	503	5/5	0.91	0.16	127,130,133,134	0
7	SO4	B	503	5/5	0.92	0.09	157,160,165,165	0
9	LOC	B	502	29/29	0.94	0.18	75,79,83,85	0
6	MG	A	502	1/1	0.95	0.16	74,74,74,74	0
7	SO4	D	503	5/5	0.95	0.30	142,143,148,149	0
6	MG	C	601	1/1	0.96	0.13	56,56,56,56	0
9	LOC	D	502	29/29	0.97	0.18	51,63,67,67	0
8	GDP	B	501	28/28	0.98	0.18	69,71,76,76	0
8	GDP	D	501	28/28	0.98	0.17	62,66,70,73	0
5	GTP	C	600	32/32	0.98	0.17	57,59,66,68	0
5	GTP	A	501	32/32	0.98	0.20	63,70,76,76	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around LOC B 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around LOC D 502:**




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GDP D 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.