



## Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 03:14 am BST

PDB ID : 3V96  
Title : Complex of matrix metalloproteinase-10 catalytic domain (MMP-10cd) with tissue inhibitor of metalloproteinases-1 (TIMP-1)  
Authors : Batra, J.; Soares, A.S.; Radisky, E.S.  
Deposited on : 2011-12-23  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	184	
2	B	165	

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 2789 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Metalloproteinase inhibitor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	164	1308	834	227	232	15	0	0	0


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	30	ALA	ASN	ENGINEERED MUTATION	UNP P01033

- Molecule 2 is a protein called Stromelysin-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	159	1278	821	207	247	3	0	2	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	Zn	0	0
			2	2		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	3	Total	Ca	0	0
			3	3		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	105	Total	O	0	2
			105	105		
6	B	83	Total	O	0	0
			83	83		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Metalloproteinase inhibitor 1

Chain A: 7% 61% 24% 11%



- Molecule 2: Stromelysin-2

Chain B: 6% 78% 15%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	231.72Å 37.68Å 39.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.00 – 1.90 35.83 – 1.89	Depositor EDS
% Data completeness (in resolution range)	91.8 (35.00-1.90) 91.8 (35.83-1.89)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.83 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, $R_{free}$	0.215 , 0.266 0.215 , 0.270	Depositor DCC
$R_{free}$ test set	1326 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.0	Xtriage
Anisotropy	0.231	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 50.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.054 for -h,l,k	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2789	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.43% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, ZN, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.04	1/1342 (0.1%)	1.00	3/1815 (0.2%)
2	B	0.93	1/1324 (0.1%)	0.87	0/1803
All	All	0.98	2/2666 (0.1%)	0.94	3/3618 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	137	CYS	CB-SG	-6.86	1.70	1.82
2	B	218	GLU	CB-CG	5.76	1.63	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	20	ARG	NE-CZ-NH2	-8.65	115.97	120.30
1	A	20	ARG	NE-CZ-NH1	7.29	123.95	120.30
1	A	84	LEU	CA-CB-CG	-5.94	101.63	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	239	TYR	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1308	0	1277	48	0
2	B	1278	0	1196	34	0
3	A	10	0	0	0	0
4	B	2	0	0	0	0
5	B	3	0	0	0	0
6	A	105	0	0	5	0
6	B	83	0	0	3	0
All	All	2789	0	2473	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:5:PRO:HG3	1:A:150:GLN:NE2	1.61	1.12
1:A:5:PRO:HG3	1:A:150:GLN:HE22	1.07	1.12
1:A:135:ILE:HG23	1:A:136:PRO:CD	1.86	1.05
1:A:135:ILE:HG23	1:A:136:PRO:HD2	1.04	1.03
1:A:136:PRO:O	1:A:137:CYS:HB3	1.56	1.03
1:A:31:GLN:HA	1:A:31:GLN:NE2	1.69	1.03
1:A:31:GLN:HE21	1:A:31:GLN:CA	1.71	1.03
1:A:31:GLN:HA	1:A:31:GLN:HE21	0.86	1.02
1:A:135:ILE:HG22	2:B:207:SER:HA	1.41	0.99
2:B:150:ARG:HG3	2:B:150:ARG:HH11	1.28	0.95
1:A:70:CYS:HA	1:A:98:THR:CG2	2.01	0.89
2:B:116:ARG:HB3	2:B:159:MET:CE	2.04	0.87
2:B:116:ARG:HB3	2:B:159:MET:HE1	1.58	0.85
2:B:230:ASN:ND2	2:B:232:GLU:H	1.78	0.82
2:B:213:LEU:HD22	2:B:245:LEU:HD23	1.60	0.81
1:A:6:PRO:O	6:A:370:HOH:O	1.98	0.80

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:5:PRO:CG	1:A:150:GLN:NE2	2.46	0.75
1:A:5:PRO:CG	1:A:150:GLN:HE22	1.94	0.75
1:A:17:LEU:HD23	1:A:48:GLY:HA3	1.67	0.74
1:A:114:ARG:HD3	6:A:383:HOH:O	1.86	0.74
1:A:136:PRO:O	1:A:137:CYS:CB	2.33	0.73
1:A:135:ILE:CG2	1:A:136:PRO:HD2	2.00	0.72
1:A:5:PRO:HB3	2:B:238:LEU:HD13	1.74	0.69
1:A:64:PRO:HD2	1:A:71:GLY:HA2	1.76	0.68
1:A:70:CYS:CA	1:A:98:THR:CG2	2.73	0.67
2:B:213:LEU:CD2	2:B:245:LEU:HD23	2.27	0.65
1:A:137:CYS:SG	1:A:146:LEU:HD11	2.37	0.64
1:A:70:CYS:HA	1:A:98:THR:HG23	1.78	0.64
2:B:230:ASN:HD22	2:B:232:GLU:H	1.45	0.63
2:B:238:LEU:O	2:B:240:ASN:HB3	1.98	0.62
1:A:156:GLU:OE2	1:A:157:LYS:HE3	2.00	0.61
1:A:135:ILE:CG2	2:B:207:SER:HA	2.23	0.60
1:A:70:CYS:HA	1:A:98:THR:HG21	1.79	0.60
1:A:135:ILE:HG22	2:B:207:SER:CA	2.25	0.59
1:A:64:PRO:O	1:A:71:GLY:HA3	2.03	0.58
1:A:70:CYS:CA	1:A:98:THR:HG23	2.34	0.58
1:A:128:THR:HG22	6:A:319:HOH:O	2.04	0.57
1:A:70:CYS:C	1:A:98:THR:HG22	2.24	0.57
2:B:230:ASN:HD22	2:B:230:ASN:C	2.08	0.57
1:A:20:ARG:HD2	1:A:82:GLU:OE1	2.06	0.56
2:B:247:GLN:O	2:B:247:GLN:HG2	2.07	0.54
2:B:230:ASN:HD21	2:B:232:GLU:HG2	1.75	0.52
2:B:159:MET:HA	2:B:159:MET:CE	2.40	0.51
1:A:61:VAL:HG22	1:A:96:ILE:HG12	1.93	0.51
1:A:158:GLY:HA2	6:A:377:HOH:O	2.10	0.50
1:A:70:CYS:C	1:A:98:THR:CG2	2.79	0.50
2:B:163:ALA:HB1	2:B:167:HIS:CB	2.41	0.50
2:B:210:ASN:HD22	2:B:213:LEU:H	1.57	0.50
2:B:150:ARG:HH11	2:B:150:ARG:CG	2.08	0.49
2:B:169:ASP:HB2	6:B:445:HOH:O	2.11	0.49
1:A:104:PRO:HD2	1:A:107:SER:OG	2.13	0.49
2:B:247:GLN:OE1	6:B:466:HOH:O	2.20	0.49
1:A:4:VAL:HG13	1:A:133:LEU:HD11	1.95	0.48
2:B:150:ARG:HG3	2:B:150:ARG:NH1	2.10	0.48
2:B:210:ASN:HD22	2:B:213:LEU:HB2	1.79	0.48
2:B:116:ARG:HB3	2:B:159:MET:HE2	1.90	0.47
2:B:159:MET:HA	2:B:159:MET:HE3	1.96	0.47

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:17:LEU:CD2	1:A:48:GLY:HA3	2.42	0.46
1:A:77:HIS:CD2	1:A:77:HIS:N	2.84	0.46
2:B:230:ASN:HD22	2:B:231:THR:N	2.14	0.45
1:A:170:GLU:HB2	1:A:171:PRO:CD	2.47	0.45
1:A:178:SER:C	6:A:363:HOH:O	2.56	0.44
2:B:251:SER:CB	2:B:254:ASP:OD2	2.65	0.44
1:A:141:SER:HB3	1:A:144:HIS:CD2	2.53	0.43
2:B:156:ALA:HB3	2:B:159:MET:CE	2.48	0.43
1:A:84:LEU:HG	1:A:85:ILE:N	2.34	0.43
1:A:17:LEU:HD12	1:A:94:LEU:HD13	2.01	0.43
1:A:7:HIS:HD2	1:A:9:GLN:H	1.65	0.43
1:A:88:LYS:HE2	1:A:90:GLN:HG3	2.01	0.43
2:B:128:ALA:CB	6:B:441:HOH:O	2.67	0.43
2:B:163:ALA:HB1	2:B:167:HIS:HB3	2.01	0.42
1:A:20:ARG:HD3	1:A:105:TRP:CG	2.53	0.42
2:B:242:PHE:CE2	2:B:243:THR:HG22	2.55	0.42
1:A:130:PHE:HA	1:A:131:PRO:HD3	1.89	0.41
2:B:210:ASN:ND2	2:B:213:LEU:H	2.17	0.41
2:B:184:TYR:CE1	2:B:190:LEU:HD22	2.56	0.41
2:B:185:PRO:HA	2:B:221:HIS:O	2.21	0.40
1:A:135:ILE:CG2	1:A:136:PRO:CD	2.77	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	158/184 (86%)	144 (91%)	12 (8%)	2 (1%)	12	4
2	B	159/165 (96%)	151 (95%)	8 (5%)	0	100	100
All	All	317/349 (91%)	295 (93%)	20 (6%)	2 (1%)	25	15

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	32	THR
1	A	76	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	147/160 (92%)	135 (92%)	12 (8%)	11	4
2	B	136/139 (98%)	126 (93%)	10 (7%)	13	6
All	All	283/299 (95%)	261 (92%)	22 (8%)	12	5

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	GLN
1	A	33	THR
1	A	36	GLN
1	A	90	GLN
1	A	98	THR
1	A	128	THR
1	A	139	LEU
1	A	140	GLN
1	A	152	LEU
1	A	164	LEU
1	A	169	ARG
1	A	178	SER
2	B	110	LYS
2	B	111	THR
2	B	146	LEU
2	B	150	ARG
2	B	159	MET
2	B	190	LEU
2	B	213	LEU
2	B	230	ASN
2	B	247	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	251	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	7	HIS
1	A	31	GLN
1	A	77	HIS
1	A	140	GLN
1	A	144	HIS
1	A	150	GLN
2	B	112	HIS
2	B	210	ASN
2	B	230	ASN
2	B	252	GLN
2	B	259	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 5 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PO4	A	201	-	4,4,4	0.55	0	6,6,6	1.23	1 (16%)
3	PO4	A	202	-	4,4,4	1.15	1 (25%)	6,6,6	2.33	2 (33%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	202	PO4	P-O1	2.21	1.56	1.50

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	202	PO4	O4-P-O3	4.50	122.42	107.97
3	A	201	PO4	O4-P-O1	-2.26	102.63	110.89
3	A	202	PO4	O4-P-O1	-2.14	103.08	110.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	164/184 (89%)	0.50	13 (7%) <b>12</b> <b>14</b>	22, 43, 89, 135	0
2	B	159/165 (96%)	0.64	10 (6%) <b>20</b> <b>22</b>	20, 46, 75, 99	0
All	All	323/349 (92%)	0.57	23 (7%) <b>16</b> <b>17</b>	20, 44, 81, 135	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	32	THR	5.6
1	A	77	HIS	5.5
1	A	31	GLN	4.9
2	B	154	GLY	4.5
2	B	246	ALA	4.5
1	A	135	ILE	4.2
1	A	122	VAL	4.1
1	A	138	LYS	3.4
1	A	173	LEU	3.1
1	A	47	LYS	3.0
2	B	249	ARG	2.9
2	B	109	ARG	2.8
2	B	180	LEU	2.8
1	A	58	ILE	2.7
1	A	48	GLY	2.6
2	B	155	GLU	2.5
1	A	49	PHE	2.4
2	B	214	VAL	2.3
2	B	188	PRO	2.3
2	B	194	ILE	2.1
1	A	24	VAL	2.1
1	A	136	PRO	2.0
2	B	196	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	PO4	A	202	5/5	0.85	0.21	53,55,58,61	0
3	PO4	A	201	5/5	0.94	0.12	48,51,52,54	0
5	CA	B	305	1/1	0.95	0.07	56,56,56,56	0
4	ZN	B	302	1/1	0.99	0.10	42,42,42,42	0
5	CA	B	303	1/1	0.99	0.13	33,33,33,33	0
5	CA	B	304	1/1	0.99	0.09	49,49,49,49	0
4	ZN	B	301	1/1	1.00	0.12	22,22,22,22	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.