



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 10, 2020 – 01:28 PM BST

PDB ID : 3ZP6
Title : INFLUENZA VIRUS (VN1194) H5 E190D mutant HA with LSTc
Authors : Liu, J.; Stevens, D.J.; Gamblin, S.J.; Skehel, J.J.
Deposited on : 2013-02-26
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	340	
2	F	160	
3	A	2	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4007 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HAEMAGGLUTININ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	E	321	2548	1610	440	483	15	0	0	0


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	40	LYS	THR	conflict	UNP Q6DQ34
E	190	ASP	GLU	engineered mutation	UNP Q6DQ34

- Molecule 2 is a protein called HAEMAGGLUTININ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	F	158	1272	791	221	252	8	0	0	0

- Molecule 3 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	A	2	25	14	1	10	0	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	F	1	14	8	1	5	0	0
4	F	1	14	8	1	5	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	E	72	72	72	0	0
5	F	62	62	62	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HAEMAGGLUTININ



- Molecule 2: HAEMAGGLUTININ



- Molecule 3: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	101.39Å 101.39Å 450.53Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.60 29.95 – 2.52	Depositor EDS
% Data completeness (in resolution range)	99.1 (30.00-2.60) 91.8 (29.95-2.52)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.72 (at 2.51Å)	Xtrriage
Refinement program	REFMAC 5	Depositor
R, R_{free}	0.223 , (Not available) 0.245 , 0.267	Depositor DCC
R_{free} test set	1438 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	37.5	Xtrriage
Anisotropy	0.030	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 56.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	4007	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SIA, GAL, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	E	0.38	0/2610	0.54	0/3545
2	F	0.45	0/1296	0.54	0/1742
All	All	0.40	0/3906	0.54	0/5287

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	2548	0	2490	57	1
2	F	1272	0	1186	15	0
3	A	25	0	19	1	0
4	F	28	0	26	0	0
5	E	72	0	0	6	0
5	F	62	0	0	2	0
All	All	4007	0	3721	68	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:168:TYR:HD1	1:E:169:ASN:N	1.51	1.08
1:E:159:SER:O	1:E:196:GLN:OE1	1.83	0.97
1:E:57:ARG:O	5:E:2026:HOH:O	1.83	0.94
1:E:168:TYR:CD1	1:E:169:ASN:N	2.36	0.93
1:E:168:TYR:C	1:E:168:TYR:HD1	1.75	0.90
1:E:120:ILE:HG22	1:E:168:TYR:CE2	2.09	0.87
1:E:120:ILE:HG22	1:E:168:TYR:HE2	1.40	0.85
1:E:168:TYR:CE1	1:E:169:ASN:O	2.30	0.84
1:E:159:SER:C	1:E:196:GLN:OE1	2.17	0.83
1:E:168:TYR:HE1	1:E:169:ASN:O	1.62	0.81
1:E:299:HIS:HD2	1:E:301:LEU:H	1.30	0.80
1:E:168:TYR:CD1	1:E:168:TYR:C	2.51	0.79
2:F:19:ASP:HB3	2:F:36:ALA:HB2	1.66	0.78
1:E:72:ASP:O	1:E:75:ILE:HG12	1.86	0.76
1:E:110:SER:HB3	5:E:2042:HOH:O	1.84	0.75
1:E:60:SER:OG	1:E:92:ASP:OD1	2.04	0.75
1:E:128:SER:HB3	1:E:162:PRO:HG2	1.73	0.69
1:E:168:TYR:CE1	1:E:169:ASN:C	2.67	0.67
1:E:9:ILE:HD11	2:F:122:VAL:HG21	1.78	0.66
1:E:56:LEU:O	1:E:57:ARG:HB2	1.96	0.65
1:E:120:ILE:HB	1:E:168:TYR:HD2	1.62	0.65
1:E:92:ASP:O	5:E:2036:HOH:O	2.15	0.64
1:E:190:ASP:OD2	3:A:2:SIA:H92	1.99	0.63
1:E:170:ASN:OD1	1:E:172:ASN:ND2	2.26	0.62
1:E:168:TYR:HE1	1:E:169:ASN:C	2.02	0.62
1:E:120:ILE:HB	1:E:168:TYR:CD2	2.35	0.61
1:E:169:ASN:OD1	1:E:169:ASN:C	2.39	0.60
1:E:299:HIS:CD2	1:E:301:LEU:H	2.18	0.59
1:E:113:ASN:OD1	1:E:264:GLY:HA3	2.03	0.57
1:E:134:GLY:HA3	1:E:153:TRP:HB3	1.87	0.57
1:E:168:TYR:HE1	1:E:170:ASN:HA	1.68	0.57
1:E:173:GLN:NE2	1:E:173:GLN:HA	2.21	0.55
1:E:120:ILE:CG2	1:E:168:TYR:CE2	2.87	0.55
2:F:157:TYR:O	2:F:158:ASP:HB2	2.07	0.55
2:F:156:THR:HG22	2:F:156:THR:O	2.08	0.54
1:E:115:PHE:HE1	1:E:260:ILE:HG12	1.72	0.54
1:E:95:TYR:CD1	1:E:230:MET:HG2	2.44	0.53
1:E:168:TYR:CE1	1:E:170:ASN:HA	2.44	0.52
1:E:151:VAL:HB	1:E:252:ILE:HG22	1.91	0.52
1:E:20:VAL:HG21	1:E:318:ALA:HB2	1.93	0.51
1:E:164:ILE:O	1:E:246:GLU:HA	2.11	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:168:TYR:CD1	1:E:169:ASN:C	2.84	0.50
1:E:169:ASN:HA	1:E:242:ALA:HA	1.94	0.50
2:F:126:LEU:O	2:F:129:ASN:HB2	2.11	0.50
1:E:62:ALA:HB3	1:E:92:ASP:OD1	2.12	0.49
1:E:285:PRO:HD3	1:E:301:LEU:O	2.13	0.48
1:E:97:GLY:HA3	1:E:230:MET:O	2.14	0.48
1:E:309:TYR:HD2	2:F:89:LEU:HD22	1.77	0.48
1:E:58:ASP:HB3	1:E:90:VAL:HG22	1.94	0.48
2:F:19:ASP:HB2	5:F:2012:HOH:O	2.13	0.48
1:E:115:PHE:HE1	1:E:260:ILE:CG1	2.27	0.47
2:F:75:ARG:NH1	2:F:78:GLU:OE1	2.47	0.45
1:E:291:SER:HB2	5:E:2021:HOH:O	2.17	0.45
1:E:9:ILE:HG13	2:F:119:TYR:HA	1.98	0.45
1:E:120:ILE:HD11	1:E:254:PRO:HB2	2.00	0.44
1:E:129:HIS:HE1	5:E:2050:HOH:O	2.00	0.44
1:E:324:SER:HA	5:E:2069:HOH:O	2.18	0.44
2:F:72:ASN:HD22	2:F:75:ARG:HH21	1.65	0.44
1:E:203:SER:OG	1:E:246:GLU:HB3	2.18	0.43
1:E:169:ASN:HB2	1:E:242:ALA:HB2	2.00	0.43
2:F:72:ASN:ND2	2:F:75:ARG:HH21	2.17	0.43
2:F:101:LEU:HD13	5:F:2035:HOH:O	2.19	0.43
2:F:68:ARG:NH1	2:F:81:ASN:OD1	2.52	0.43
2:F:126:LEU:HD23	2:F:126:LEU:N	2.33	0.42
1:E:214:VAL:HA	1:E:215:PRO:HD3	1.88	0.41
1:E:115:PHE:CE1	1:E:260:ILE:HG12	2.54	0.41
1:E:168:TYR:HD1	1:E:169:ASN:CA	2.26	0.40
1:E:295:PHE:HE1	2:F:62:GLN:OE1	2.05	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:259:LYS:NZ	1:E:262:LYS:NZ[11_565]	1.65	0.55

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	319/340 (94%)	306 (96%)	12 (4%)	1 (0%)	41	64
2	F	156/160 (98%)	153 (98%)	3 (2%)	0	100	100
All	All	475/500 (95%)	459 (97%)	15 (3%)	1 (0%)	47	71

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	198	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	289/307 (94%)	272 (94%)	17 (6%)	19	39
2	F	134/136 (98%)	128 (96%)	6 (4%)	27	52
All	All	423/443 (96%)	400 (95%)	23 (5%)	22	44

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	49	ASP
1	E	78	PRO
1	E	168	TYR
1	E	169	ASN
1	E	190	ASP
1	E	196	GLN
1	E	220	ARG
1	E	226	GLN
1	E	262	LYS
1	E	263	LYS
1	E	274	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	278	CYS
1	E	284	THR
1	E	297	ASN
1	E	301	LEU
1	E	315	LEU
1	E	321	LEU
2	F	57	ASP
2	F	58	LYS
2	F	101	LEU
2	F	128	ASP
2	F	133	LEU
2	F	158	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	19	GLN
1	E	173	GLN
1	E	226	GLN
1	E	297	ASN
1	E	299	HIS
2	F	26	HIS
2	F	62	GLN
2	F	72	ASN
2	F	125	GLN
2	F	146	ASN
2	F	154	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GAL	A	1	3	3,4,12	0.39	0	1,4,17	0.64	0
3	SIA	A	2	3	17,20,21	0.48	0	21,28,31	0.93	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GAL	A	1	3	-	1/2/2/22	-
3	SIA	A	2	3	-	0/14/34/38	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	2	SIA	C6-O6-C2	2.05	115.72	111.34

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1	GAL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	2	SIA	1	0

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	F	411	1	14,14,15	0.41	0	17,19,21	0.75	0
4	NAG	F	421	1	14,14,15	0.44	0	17,19,21	1.02	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	F	411	1	-	2/6/23/26	0/1/1/1
4	NAG	F	421	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	F	421	NAG	C4-C3-C2	-2.02	108.06	111.02

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	F	411	NAG	O5-C5-C6-O6
4	F	411	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	E	321/340 (94%)	0.96	55 (17%) 1 1	26, 70, 95, 100	0
2	F	158/160 (98%)	0.56	17 (10%) 5 3	15, 45, 71, 95	0
All	All	479/500 (95%)	0.83	72 (15%) 2 1	15, 61, 93, 100	0

All (72) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	192	THR	7.3
1	E	158	ASN	7.3
2	F	148	CYS	5.4
1	E	160	THR	5.0
1	E	189	ALA	4.8
2	F	142	HIS	4.7
2	F	60	ASN	4.6
2	F	151	SER	4.6
1	E	171	THR	4.3
1	E	75	ILE	4.2
1	E	251	PHE	4.2
2	F	154	ASN	4.0
1	E	159	SER	3.9
2	F	158	ASP	3.8
1	E	252	ILE	3.8
2	F	57	ASP	3.7
1	E	190	ASP	3.7
1	E	66	LEU	3.7
1	E	125	SER	3.6
1	E	179	LEU	3.5
1	E	325	PRO	3.5
2	F	155	GLY	3.2
2	F	147	GLU	3.2
1	E	49	ASP	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	50	GLY	3.0
1	E	134	GLY	3.0
1	E	242	ALA	3.0
1	E	51	VAL	2.9
2	F	129	ASN	2.9
1	E	178	VAL	2.8
1	E	254	PRO	2.8
1	E	187	ASP	2.8
1	E	241	ASP	2.7
1	E	127	SER	2.7
2	F	128	ASP	2.7
1	E	133	LEU	2.7
1	E	164	ILE	2.7
1	E	157	LYS	2.7
1	E	278	CYS	2.7
1	E	305	GLU	2.7
1	E	147	PHE	2.7
2	F	143	LYS	2.7
1	E	76	ASN	2.7
1	E	47	ASP	2.6
1	E	141	TYR	2.6
1	E	253	ALA	2.6
1	E	169	ASN	2.6
1	E	177	LEU	2.6
1	E	151	VAL	2.6
1	E	130	GLU	2.6
1	E	124	SER	2.5
2	F	150	GLU	2.5
1	E	170	ASN	2.5
1	E	126	TRP	2.5
1	E	201	TYR	2.5
1	E	153	TRP	2.4
1	E	154	LEU	2.3
1	E	144	LYS	2.3
1	E	196	GLN	2.3
1	E	258	TYR	2.3
2	F	141	TYR	2.3
1	E	182	ILE	2.3
1	E	243	ILE	2.2
1	E	240	ASN	2.2
1	E	225	GLY	2.2
2	F	152	VAL	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	139	CYS	2.2
1	E	277	ASN	2.1
2	F	126	LEU	2.1
1	E	232	PHE	2.1
1	E	180	TRP	2.1
2	F	50	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GAL	A	1	5/12	0.51	0.32	130,130,131,131	0
3	SIA	A	2	20/21	0.67	0.33	130,130,130,130	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	F	421	14/15	0.80	0.39	73,73,74,74	0
4	NAG	F	411	14/15	0.82	0.48	79,80,80,80	0

6.5 Other polymers [i](#)

There are no such residues in this entry.