



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 12, 2023 – 06:32 pm GMT

PDB ID : 4B7X
Title : Crystal structure of hypothetical protein PA1648 from *Pseudomonas aeruginosa*.
Authors : Alpey, M.S.; McMahon, S.A.; Duthie, F.; Naismith, J.H.
Deposited on : 2012-08-24
Resolution : 2.20 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	336	
1	B	336	
1	C	336	
1	D	336	
1	E	336	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	336	<p>5% 88% 10% .</p>
1	G	336	<p>12% 84% 11% . .</p>
1	H	336	<p>4% 84% 13% .</p>
1	I	336	<p>22% 60% 9% . 30%</p>
1	J	336	<p>7% 84% 13% . .</p>
1	K	336	<p>16% 71% 18% . 10%</p>
1	L	336	<p>8% 79% 15% 5%</p>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 30208 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROBABLE OXIDOREDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	329	2504	1597	431	463	13	0	0	0
1	B	331	2521	1608	434	466	13	0	0	0
1	C	325	2473	1581	423	456	13	0	0	0
1	D	327	2490	1590	428	459	13	0	0	0
1	E	331	2521	1608	434	466	13	0	0	0
1	F	331	2521	1608	434	466	13	0	0	0
1	G	321	2446	1564	417	452	13	0	0	0
1	H	327	2489	1591	426	459	13	0	0	0
1	I	234	1768	1129	303	324	12	0	0	0
1	J	327	2486	1586	428	459	13	0	0	0
1	K	302	2316	1484	399	421	12	0	0	0
1	L	320	2443	1563	417	450	13	0	0	0

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP Q9I377
A	0	ALA	-	expression tag	UNP Q9I377
B	-1	GLY	-	expression tag	UNP Q9I377
B	0	ALA	-	expression tag	UNP Q9I377
C	-1	GLY	-	expression tag	UNP Q9I377

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	ALA	-	expression tag	UNP Q9I377
D	-1	GLY	-	expression tag	UNP Q9I377
D	0	ALA	-	expression tag	UNP Q9I377
E	-1	GLY	-	expression tag	UNP Q9I377
E	0	ALA	-	expression tag	UNP Q9I377
F	-1	GLY	-	expression tag	UNP Q9I377
F	0	ALA	-	expression tag	UNP Q9I377
G	-1	GLY	-	expression tag	UNP Q9I377
G	0	ALA	-	expression tag	UNP Q9I377
H	-1	GLY	-	expression tag	UNP Q9I377
H	0	ALA	-	expression tag	UNP Q9I377
I	-1	GLY	-	expression tag	UNP Q9I377
I	0	ALA	-	expression tag	UNP Q9I377
J	-1	GLY	-	expression tag	UNP Q9I377
J	0	ALA	-	expression tag	UNP Q9I377
K	-1	GLY	-	expression tag	UNP Q9I377
K	0	ALA	-	expression tag	UNP Q9I377
L	-1	GLY	-	expression tag	UNP Q9I377
L	0	ALA	-	expression tag	UNP Q9I377

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: C₂₁H₂₈N₇O₁₇P₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	48	21	7	17	3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	B	1	Total 48	C 21	N 7	O 17	P 3	0	0
2	C	1	Total 48	C 21	N 7	O 17	P 3	0	0
2	D	1	Total 48	C 21	N 7	O 17	P 3	0	0
2	E	1	Total 48	C 21	N 7	O 17	P 3	0	0
2	F	1	Total 48	C 21	N 7	O 17	P 3	0	0
2	G	1	Total 48	C 21	N 7	O 17	P 3	0	0
2	H	1	Total 48	C 21	N 7	O 17	P 3	0	0
2	I	1	Total 48	C 21	N 7	O 17	P 3	0	0
2	J	1	Total 48	C 21	N 7	O 17	P 3	0	0
2	K	1	Total 48	C 21	N 7	O 17	P 3	0	0
2	L	1	Total 48	C 21	N 7	O 17	P 3	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	118	Total 118	O 118	0	0
3	B	68	Total 68	O 68	0	0
3	C	46	Total 46	O 46	0	0
3	D	60	Total 60	O 60	0	0
3	E	81	Total 81	O 81	0	0
3	F	67	Total 67	O 67	0	0
3	G	43	Total 43	O 43	0	0
3	H	41	Total 41	O 41	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	I	20	Total 20	O 20	0	0
3	J	37	Total 37	O 37	0	0
3	K	30	Total 30	O 30	0	0
3	L	43	Total 43	O 43	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PROBABLE OXIDOREDUCTASE




- Molecule 1: PROBABLE OXIDOREDUCTASE



- Molecule 1: PROBABLE OXIDOREDUCTASE




- Molecule 1: PROBABLE OXIDOREDUCTASE









● Molecule 1: PROBABLE OXIDOREDUCTASE





● Molecule 1: PROBABLE OXIDOREDUCTASE




● Molecule 1: PROBABLE OXIDOREDUCTASE



● Molecule 1: PROBABLE OXIDOREDUCTASE





4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	169.26Å 175.97Å 182.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	56.42 – 2.20 55.83 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.8 (56.42-2.20) 99.8 (55.83-2.20)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.79 (at 2.20Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.237 , 0.276 0.234 , 0.272	Depositor DCC
R_{free} test set	13815 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	34.8	Xtrriage
Anisotropy	0.421	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 44.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.013 for -h,l,k	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	30208	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.56	2/2551 (0.1%)	0.67	0/3443
1	B	0.50	1/2569 (0.0%)	0.63	0/3469
1	C	0.44	0/2518	0.59	0/3397
1	D	0.48	1/2537 (0.0%)	0.61	0/3425
1	E	0.53	1/2569 (0.0%)	0.67	0/3469
1	F	0.49	0/2569	0.60	0/3469
1	G	0.44	2/2491 (0.1%)	0.58	0/3362
1	H	0.42	1/2536 (0.0%)	0.56	0/3423
1	I	0.42	2/1795 (0.1%)	0.53	0/2416
1	J	0.43	1/2533 (0.0%)	0.58	0/3420
1	K	0.44	2/2357 (0.1%)	0.56	0/3177
1	L	0.44	1/2488 (0.0%)	0.58	0/3356
All	All	0.47	14/29513 (0.0%)	0.60	0/39826

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	295	TRP	CD2-CE2	5.67	1.48	1.41
1	K	295	TRP	CD2-CE2	5.32	1.47	1.41
1	E	55	TRP	CD2-CE2	5.20	1.47	1.41
1	B	295	TRP	CD2-CE2	5.18	1.47	1.41
1	A	55	TRP	CD2-CE2	5.18	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2504	0	2524	16	0
1	B	2521	0	2544	14	0
1	C	2473	0	2499	24	0
1	D	2490	0	2513	22	0
1	E	2521	0	2544	18	0
1	F	2521	0	2544	25	0
1	G	2446	0	2469	24	0
1	H	2489	0	2515	29	0
1	I	1768	0	1794	18	0
1	J	2486	0	2505	35	0
1	K	2316	0	2334	33	0
1	L	2443	0	2467	28	0
2	A	48	0	25	1	0
2	B	48	0	25	4	0
2	C	48	0	25	4	0
2	D	48	0	25	3	0
2	E	48	0	25	2	0
2	F	48	0	25	3	0
2	G	48	0	25	4	0
2	H	48	0	25	2	0
2	I	48	0	25	4	0
2	J	48	0	25	3	0
2	K	48	0	25	2	0
2	L	48	0	25	2	0
3	A	118	0	0	0	0
3	B	68	0	0	0	0
3	C	46	0	0	1	0
3	D	60	0	0	0	0
3	E	81	0	0	0	0
3	F	67	0	0	0	0
3	G	43	0	0	0	0
3	H	41	0	0	0	0
3	I	20	0	0	0	0
3	J	37	0	0	1	0
3	K	30	0	0	0	0
3	L	43	0	0	0	0
All	All	30208	0	29552	264	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 264 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:400:NAP:H4B	2:D:400:NAP:O2A	1.26	1.02
1:F:16:SER:O	1:F:254:LYS:HG3	1.65	0.96
1:C:48:LEU:HD11	1:C:332:LEU:HB2	1.51	0.91
2:D:400:NAP:O2A	2:D:400:NAP:C4B	2.19	0.84
1:F:206:ALA:O	1:F:210:LYS:HG3	1.80	0.81

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	325/336 (97%)	322 (99%)	3 (1%)	0	100	100
1	B	329/336 (98%)	324 (98%)	5 (2%)	0	100	100
1	C	319/336 (95%)	310 (97%)	9 (3%)	0	100	100
1	D	323/336 (96%)	316 (98%)	6 (2%)	1 (0%)	41	46
1	E	329/336 (98%)	323 (98%)	6 (2%)	0	100	100
1	F	329/336 (98%)	323 (98%)	6 (2%)	0	100	100
1	G	317/336 (94%)	307 (97%)	9 (3%)	1 (0%)	41	46
1	H	323/336 (96%)	313 (97%)	10 (3%)	0	100	100
1	I	226/336 (67%)	216 (96%)	10 (4%)	0	100	100
1	J	323/336 (96%)	314 (97%)	8 (2%)	1 (0%)	41	46
1	K	292/336 (87%)	277 (95%)	12 (4%)	3 (1%)	15	14
1	L	314/336 (94%)	308 (98%)	6 (2%)	0	100	100
All	All	3749/4032 (93%)	3653 (97%)	90 (2%)	6 (0%)	47	55

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	21	ARG
1	K	262	ASN
1	K	261	ALA
1	G	258	ARG
1	K	109	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	258/263 (98%)	253 (98%)	5 (2%)	57	71
1	B	260/263 (99%)	254 (98%)	6 (2%)	50	63
1	C	255/263 (97%)	249 (98%)	6 (2%)	49	62
1	D	257/263 (98%)	251 (98%)	6 (2%)	50	63
1	E	260/263 (99%)	255 (98%)	5 (2%)	57	71
1	F	260/263 (99%)	257 (99%)	3 (1%)	71	83
1	G	252/263 (96%)	247 (98%)	5 (2%)	55	69
1	H	257/263 (98%)	250 (97%)	7 (3%)	44	57
1	I	180/263 (68%)	175 (97%)	5 (3%)	43	56
1	J	256/263 (97%)	249 (97%)	7 (3%)	44	57
1	K	239/263 (91%)	231 (97%)	8 (3%)	38	49
1	L	252/263 (96%)	243 (96%)	9 (4%)	35	45
All	All	2986/3156 (95%)	2914 (98%)	72 (2%)	49	62

5 of 72 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	K	118	ARG
1	L	327	PHE
1	K	230	ASP
1	L	121	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	53	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	J	8	GLN
1	J	250	GLN
1	L	326	ASN
1	K	8	GLN
1	G	8	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAP	K	400	-	45,52,52	1.62	8 (17%)	56,80,80	1.29	7 (12%)
2	NAP	B	400	-	45,52,52	1.59	7 (15%)	56,80,80	1.26	6 (10%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAP	C	400	-	45,52,52	1.45	7 (15%)	56,80,80	1.28	6 (10%)
2	NAP	F	400	-	45,52,52	1.54	7 (15%)	56,80,80	1.25	5 (8%)
2	NAP	A	400	-	45,52,52	1.52	8 (17%)	56,80,80	1.21	7 (12%)
2	NAP	H	400	-	45,52,52	1.59	7 (15%)	56,80,80	1.17	2 (3%)
2	NAP	L	400	-	45,52,52	1.40	5 (11%)	56,80,80	1.35	7 (12%)
2	NAP	E	400	-	45,52,52	1.49	6 (13%)	56,80,80	1.35	8 (14%)
2	NAP	G	400	-	45,52,52	1.42	6 (13%)	56,80,80	1.20	4 (7%)
2	NAP	I	400	-	45,52,52	1.50	9 (20%)	56,80,80	1.20	5 (8%)
2	NAP	J	400	-	45,52,52	1.42	8 (17%)	56,80,80	1.27	5 (8%)
2	NAP	D	400	-	45,52,52	1.93	8 (17%)	56,80,80	1.44	7 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	K	400	-	-	9/31/67/67	0/5/5/5
2	NAP	B	400	-	-	8/31/67/67	0/5/5/5
2	NAP	C	400	-	-	8/31/67/67	0/5/5/5
2	NAP	F	400	-	-	7/31/67/67	0/5/5/5
2	NAP	A	400	-	-	7/31/67/67	0/5/5/5
2	NAP	H	400	-	-	9/31/67/67	0/5/5/5
2	NAP	L	400	-	-	11/31/67/67	0/5/5/5
2	NAP	E	400	-	-	9/31/67/67	0/5/5/5
2	NAP	G	400	-	-	7/31/67/67	0/5/5/5
2	NAP	I	400	-	-	6/31/67/67	0/5/5/5
2	NAP	J	400	-	-	7/31/67/67	0/5/5/5
2	NAP	D	400	-	-	11/31/67/67	0/5/5/5

The worst 5 of 86 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	400	NAP	O4B-C1B	6.34	1.49	1.41
2	D	400	NAP	C2N-N1N	5.29	1.41	1.35
2	K	400	NAP	C2N-N1N	5.18	1.41	1.35
2	B	400	NAP	O4D-C1D	5.11	1.48	1.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	400	NAP	O4D-C1D	4.94	1.48	1.41

The worst 5 of 69 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	400	NAP	N3A-C2A-N1A	-5.10	120.70	128.68
2	C	400	NAP	N3A-C2A-N1A	-4.75	121.25	128.68
2	K	400	NAP	N3A-C2A-N1A	-4.73	121.29	128.68
2	I	400	NAP	N3A-C2A-N1A	-4.55	121.57	128.68
2	D	400	NAP	N3A-C2A-N1A	-4.53	121.59	128.68

There are no chirality outliers.

5 of 99 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	400	NAP	O4D-C1D-N1N-C6N
2	B	400	NAP	O4D-C1D-N1N-C6N
2	C	400	NAP	O4D-C1D-N1N-C6N
2	D	400	NAP	C5B-O5B-PA-O1A
2	D	400	NAP	C5B-O5B-PA-O2A


There are no ring outliers.


12 monomers are involved in 34 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	K	400	NAP	2	0
2	B	400	NAP	4	0
2	C	400	NAP	4	0
2	F	400	NAP	3	0
2	A	400	NAP	1	0
2	H	400	NAP	2	0
2	L	400	NAP	2	0
2	E	400	NAP	2	0
2	G	400	NAP	4	0
2	I	400	NAP	4	0
2	J	400	NAP	3	0
2	D	400	NAP	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will


also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
































5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	329/336 (97%)	-0.15	2 (0%) 89 88	14, 23, 42, 62	0
1	B	331/336 (98%)	-0.10	1 (0%) 94 93	20, 33, 51, 61	0
1	C	325/336 (96%)	0.47	21 (6%) 18 17	23, 45, 79, 98	0
1	D	327/336 (97%)	0.25	22 (6%) 17 16	20, 33, 83, 109	0
1	E	331/336 (98%)	-0.00	9 (2%) 54 52	14, 26, 55, 87	0
1	F	331/336 (98%)	0.06	16 (4%) 30 29	25, 37, 59, 77	0
1	G	321/336 (95%)	0.52	39 (12%) 4 3	23, 43, 96, 120	0
1	H	327/336 (97%)	0.34	14 (4%) 35 33	27, 49, 69, 92	0
1	I	234/336 (69%)	1.49	75 (32%) 0 0	31, 63, 122, 140	0
1	J	327/336 (97%)	0.36	24 (7%) 15 14	25, 47, 81, 109	0
1	K	302/336 (89%)	0.92	53 (17%) 1 1	27, 55, 91, 112	0
1	L	320/336 (95%)	0.54	27 (8%) 11 9	27, 47, 77, 105	0
All	All	3805/4032 (94%)	0.36	303 (7%) 12 11	14, 40, 84, 140	0

The worst 5 of 303 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	315	PRO	11.9
1	K	5	ILE	9.1
1	I	312	GLU	7.8
1	D	16	SER	7.5
1	J	21	ARG	7.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAP	L	400	48/48	0.90	0.18	35,48,63,65	0
2	NAP	K	400	48/48	0.92	0.14	38,49,60,61	0
2	NAP	I	400	48/48	0.92	0.14	47,59,65,70	0
2	NAP	G	400	48/48	0.95	0.15	30,48,64,71	0
2	NAP	C	400	48/48	0.95	0.12	25,36,40,44	0
2	NAP	J	400	48/48	0.95	0.13	30,48,58,61	0
2	NAP	D	400	48/48	0.95	0.13	28,35,51,53	0
2	NAP	F	400	48/48	0.95	0.12	26,31,38,40	0
2	NAP	E	400	48/48	0.96	0.13	19,32,38,40	0
2	NAP	H	400	48/48	0.96	0.11	33,40,43,46	0
2	NAP	B	400	48/48	0.96	0.11	25,30,33,34	0
2	NAP	A	400	48/48	0.97	0.11	15,21,26,26	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NAP L 400:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAP K 400:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAP I 400:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAP G 400:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAP C 400:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAP J 400:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around NAP D 400:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAP F 400:**




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)








Electron density around NAP B 400:

$2mF_o-DF_c$ (at 0.7 rnsd) in gray
 mF_o-DF_c (at 3 rnsd) in purple (negative)
and green (positive)

**Electron density around NAP A 400:**

$2mF_o-DF_c$ (at 0.7 rnsd) in gray
 mF_o-DF_c (at 3 rnsd) in purple (negative)
and green (positive)



6.5 Other polymers

There are no such residues in this entry.