

wwPDB X-ray Structure Validation Summary Report (i)

May 13, 2020 – 01:23 pm BST

PDB ID	:	4CMB
Title	:	Crystal structure of pteridine reductase 1 (PTR1) from Trypanosoma brucei
		in ternary complex with cofactor and inhibitor
Authors	:	Barrack, K.L.; Hunter, W.N.
Deposited on		
$\operatorname{Resolution}$:	1.96 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
$\operatorname{CCP4}$:	$7.0.044 (\mathrm{Gargrove})$
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} {f Whole archive}\ (\#{f Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries},{ m resolution\ range}({ m \AA}))$
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539(1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	288	4%	8%	13%
1	В	288	80%	7%	13%
1	С	288	75%	11%	14%
1	D	288	78%	8% •	14%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 8567 atoms, of which 0 are hydrogens and 0 are deuteriums.


In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	Δ	251	Total	С	Ν	Ο	S	0	7	0
	A	201	1907	1199	335	362	11	0		
1	В	250	Total	С	Ν	Ο	S	0	1	0
	I D	230	1868	1174	329	353	12	0	1	0
1	С	249	Total	С	Ν	Ο	S	0	3	0
		249	1867	1176	327	352	12	0	່ <u>ບ</u>	0
1	П	249	Total	С	Ν	Ο	S	0	2	0
	I D	249	1866	1173	328	354	11			0

• Molecule 1 is a protein called PTERIDINE REDUCTASE 1.

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	-19	MET	-	expression tag	UNP 076290
A	-18	GLY	-	expression tag	UNP 076290
A	-17	SER	-	expression tag	UNP 076290
А	-16	SER	-	expression tag	UNP 076290
A	-15	HIS	-	expression tag	UNP 076290
A	-14	HIS	-	expression tag	UNP 076290
А	-13	HIS	-	expression tag	UNP 076290
A	-12	HIS	-	expression tag	UNP 076290
А	-11	HIS	-	expression tag	UNP 076290
A	-10	HIS	-	expression tag	UNP 076290
А	-9	SER	-	expression tag	UNP 076290
A	-8	SER	-	expression tag	UNP 076290
А	-7	GLY	-	expression tag	UNP 076290
А	-6	LEU	-	expression tag	UNP 076290
A	-5	VAL	-	expression tag	UNP 076290
A	-4	PRO	-	expression tag	UNP 076290
A	-3	ARG	-	expression tag	UNP 076290
А	-2	GLY	-	expression tag	UNP 076290
А	-1	SER	-	expression tag	UNP 076290
А	0	HIS	-	expression tag	UNP 076290
В	-19	MET	-	expression tag	UNP 076290



	ed from pre	Modelled	Actual	Comment	Reference
B	-18	GLY	-	expression tag	UNP O76290
B	-17	SER	_	expression tag	UNP 076290
B	-16	SER	_	expression tag	UNP 076290
B	-15	HIS	_	expression tag	UNP 076290
B	-14	HIS	_	expression tag	UNP 076290
B	-13	HIS	_	expression tag	UNP 076290
B	-12	HIS	_	expression tag	UNP 076290
B	-11	HIS	_	expression tag	UNP 076290
В	-10	HIS	_	expression tag	UNP 076290
В	-9	SER	-	expression tag	UNP 076290
В	-8	SER	_	expression tag	UNP 076290
В	-7	GLY	-	expression tag	UNP 076290
В	-6	LEU	-	expression tag	UNP 076290
В	-5	VAL	-	expression tag	UNP 076290
В	-4	PRO	_	expression tag	UNP 076290
В	-3	ARG	_	expression tag	UNP 076290
В	-2	GLY	_	expression tag	UNP 076290
В	-1	SER	-	expression tag	UNP 076290
В	0	HIS	-	expression tag	UNP 076290
С	-19	MET	-	expression tag	UNP 076290
С	-18	GLY	-	expression tag	UNP 076290
С	-17	SER	-	expression tag	UNP 076290
С	-16	SER	-	expression tag	UNP 076290
С	-15	HIS	-	expression tag	UNP 076290
С	-14	HIS	-	expression tag	UNP 076290
С	-13	HIS	-	expression tag	UNP 076290
С	-12	HIS	-	expression tag	UNP 076290
С	-11	HIS	-	expression tag	UNP 076290
С	-10	HIS	-	expression tag	UNP 076290
С	-9	SER	-	expression tag	UNP 076290
С	-8	SER	-	expression tag	UNP 076290
С	-7	GLY	-	expression tag	UNP 076290
С	-6	LEU	-	expression tag	UNP 076290
С	-5	VAL	-	expression tag	UNP 076290
С	-4	PRO	-	expression tag	UNP 076290
С	-3	ARG	-	expression tag	UNP 076290
С	-2	GLY	-	expression tag	UNP 076290
С	-1	SER	-	expression tag	UNP 076290
С	0	HIS	-	expression tag	UNP 076290
D	-19	MET	-	expression tag	UNP 076290
D	-18	GLY	-	expression tag	UNP 076290
D	-17	SER	-	expression tag	UNP 076290

Continued on next page...




4CMB

Chain	Residue	Modelled	Actual	Comment	Reference
D	-16	SER	-	expression tag	UNP 076290
D	-15	HIS	-	expression tag	UNP 076290
D	-14	HIS	-	expression tag	UNP 076290
D	-13	HIS	-	expression tag	UNP 076290
D	-12	HIS	-	expression tag	UNP 076290
D	-11	HIS	-	expression tag	UNP 076290
D	-10	HIS	-	expression tag	UNP 076290
D	-9	SER	-	expression tag	UNP 076290
D	-8	SER	-	expression tag	UNP 076290
D	-7	GLY	-	expression tag	UNP 076290
D	-6	LEU	-	expression tag	UNP 076290
D	-5	VAL	-	expression tag	UNP 076290
D	-4	PRO	-	expression tag	UNP 076290
D	-3	ARG	-	expression tag	UNP 076290
D	-2	GLY	-	expression tag	UNP 076290
D	-1	SER	-	expression tag	UNP 076290
D	0	HIS	-	expression tag	UNP 076290

Continued from previous page...

• Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
0	Δ	1	Total	С	Ν	Ο	Р	0	0
		L	48	21	7	17	3	0	0
0	D	1	Total	С	Ν	Ο	Р	0	0
	D		48	21	7	17	3	U	U


Continued on next page...



Continued	fmom	maniono	maao
Continued	jrom	previous	page

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf
0	C	1	Total	С	Ν	Ο	Р	0	0
	U	L	48	21	7	17	3	0	0
0	р	1	Total	С	Ν	Ο	Р	0	0
	D	L	48	21	7	17	3	0	0


• Molecule 3 is N4-cyclohexyl-5-(4-fluorophenyl)-7H-pyrrolo[2,3-d]pyrimidine-2,4-diamine (three-letter code: UEN) (formula: C₁₈H₂₀FN₅).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total C F N 24 18 1 5	0	0
3	В	1	Total C F N 24 18 1 5	0	0
3	С	1	Total C F N 24 18 1 5	0	0
3	D	1	Total C F N 24 18 1 5	0	0

• Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).





Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	А	1	Total C 4 2	O 2	0	0


• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	230	Total O 230 230	0	0
5	В	201	Total O 201 201	0	0
5	С	181	Total O 181 181	0	0
5	D	155	Total O 155 155	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: PTERIDINE REDUCTASE 1





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	74.06Å 89.68 Å 84.35 Å	Deperitor
a, b, c, α , β , γ	90.00° 115.44° 90.00°	Depositor
Resolution (Å)	37.50 - 1.96	Depositor
Resolution (A)	37.80 - 2.00	EDS
% Data completeness	91.1 (37.50-1.96)	Depositor
(in resolution range)	99.3 (37.80 - 2.00)	EDS
R _{merge}	0.08	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$3.36 (at 2.00 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
D D	0.172 , 0.223	Depositor
R, R_{free}	0.184 , 0.233	DCC
R_{free} test set	3374 reflections $(5.05%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	15.7	Xtriage
Anisotropy	0.032	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.36 , 46.9	EDS
L-test for twinning ²	$< L >=0.48, < L^2>=0.32$	Xtriage
Estimated twinning fraction	0.022 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8567	wwPDB-VP
Average B, all atoms $(Å^2)$	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 21.43 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.0319e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, UEN, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Chain		lengths	Bond angles		
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.72	0/1949	0.80	0/2643	
1	В	0.73	0/1898	0.80	1/2574~(0.0%)	
1	С	0.70	0/1903	0.81	2/2581~(0.1%)	
1	D	0.73	0/1899	0.83	4/2576~(0.2%)	
All	All	0.72	0/7649	0.81	7/10374~(0.1%)	

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$\mathbf{Ideal}(^{o})$
1	D	230	ARG	NE-CZ-NH1	7.80	124.20	120.30
1	D	230	ARG	NE-CZ-NH2	-7.64	116.48	120.30
1	С	159	LEU	CA-CB-CG	-7.43	98.20	115.30
1	D	165	ASP	CB-CG-OD2	-7.25	111.78	118.30
1	С	198	ARG	NE-CZ-NH1	6.76	123.68	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1907	0	1943	18	0
1	В	1868	0	1894	12	0

Continued on next page...



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	С	1867	0	1902	25	0
1	D	1866	0	1891	17	0
2	А	48	0	25	1	0
2	В	48	0	25	0	0
2	С	48	0	25	1	0
2	D	48	0	25	1	0
3	А	24	0	20	1	0
3	В	24	0	20	2	0
3	С	24	0	20	0	0
3	D	24	0	20	0	0
4	А	4	0	3	1	0
5	А	230	0	0	9	0
5	В	201	0	0	3	0
5	С	181	0	0	2	0
5	D	155	0	0	1	0
All	All	8567	0	7813	65	0

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 65 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:167:PRO:HG2	1:C:190[B]:LEU:HD12	1.68	0.76
1:D:164:VAL:HG22	1:D:179:HIS:CD2	2.22	0.74
1:B:103:VAL:HA	1:D:140[B]:GLN:HE22	1.53	0.72
5:A:2132:HOH:O	1:C:140:GLN:NE2	2.22	0.71
1:A:206[B]:VAL:HG11	1:A:221:TRP:CE3	2.29	0.68

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	252/288~(88%)	241~(96%)	11 (4%)	0	100 100
1	В	245/288~(85%)	237~(97%)	7(3%)	1 (0%)	34 22
1	С	246/288~(85%)	236~(96%)	10 (4%)	0	100 100
1	D	245/288~(85%)	239~(98%)	5(2%)	1 (0%)	34 22
All	All	988/1152~(86%)	953~(96%)	33~(3%)	2(0%)	47 38

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	208	LEU
1	D	208	LEU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	\mathbf{ntiles}
1	А	206/231~(89%)	203~(98%)	3~(2%)	65	60
1	В	199/231~(86%)	196~(98%)	3~(2%)	65	60
1	С	200/231~(87%)	199~(100%)	1 (0%)	88	88
1	D	199/231~(86%)	197~(99%)	2(1%)	76	74
All	All	804/924~(87%)	795~(99%)	9 (1%)	73	71

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	43	SER
1	D	216	GLU
1	С	211	VAL
1	А	166	GLN
1	В	152	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:



Mol	Chain	Res	Type
1	С	140	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain	Res	Link	В	ond leng	gths	Bond angles		
	Type	Chain	nes	LINK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAP	D	1269	-	45,52,52	1.05	2 (4%)	56,80,80	1.24	7 (12%)
2	NAP	В	1269	-	45,52,52	1.10	4 (8%)	56,80,80	1.30	6(10%)
4	ACT	А	1271	-	$1,\!3,\!3$	1.98	0	0,3,3	0.00	-
3	UEN	А	1270	-	25,27,27	2.50	11 (44%)	29,38,38	2.75	12 (41%)
3	UEN	D	1270	-	25,27,27	2.23	10 (40%)	29,38,38	2.85	11 (37%)
2	NAP	А	1269	-	45,52,52	0.89	1 (2%)	56,80,80	1.37	7 (12%)
3	UEN	С	1270	-	25,27,27	2.38	10 (40%)	29,38,38	2.76	11 (37%)
3	UEN	В	1270	-	25,27,27	2.51	10 (40%)	29,38,38	2.61	11 (37%)
2	NAP	С	1269	-	45,52,52	0.99	3 (6%)	56,80,80	1.19	5 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the



Mol	Type	Chain	\mathbf{Res}	Link	Chirals	Torsions	Rings
2	NAP	D	1269	-	-	0/31/67/67	0/5/5/5
2	NAP	В	1269	-	-	1/31/67/67	0/5/5/5
3	UEN	А	1270	-	-	0/8/16/16	0/4/4/4
3	UEN	D	1270	-	-	0/8/16/16	0/4/4/4
2	NAP	А	1269	-	-	0/31/67/67	0/5/5/5
3	UEN	С	1270	-	-	0/8/16/16	0/4/4/4
3	UEN	В	1270	-	-	1/8/16/16	0/4/4/4
2	NAP	С	1269	-	-	1/31/67/67	0/5/5/5

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

The worst 5 of 51 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
3	А	1270	UEN	CAT-C5	-6.82	1.30	1.41
3	В	1270	UEN	CAT-C5	-5.86	1.31	1.41
3	С	1270	UEN	CAT-C5	-5.79	1.31	1.41
3	В	1270	UEN	CAF-CAS	5.26	1.50	1.39
3	А	1270	UEN	CAE-CAS	4.95	1.49	1.39

The worst 5 of 70 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
3	А	1270	UEN	CAT-C5-C4	8.41	114.25	107.54
3	D	1270	UEN	CAT-C5-C4	7.80	113.76	107.54
3	С	1270	UEN	CAT-C5-C4	6.61	112.81	107.54
3	С	1270	UEN	C5-C6-NAO	6.40	126.67	120.66
3	D	1270	UEN	C5-C6-NAO	6.05	126.34	120.66

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	В	1269	NAP	C3B-C2B-O2B-P2B
2	С	1269	NAP	C5B-O5B-PA-O1A
3	В	1270	UEN	CAE-CAS-CAT-C5


There are no ring outliers.

6 monomers are involved in 7 short contacts:




Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	1269	NAP	1	0
4	А	1271	ACT	1	0
3	А	1270	UEN	1	0
2	А	1269	NAP	1	0
3	В	1270	UEN	2	0
2	С	1269	NAP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\# RSRZ {>}2$	$OWAB(A^2)$	Q<0.9
1	А	251/288~(87%)	0.00	11 (4%) 34 44	9, 15, 34, 56	0
1	В	250/288~(86%)	-0.08	7 (2%) 53 62	9, 15, 31, 54	0
1	С	249/288~(86%)	0.01	10 (4%) 38 48	9, 15, 36, 62	0
1	D	249/288~(86%)	0.03	9 (3%) 42 52	8, 16, 35, 55	0
All	All	999/1152~(86%)	-0.01	37 (3%) 41 51	8, 15, 35, 62	0

The worst 5 of 37 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	113	GLY	8.0
1	В	113	GLY	4.8
1	С	212	ALA	4.7
1	D	211	VAL	4.4
1	А	151	SER	4.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
3	UEN	D	1270	24/24	0.91	0.12	14, 19, 33, 44	0
3	UEN	С	1270	24/24	0.91	0.12	13, 18, 34, 44	0
3	UEN	В	1270	24/24	0.93	0.10	11, 16, 26, 33	0
3	UEN	А	1270	24/24	0.94	0.09	14, 19, 26, 33	0
2	NAP	В	1269	48/48	0.97	0.08	$9,\!12,\!16,\!19$	0
2	NAP	А	1269	48/48	0.98	0.07	$8,\!12,\!15,\!15$	0
2	NAP	D	1269	48/48	0.98	0.07	$10,\!13,\!16,\!17$	0
4	ACT	А	1271	4/4	0.98	0.08	$15,\!15,\!16,\!16$	0
2	NAP	С	1269	48/48	0.98	0.07	$10,\!12,\!13,\!14$	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.































6.5 Other polymers (i)

There are no such residues in this entry.

