

wwPDB X-ray Structure Validation Summary Report (i)

May 24, 2020 – 05:36 am BST

PDB ID : 4CMG

Title : Crystal structure of pteridine reductase 1 (PTR1) from Trypanosoma brucei

in ternary complex with cofactor and inhibitor

Authors : Barrack, K.L.; Hunter, W.N.

Deposited on : 2014-01-16

Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.11

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\ DIFFRACTION$

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$egin{aligned} ext{Similar resolution} \ (\# ext{Entries}, ext{resolution range}(\AA)) \end{aligned}$		
R_{free}	130704	8085 (2.00-2.00)		
Clashscore	141614	9178 (2.00-2.00)		
Ramachandran outliers	138981	9054 (2.00-2.00)		
Sidechain outliers	138945	9053 (2.00-2.00)		
RSRZ outliers	127900	7900 (2.00-2.00)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	288	80%	% 13%
1	В	288	76% 10%	• 13%
1	С	288	79% 7%	13%
1	D	288	76% 11%	13%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 8279 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called PTERIDINE REDUCTASE 1.

Mol	Chain	Residues	${f Atoms}$			ZeroOcc	AltConf	Trace		
1	Λ	250	Total	С	N	О	S	0	0	0
1	A	250	1860	1168	328	353	11	0	0	
1	В	251	Total	С	N	О	S	0	0	0
1		201	1869	1174	330	354	11	0	0	0
1	С	250	Total	С	N	О	S	0	1	0
1		250	1869	1173	329	356	11	0		
1	D	250	Total	С	N	О	S	0	0	0
1	ש	250	1860	1168	328	353	11		U	

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP O76290
A	-18	GLY	-	expression tag	UNP O76290
A	-17	SER	-	expression tag	UNP O76290
A	-16	SER	_	expression tag	UNP O76290
A	-15	HIS	-	expression tag	UNP O76290
A	-14	HIS	_	expression tag	UNP O76290
A	-13	HIS	_	expression tag	UNP O76290
A	-12	HIS	_	expression tag	UNP O76290
A	-11	HIS	_	expression tag	UNP O76290
A	-10	HIS	-	expression tag	UNP O76290
A	-9	SER	_	expression tag	UNP O76290
A	-8	SER	_	expression tag	UNP O76290
A	-7	GLY	_	expression tag	UNP O76290
A	-6	LEU	_	expression tag	UNP O76290
A	-5	VAL	_	expression tag	UNP O76290
A	-4	PRO	-	expression tag	UNP O76290
A	-3	ARG	=	expression tag	UNP O76290
A	-2	GLY	-	expression tag	UNP O76290
A	-1	SER		expression tag	UNP O76290
A	0	HIS	-	expression tag	UNP O76290
В	-19	MET	_	expression tag	UNP O76290



Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
В	-18	GLY	-	expression tag	UNP O76290
В	-17	SER	-	expression tag	UNP O76290
В	-16	SER	-	expression tag	UNP O76290
В	-15	HIS	-	expression tag	UNP O76290
В	-14	HIS	-	expression tag	UNP O76290
В	-13	HIS	-	expression tag	UNP O76290
В	-12	HIS	-	expression tag	UNP O76290
В	-11	HIS	-	expression tag	UNP O76290
В	-10	HIS	-	expression tag	UNP O76290
В	-9	SER	-	expression tag	UNP O76290
В	-8	SER	-	expression tag	UNP O76290
В	-7	GLY	-	expression tag	UNP O76290
В	-6	LEU	-	expression tag	UNP O76290
В	-5	VAL	-	expression tag	UNP O76290
В	-4	PRO	_	expression tag	UNP O76290
В	-3	ARG	-	expression tag	UNP O76290
В	-2	GLY	_	expression tag	UNP O76290
В	-1	SER	-	expression tag	UNP O76290
В	0	HIS	-	expression tag	UNP O76290
С	-19	MET	-	expression tag	UNP O76290
С	-18	GLY	-	expression tag	UNP O76290
С	-17	SER	_	expression tag	UNP O76290
С	-16	SER	-	expression tag	UNP O76290
С	-15	HIS	_	expression tag	UNP O76290
С	-14	HIS	-	expression tag	UNP O76290
С	-13	HIS	-	expression tag	UNP O76290
С	-12	HIS	-	expression tag	UNP O76290
С	-11	HIS	-	expression tag	UNP O76290
С	-10	HIS	-	expression tag	UNP O76290
С	-9	SER	-	expression tag	UNP O76290
С	-8	SER	-	expression tag	UNP O76290
С	-7	GLY	-	expression tag	UNP O76290
С	-6	LEU	-	expression tag	UNP O76290
С	-5	VAL	-	expression tag	UNP O76290
С	-4	PRO	-	expression tag	UNP O76290
С	-3	ARG	-	expression tag	UNP O76290
С	-2	GLY	-	expression tag	UNP O76290
С	-1	SER	-	expression tag	UNP O76290
С	0	HIS	-	expression tag	UNP O76290
D	-19	MET	-	expression tag	UNP O76290
D	-18	GLY	-	expression tag	UNP O76290
D	-17	SER	-	expression tag	UNP O76290



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-16	SER	_	expression tag	UNP O76290
D	-15	HIS	-	expression tag	UNP O76290
D	-14	HIS	-	expression tag	UNP O76290
D	-13	HIS	-	expression tag	UNP O76290
D	-12	HIS	_	expression tag	UNP O76290
D	-11	HIS	_	expression tag	UNP O76290
D	-10	HIS	-	expression tag	UNP O76290
D	-9	SER	_	expression tag	UNP O76290
D	-8	SER	-	expression tag	UNP O76290
D	-7	GLY	-	expression tag	UNP O76290
D	-6	LEU	_	expression tag	UNP O76290
D	-5	VAL	-	expression tag	UNP O76290
D	-4	PRO	_	expression tag	UNP O76290
D	-3	ARG	-	expression tag	UNP O76290
D	-2	GLY	-	expression tag	UNP O76290
D	-1	SER	=	expression tag	UNP O76290
D	0	HIS	-	expression tag	UNP O76290

• Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).




Mol	Chain	Residues		At	oms			ZeroOcc	AltConf
9	Λ	1	Total	С	N	О	Р	0	0
	2 A	1	48	21	7	17	3		0
9	D	1	Total	С	N	О	Р	0	0
	Б	1	48	21	7	17	3	U	U



Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
9	2 C	1	Total	С	N	О	Р	0	0
2		1	48	21	7	17	3	U	0
9	D	1	Total	С	N	О	Р	0	0
	D	1	48	21	7	17	3		U


• Molecule 3 is 6-(4-fluorophenyl)-5-(4-methoxyphenyl)-7H-pyrrolo[2,3-d]pyrimidine-2,4-diam ine (three-letter code: UHX) (formula: $C_{19}H_{16}FN_5O$).



Mol	Chain	Residues		Ato	ms			ZeroOcc	AltConf
3	Δ	1	Total	С	F	N	О	0	0
	11	1	26	19	1	5	1		U
3	R	1	Total	С	F	Ν	Ο	0	0
'	Б	1	26	19	1	5	1		U
3	С	1	Total	С	F	N	О	0	0
'		1	26	19	1	5	1	0	0
3	D	1	Total	С	F	N	О	0	0
)	ש	1	26	19	1	5	1	U	U

• Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	В	1	Total C O 4 2 2	0	0
4	С	1	Total C O 4 2 2	0	0


• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	150	Total O 150 150	0	0
5	В	133	Total O 133 133	0	0
5	С	114	Total O 114 114	0	0
5	D	120	Total O 120 120	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	73.92Å 89.66Å 84.08Å	Donositor
a, b, c, α , β , γ	90.00° 115.97° 90.00°	Depositor
Resolution (Å)	44.83 - 2.00	Depositor
Resolution (A)	44.83 - 2.00	EDS
% Data completeness	95.8 (44.83-2.00)	Depositor
(in resolution range)	95.8 (44.83-2.00)	EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.78 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
D D	0.161 , 0.213	Depositor
R, R_{free}	0.172 , 0.221	DCC
R_{free} test set	3272 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	15.4	Xtriage
Anisotropy	0.048	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.38 , 45.5	EDS
L-test for twinning ²	$< L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	0.016 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8279	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 18.66% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, UHX, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
10101	Chain	RMSZ	# Z >5	RMSZ	# Z > 5	
1	A	0.76	0/1887	0.86	3/2560~(0.1%)	
1	В	0.75	0/1896	0.88	6/2571~(0.2%)	
1	С	0.73	0/1896	0.89	3/2572~(0.1%)	
1	D	0.75	0/1887	0.88	3/2560~(0.1%)	
All	All	0.75	0/7566	0.88	$15/10263 \; (0.1\%)$	

There are no bond length outliers.

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^o)$
1	С	198	ARG	NE-CZ-NH2	-9.65	115.47	120.30
1	D	222	ARG	NE-CZ-NH1	9.08	124.84	120.30
1	В	230	ARG	NE-CZ-NH1	8.03	124.31	120.30
1	С	198	ARG	NE-CZ-NH1	8.01	124.30	120.30
1	D	222	ARG	NE-CZ-NH2	-7.67	116.46	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1860	0	1882	10	0
1	В	1869	0	1895	20	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	С	1869	0	1887	11	0
1	D	1860	0	1882	23	0
2	A	48	0	25	0	0
2	В	48	0	25	0	0
2	С	48	0	25	0	0
2	D	48	0	25	0	0
3	A	26	0	16	0	0
3	В	26	0	16	0	0
3	С	26	0	16	0	0
3	D	26	0	16	1	0
4	В	4	0	3	0	0
4	С	4	0	3	0	0
5	A	150	0	0	1	0
5	В	133	0	0	8	0
5	С	114	0	0	4	0
5	D	120	0	0	5	0
All	All	8279	0	7716	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 63 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{aligned}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:D:78:ASN:HB3	5:D:2041:HOH:O	1.43	1.19
1:C:215[A]:GLU:HG3	5:C:2084:HOH:O	1.71	0.91
1:D:78:ASN:OD1	1:D:141:ARG:NH1	2.08	0.87
1:B:103:VAL:HG22	5:B:2068:HOH:O	1.83	0.78
1:B:175:ASN:HB2	5:B:2099:HOH:O	1.85	0.77

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the	number of residues	for which t	the backbone	conformation	was
analysed, and the total number of	residues.				

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	$_{ m ntiles}$
1	A	244/288 (85%)	233 (96%)	11 (4%)	0	100	100
1	В	$245/288 \; (85\%)$	237 (97%)	8 (3%)	0	100	100
1	С	$245/288 \ (85\%)$	237 (97%)	8 (3%)	0	100	100
1	D	$244/288 \; (85\%)$	235 (96%)	9 (4%)	0	100	100
All	All	978/1152 (85%)	942 (96%)	36 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	198/231~(86%)	195 (98%)	3 (2%)	65 69		
1	В	199/231~(86%)	195 (98%)	4 (2%)	55 58		
1	С	199/231~(86%)	192 (96%)	7 (4%)	36 35		
1	D	198/231~(86%)	197 (100%)	1 (0%)	88 92		
All	All	794/924~(86%)	779 (98%)	15 (2%)	59 61		

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	${f Res}$	Type
1	В	229	ARG
1	С	166	GLN
1	С	242	ILE
1	В	213	MET
1	С	216	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:



Mol	Chain	Res	Type
1	A	250	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trens	Chain	Res	Link	Во	ond leng	ths	Bond angles		
MIOI	Type	Chain	nes	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
3	UHX	D	1270	-	29,29,29	1.59	7 (24%)	32,42,42	1.82	6 (18%)
2	NAP	С	1269	-	45,52,52	0.90	1 (2%)	56,80,80	1.22	5 (8%)
2	NAP	A	1269	-	45,52,52	0.92	3 (6%)	56,80,80	1.17	5 (8%)
4	ACT	С	1271	-	1,3,3	1.32	0	0,3,3	0.00	-
4	ACT	В	1271	-	1,3,3	0.44	0	0,3,3	0.00	-
2	NAP	D	1269	-	45,52,52	1.01	3 (6%)	56,80,80	1.19	4 (7%)
3	UHX	A	1270	-	29,29,29	1.24	3 (10%)	32,42,42	1.35	5 (15%)
3	UHX	С	1270	-	29,29,29	1.29	4 (13%)	32,42,42	1.43	4 (12%)
2	NAP	В	1269	-	45,52,52	0.94	1 (2%)	56,80,80	1.43	9 (16%)
3	UHX	В	1270	-	29,29,29	1.19	3 (10%)	32,42,42	1.84	10 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral



centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	UHX	D	1270	-	-	2/10/10/10	0/4/4/4
2	NAP	С	1269	-	-	0/31/67/67	0/5/5/5
2	NAP	A	1269	-	-	0/31/67/67	0/5/5/5
2	NAP	D	1269	_	-	0/31/67/67	0/5/5/5
3	UHX	A	1270	-	-	2/10/10/10	0/4/4/4
3	UHX	С	1270	-	-	2/10/10/10	0/4/4/4
2	NAP	В	1269	-	-	0/31/67/67	0/5/5/5
3	UHX	В	1270	_	-	2/10/10/10	0/4/4/4

The worst 5 of 25 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(ext{\AA})$
3	D	1270	UHX	CAX-CAW	-4.53	1.36	1.41
3	D	1270	UHX	C2-NAB	3.64	1.41	1.33
2	D	1269	NAP	P2B-O2B	3.37	1.65	1.59
3	В	1270	UHX	C2-NAB	3.25	1.40	1.33
3	С	1270	UHX	C2-NAB	3.16	1.40	1.33

The worst 5 of 48 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\mathbf{Ideal}(^{o})$
3	В	1270	UHX	N3-C2-N1	-5.60	119.75	127.22
3	С	1270	UHX	N3-C2-N1	-4.93	120.65	127.22
3	D	1270	UHX	C2-N3-C4	4.71	120.74	115.36
3	D	1270	UHX	N3-C2-N1	-4.57	121.13	127.22
3	A	1270	UHX	N3-C2-N1	-4.04	121.83	127.22

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	В	1270	UHX	CAG-CAT-OAP-CAA
3	В	1270	UHX	CAH-CAT-OAP-CAA
3	D	1270	UHX	CAG-CAT-OAP-CAA
3	D	1270	UHX	CAH-CAT-OAP-CAA
3	С	1270	UHX	CAG-CAT-OAP-CAA

There are no ring outliers.




1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	1270	UHX	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	A	250/288~(86%)	-0.53	3 (1%) 79 78	8, 14, 36, 67	0
1	В	251/288 (87%)	-0.49	2 (0%) 86 85	7, 15, 34, 67	0
1	С	$250/288 \; (86\%)$	-0.43	6 (2%) 59 57	8, 16, 35, 57	0
1	D	$250/288 \; (86\%)$	-0.54	1 (0%) 92 92	9, 15, 32, 48	0
All	All	1001/1152~(86%)	-0.50	12 (1%) 79 78	7, 15, 34, 67	0

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	151	SER	6.0
1	С	151	SER	3.5
1	С	212	ALA	3.2
1	С	2	GLU	3.1
1	D	151	SER	2.8

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
3	UHX	D	1270	26/26	0.95	0.09	12,16,21,21	0
3	UHX	A	1270	26/26	0.95	0.10	13,20,28,35	0
3	UHX	С	1270	26/26	0.95	0.08	14,20,24,25	0
3	UHX	В	1270	26/26	0.96	0.07	14,16,23,30	0
2	NAP	D	1269	48/48	0.98	0.07	9,12,15,15	0
2	NAP	С	1269	48/48	0.98	0.07	9,13,14,15	0
2	NAP	A	1269	48/48	0.98	0.07	9,11,13,14	0
2	NAP	В	1269	48/48	0.98	0.07	9,12,15,17	0
4	ACT	С	1271	4/4	0.98	0.09	17,17,18,18	0
4	ACT	В	1271	4/4	0.99	0.14	18,20,21,22	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





















Electron density around NAP D 1269:

 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NAP C 1269:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around NAP A 1269:

 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NAP B 1269:

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

