



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 19, 2024 – 06:06 AM EDT

PDB ID : 4FFW  
Title : Crystal Structure of Dipeptidyl Peptidase IV (DPP4, DPP-IV, CD26) in Complex with Fab + sitagliptin  
Authors : Wang, Z.; Sudom, A.; Walker, N.P.; Min, X.  
Deposited on : 2012-06-01  
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 1.20.1  
EDS : 2.37.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1957 (2.90-2.90)
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	730	
1	B	730	
2	C	210	
2	L	210	
3	D	217	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	H	217	 <p>A horizontal bar chart representing the quality of chain H. The bar is divided into three segments: a red segment on the left labeled '23%', a large green segment in the middle labeled '87%', and a small yellow segment on the right labeled '8%'. The yellow segment is followed by two small grey dots.</p>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 18266 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dipeptidyl peptidase 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	727	Total	C	N	O	S	0	0	0
			5921	3792	980	1123	26			
1	B	729	Total	C	N	O	S	0	0	0
			5940	3804	985	1125	26			


- Molecule 2 is a protein called Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	207	Total	C	N	O	S	0	0	0
			1588	987	272	322	7			
2	L	202	Total	C	N	O	S	0	0	0
			1535	956	260	312	7			

- Molecule 3 is a protein called Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	D	209	Total	C	N	O	S	0	0	0
			1597	1019	259	314	5			
3	H	208	Total	C	N	O	S	0	0	0
			1590	1014	258	313	5			

- Molecule 4 is (2R)-4-OXO-4-[3-(TRIFLUOROMETHYL)-5,6-DIHYDRO[1,2,4]TRIAZOLO[4,3-A]PYRAZIN-7(8H)-YL]-1-(2,4,5-TRIFLUOROPHENYL)BUTAN-2-A MINE (three-letter code: 715) (formula: C<sub>16</sub>H<sub>15</sub>F<sub>6</sub>N<sub>5</sub>O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	F	N	O	0	0
			28	16	6	5	1		
4	B	1	Total	C	F	N	O	0	0
			28	16	6	5	1		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	27	Total	O	0	0
			27	27		
5	B	11	Total	O	0	0
			11	11		
5	C	1	Total	O	0	0
			1	1		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Dipeptidyl peptidase 4

Chain A: 





- Molecule 1: Dipeptidyl peptidase 4

Chain B: 




- Molecule 2: Fab light chain


Chain C: 




- Molecule 2: Fab light chain



• Molecule 3: Fab heavy chain



• Molecule 3: Fab heavy chain



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	97.77Å 200.86Å 97.93Å 90.00° 93.64° 90.00°	Depositor
Resolution (Å)	29.99 – 2.90 29.99 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.1 (29.99-2.90) 98.7 (29.99-2.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.58 (at 2.90Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.287 , 0.311 0.286 , 0.311	Depositor DCC
$R_{free}$ test set	4123 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	68.8	Xtrriage
Anisotropy	0.172	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 32.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.450 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	18266	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	94.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.27% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
715

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/6090	0.44	0/8280
1	B	0.26	0/6109	0.44	0/8305
2	C	0.27	0/1628	0.41	0/2213
2	L	0.26	0/1574	0.42	0/2143
3	D	0.27	0/1637	0.43	0/2236
3	H	0.27	0/1630	0.43	0/2226
All	All	0.26	0/18668	0.43	0/25403

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5921	0	5658	10	0
1	B	5940	0	5682	12	0
2	C	1588	0	1515	4	0
2	L	1535	0	1466	2	0
3	D	1597	0	1575	11	0
3	H	1590	0	1566	9	0
4	A	28	0	15	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	28	0	15	0	0
5	A	27	0	0	0	0
5	B	11	0	0	0	0
5	C	1	0	0	0	0
All	All	18266	0	17492	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 45 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:287:ALA:HB1	1:B:288:PRO:HA	1.66	0.76
1:B:67:LEU:HB3	1:B:74:ILE:HD11	1.83	0.59
1:B:287:ALA:HB1	1:B:288:PRO:CA	2.31	0.58
3:H:12:VAL:HG11	3:H:18:VAL:HG13	1.86	0.57
1:A:547:VAL:HG21	1:A:636:VAL:HG11	1.88	0.56

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	725/730 (99%)	680 (94%)	43 (6%)	2 (0%)	41 71
1	B	727/730 (100%)	683 (94%)	40 (6%)	4 (1%)	25 58
2	C	203/210 (97%)	177 (87%)	22 (11%)	4 (2%)	7 27
2	L	198/210 (94%)	173 (87%)	21 (11%)	4 (2%)	7 27
3	D	205/217 (94%)	179 (87%)	20 (10%)	6 (3%)	4 18
3	H	204/217 (94%)	186 (91%)	15 (7%)	3 (2%)	10 34

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	2262/2314 (98%)	2078 (92%)	161 (7%)	23 (1%)	15 45

5 of 23 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	300	ASP
1	B	300	ASP
3	D	152	GLU
3	D	162	SER
3	H	152	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	648/651 (100%)	625 (96%)	23 (4%)	36 70
1	B	650/651 (100%)	628 (97%)	22 (3%)	37 71
2	C	183/185 (99%)	177 (97%)	6 (3%)	38 72
2	L	177/185 (96%)	174 (98%)	3 (2%)	60 86
3	D	184/189 (97%)	179 (97%)	5 (3%)	44 77
3	H	183/189 (97%)	176 (96%)	7 (4%)	33 67
All	All	2025/2050 (99%)	1959 (97%)	66 (3%)	38 72

5 of 66 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	H	20	ILE
3	H	62	GLN
2	L	89	GLN
1	B	109	ARG
1	B	39	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	124	HIS
1	B	440	HIS
1	B	341	GLN
1	B	711	ASN
1	A	319	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	715	A	801	-	27,30,30	0.78	0	34,45,45	1.27	4 (11%)
4	715	B	801	-	27,30,30	0.80	0	34,45,45	1.24	4 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	715	A	801	-	-	2/18/27/27	0/2/3/3
4	715	B	801	-	-	0/18/27/27	0/2/3/3

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	801	715	C4-C3-C2	3.11	119.96	116.54
4	A	801	715	C4-C3-C2	2.88	119.70	116.54
4	B	801	715	C26-N27-C39	2.62	130.69	125.38
4	B	801	715	C1-C2-C3	-2.58	119.78	123.76
4	A	801	715	C26-N27-C39	2.52	130.49	125.38

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	801	715	C15-C16-N19-C29
4	A	801	715	O22-C16-N19-C29

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	727/730 (99%)	0.31	8 (1%) 80 80	36, 73, 104, 123	0
1	B	729/730 (99%)	0.31	14 (1%) 66 65	34, 74, 104, 123	0
2	C	207/210 (98%)	2.07	89 (42%) 0 0	89, 145, 190, 196	0
2	L	202/210 (96%)	2.05	89 (44%) 0 0	86, 138, 192, 200	0
3	D	209/217 (96%)	1.52	67 (32%) 0 0	83, 126, 163, 168	0
3	H	208/217 (95%)	1.29	50 (24%) 0 0	79, 124, 158, 167	0
All	All	2282/2314 (98%)	0.83	317 (13%) 2 2	34, 89, 169, 200	0

The worst 5 of 317 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	181	LEU	10.5
2	L	110	ALA	10.2
3	D	101	ASP	10.1
2	C	140	PRO	9.7
2	L	196	THR	8.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	715	A	801	28/28	0.89	0.22	69,78,91,92	0
4	715	B	801	28/28	0.91	0.21	68,76,89,89	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







## 6.5 Other polymers [i](#)

There are no such residues in this entry.