



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 18, 2024 – 02:50 PM EST

PDB ID : 4FKB
Title : An Organic solvent tolerant lipase 42
Authors : Zaliha, R.N.; Rahman, R.A.; Khusaini, M.S.
Deposited on : 2012-06-13
Resolution : 1.22 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1232 (1.24-1.20)
Clashscore	141614	1294 (1.24-1.20)
Ramachandran outliers	138981	1251 (1.24-1.20)
Sidechain outliers	138945	1250 (1.24-1.20)
RSRZ outliers	127900	1209 (1.24-1.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	388	 88% 11% .
1	B	388	 86% 11% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	413	-	X	X	-
3	GOL	B	417	-	-	X	-
3	GOL	B	418	-	-	X	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7622 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Thermostable organic solvent tolerant lipase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	387	3154	2010	549	586	9	0	21	0
1	B	387	3286	2099	575	601	11	0	39	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	147	LEU	TRP	engineered mutation	UNP Q5U780
A	202	GLU	GLN	engineered mutation	UNP Q5U780
B	147	LEU	TRP	engineered mutation	UNP Q5U780
B	202	GLU	GLN	engineered mutation	UNP Q5U780

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O 1 1	0	0
3	A	1	Total C O 12 6 6	0	1
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 5 3 2	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 18 9 9	0	1
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	1
			12	6	6		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	4	Total	Na	0	0
			4	4		
4	B	5	Total	Na	0	0
			5	5		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	5	Total	Cl	0	0
			5	5		
5	B	5	Total	Cl	0	0
			5	5		

- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Ca	0	0
			1	1		
6	B	1	Total	Ca	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	524	Total 524	O 524	0	0
7	B	491	Total 491	O 491	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Thermostable organic solvent tolerant lipase



- Molecule 1: Thermostable organic solvent tolerant lipase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	117.82Å 81.00Å 99.53Å 90.00° 96.69° 90.00°	Depositor
Resolution (Å)	50.00 – 1.22 26.31 – 1.22	Depositor EDS
% Data completeness (in resolution range)	93.7 (50.00-1.22) 93.7 (26.31-1.22)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.55 (at 1.22Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.127 , 0.165 0.127 , 0.165	Depositor DCC
R_{free} test set	12959 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	7.3	Xtrriage
Anisotropy	0.136	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 58.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	7622	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CA, CL, GOL, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.23	7/3298 (0.2%)	1.26	22/4483 (0.5%)
1	B	1.27	7/3475 (0.2%)	1.25	30/4711 (0.6%)
All	All	1.25	14/6773 (0.2%)	1.26	52/9194 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	303	ARG	CZ-NH1	10.59	1.46	1.33
1	B	202	GLU	CD-OE2	10.02	1.36	1.25
1	A	100	GLU	CD-OE1	8.52	1.35	1.25
1	A	303	ARG	CZ-NH2	8.36	1.44	1.33
1	A	40	TRP	CD2-CE2	6.93	1.49	1.41

The worst 5 of 52 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	47	ARG	NE-CZ-NH2	-11.31	114.64	120.30
1	B	106	ARG	NE-CZ-NH1	-11.16	114.72	120.30
1	A	230	ARG	NE-CZ-NH1	10.86	125.73	120.30
1	A	330[A]	ARG	NE-CZ-NH1	-10.61	115.00	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	330[B]	ARG	NE-CZ-NH1	-10.61	115.00	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	16[A]	PHE	Peptide
1	B	16	PHE	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3154	0	3068	38	0
1	B	3286	0	3258	60	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	54	0	68	15	0
3	B	90	0	120	21	0
4	A	4	0	0	0	0
4	B	5	0	0	0	0
5	A	5	0	0	0	0
5	B	5	0	0	1	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
7	A	524	0	0	15	2
7	B	491	0	0	15	3
All	All	7622	0	6514	107	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 107 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:16[B]:PHE:CE2	7:A:710:HOH:O	1.83	1.29

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:214[A]:ARG:HH21	1:B:214[A]:ARG:CG	1.35	1.29
1:B:174[B]:VAL:HG23	7:B:968:HOH:O	1.30	1.26
1:A:306[B]:THR:HG22	7:A:683:HOH:O	1.04	1.22
1:B:173[B]:MET:CE	1:B:179[B]:ARG:HD2	1.71	1.20

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:683:HOH:O	7:B:718:HOH:O[4_545]	2.03	0.17
7:B:736:HOH:O	7:B:761:HOH:O[4_545]	2.09	0.11
7:A:684:HOH:O	7:B:682:HOH:O[3_445]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	406/388 (105%)	397 (98%)	9 (2%)	0	100	100
1	B	425/388 (110%)	414 (97%)	11 (3%)	0	100	100
All	All	831/776 (107%)	811 (98%)	20 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	337/318 (106%)	334 (99%)	3 (1%)	78	50
1	B	357/318 (112%)	350 (98%)	7 (2%)	55	17
All	All	694/636 (109%)	684 (99%)	10 (1%)	69	32

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	285[B]	LEU
1	B	307	LEU
1	B	357	ASP
1	B	202	GLU
1	B	214[A]	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	153	HIS
1	B	172	ASN
1	B	257	GLN
1	B	216	GLN
1	B	44	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 48 ligands modelled in this entry, 23 are monoatomic and 1 is modelled with single atom - leaving 24 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	B	408[B]	-	5,5,5	0.51	0	5,5,5	0.88	0
3	GOL	B	412	-	5,5,5	0.64	0	5,5,5	0.93	0
3	GOL	B	407	-	5,5,5	0.90	0	5,5,5	0.91	0
3	GOL	A	409	-	5,5,5	0.98	0	5,5,5	1.10	0
3	GOL	A	403[A]	-	5,5,5	0.70	0	5,5,5	0.84	0
3	GOL	B	415	-	5,5,5	0.80	0	5,5,5	0.94	0
3	GOL	B	413[A]	-	5,5,5	1.18	0	5,5,5	1.22	0
3	GOL	A	403[B]	-	5,5,5	0.44	0	5,5,5	1.08	0
3	GOL	B	408[C]	-	5,5,5	0.64	0	5,5,5	0.65	0
3	GOL	A	410	-	5,5,5	0.79	0	5,5,5	1.26	0
3	GOL	B	413[B]	-	5,5,5	1.12	1 (20%)	5,5,5	1.39	1 (20%)
3	GOL	A	413	-	5,5,5	1.03	0	5,5,5	2.67	2 (40%)
3	GOL	B	409	-	5,5,5	0.54	0	5,5,5	1.78	3 (60%)
3	GOL	B	411	-	5,5,5	1.22	1 (20%)	5,5,5	0.84	0
3	GOL	B	418	-	5,5,5	0.64	0	5,5,5	1.58	1 (20%)
3	GOL	B	410	-	5,5,5	0.96	0	5,5,5	0.63	0
3	GOL	B	416	-	5,5,5	0.83	0	5,5,5	1.20	0
3	GOL	A	414	-	5,5,5	0.66	0	5,5,5	0.82	0
3	GOL	B	414	-	5,5,5	0.33	0	5,5,5	0.63	0
3	GOL	A	411	-	3,4,5	0.50	0	1,4,5	0.72	0
3	GOL	A	408	-	5,5,5	0.22	0	5,5,5	0.76	0
3	GOL	A	412	-	5,5,5	0.77	0	5,5,5	1.12	1 (20%)
3	GOL	B	417	-	5,5,5	0.76	0	5,5,5	0.66	0
3	GOL	B	408[A]	-	5,5,5	0.69	0	5,5,5	0.87	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	408[B]	-	-	0/4/4/4	-
3	GOL	B	412	-	-	1/4/4/4	-
3	GOL	B	407	-	-	0/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	409	-	-	0/4/4/4	-
3	GOL	A	403[A]	-	-	4/4/4/4	-
3	GOL	B	415	-	-	2/4/4/4	-
3	GOL	B	413[A]	-	-	0/4/4/4	-
3	GOL	A	403[B]	-	-	2/4/4/4	-
3	GOL	B	408[C]	-	-	1/4/4/4	-
3	GOL	A	410	-	-	0/4/4/4	-
3	GOL	B	413[B]	-	-	0/4/4/4	-
3	GOL	A	413	-	-	4/4/4/4	-
3	GOL	B	409	-	-	2/4/4/4	-
3	GOL	B	411	-	-	0/4/4/4	-
3	GOL	B	418	-	-	4/4/4/4	-
3	GOL	B	410	-	-	0/4/4/4	-
3	GOL	B	416	-	-	2/4/4/4	-
3	GOL	A	414	-	-	4/4/4/4	-
3	GOL	B	414	-	-	1/4/4/4	-
3	GOL	A	411	-	-	2/2/2/4	-
3	GOL	A	408	-	-	4/4/4/4	-
3	GOL	A	412	-	-	2/4/4/4	-
3	GOL	B	417	-	-	0/4/4/4	-
3	GOL	B	408[A]	-	-	0/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	411	GOL	O3-C3	2.44	1.52	1.42
3	B	413[B]	GOL	O2-C2	-2.07	1.37	1.43

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	413	GOL	O2-C2-C3	3.88	126.21	109.12
3	A	413	GOL	O3-C3-C2	3.81	128.45	110.20
3	B	413[B]	GOL	O2-C2-C3	-2.34	98.83	109.12
3	A	412	GOL	O2-C2-C1	-2.28	99.09	109.12
3	B	409	GOL	C3-C2-C1	-2.16	103.31	111.70

There are no chirality outliers.

5 of 35 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	403[B]	GOL	O1-C1-C2-C3
3	A	411	GOL	O2-C2-C3-O3
3	A	412	GOL	C1-C2-C3-O3
3	A	412	GOL	O2-C2-C3-O3
3	A	414	GOL	O1-C1-C2-C3

There are no ring outliers.

10 monomers are involved in 36 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	407	GOL	2	0
3	B	415	GOL	1	0
3	B	413[A]	GOL	1	0
3	A	413	GOL	11	0
3	B	418	GOL	6	0
3	B	416	GOL	2	0
3	A	414	GOL	1	0
3	A	411	GOL	2	0
3	A	412	GOL	1	0
3	B	417	GOL	9	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	387/388 (99%)	-0.80	2 (0%) 91 90	5, 11, 23, 33	0
1	B	387/388 (99%)	-0.75	5 (1%) 77 75	4, 9, 22, 49	0
All	All	774/776 (99%)	-0.77	7 (0%) 84 82	4, 10, 23, 49	0

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	3	LEU	6.2
1	B	2	SER	5.2
1	B	174[A]	VAL	4.3
1	A	277	LEU	2.3
1	A	278	THR	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	NA	A	406	1/1	0.83	0.21	64,64,64,64	0
3	GOL	A	414	6/6	0.88	0.20	43,51,82,94	0
3	GOL	A	410	6/6	0.88	0.14	31,35,38,41	0
3	GOL	B	409	6/6	0.89	0.14	34,39,49,60	0
3	GOL	B	418	6/6	0.90	0.26	24,35,44,46	0
3	GOL	B	414	6/6	0.90	0.17	44,50,60,71	0
3	GOL	B	415	6/6	0.91	0.14	27,33,35,53	0
3	GOL	B	416	6/6	0.91	0.15	20,35,50,56	0
3	GOL	B	417	6/6	0.92	0.19	20,31,43,51	0
3	GOL	A	403[A]	6/6	0.92	0.19	21,34,37,39	6
3	GOL	A	403[B]	6/6	0.92	0.19	21,25,27,34	6
3	GOL	B	407	6/6	0.94	0.17	16,22,25,28	0
4	NA	B	405	1/1	0.94	0.17	42,42,42,42	0
4	NA	A	407	1/1	0.95	0.33	37,37,37,37	0
3	GOL	A	408	6/6	0.96	0.15	41,43,57,72	0
3	GOL	A	411	5/6	0.96	0.14	30,33,43,44	0
3	GOL	A	413	6/6	0.96	0.18	16,22,32,44	0
3	GOL	B	410	6/6	0.96	0.09	14,17,24,32	0
3	GOL	A	412	6/6	0.97	0.21	19,42,51,61	0
3	GOL	B	412	6/6	0.97	0.08	15,21,25,25	0
3	GOL	B	413[A]	6/6	0.97	0.17	13,17,21,23	6
4	NA	B	404	1/1	0.97	0.11	31,31,31,31	0
3	GOL	B	413[B]	6/6	0.97	0.17	8,17,24,30	6
4	NA	A	405	1/1	0.98	0.11	25,25,25,25	0
4	NA	B	403	1/1	0.98	0.18	40,40,40,40	0
4	NA	A	404	1/1	0.99	0.17	26,26,26,26	0
3	GOL	B	408[C]	6/6	0.99	0.07	6,14,18,18	6
3	GOL	A	402	1/6	0.99	0.04	15,15,15,15	0
3	GOL	A	409	6/6	0.99	0.04	12,14,15,19	0
4	NA	B	402	1/1	0.99	0.10	26,26,26,26	0
3	GOL	B	411	6/6	0.99	0.07	11,14,20,23	0
3	GOL	B	408[A]	6/6	0.99	0.07	7,14,18,24	6
3	GOL	B	408[B]	6/6	0.99	0.07	8,12,13,14	6
2	ZN	B	401	1/1	1.00	0.01	6,6,6,6	0
2	ZN	A	401	1/1	1.00	0.01	7,7,7,7	0
4	NA	B	406	1/1	1.00	0.02	12,12,12,12	0
5	CL	A	415	1/1	1.00	0.10	20,20,20,20	0
5	CL	A	416	1/1	1.00	0.04	15,15,15,15	1
5	CL	A	417	1/1	1.00	0.03	14,14,14,14	0
5	CL	A	418	1/1	1.00	0.02	18,18,18,18	0
5	CL	A	419	1/1	1.00	0.06	18,18,18,18	0
5	CL	B	419	1/1	1.00	0.03	12,12,12,12	0
5	CL	B	420	1/1	1.00	0.02	14,14,14,14	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	CL	B	421	1/1	1.00	0.01	13,13,13,13	0
5	CL	B	422	1/1	1.00	0.03	15,15,15,15	0
5	CL	B	423	1/1	1.00	0.06	16,16,16,16	0
6	CA	A	420	1/1	1.00	0.05	8,8,8,8	1
6	CA	B	424	1/1	1.00	0.05	7,7,7,7	1

6.5 Other polymers [i](#)

There are no such residues in this entry.