



## wwPDB X-ray Structure Validation Summary Report

Nov 5, 2023 – 09:54 AM EST

PDB ID : 4IYO  
Title : Crystal structure of cystathionine gamma lyase from *Xanthomonas oryzae* pv. *oryzae* (XometC) in complex with E-site serine, A-site serine, A-site external aldimine structure with aminoacrylate and A-site iminopropionate intermediates  
Authors : Ngo, H.P.T.; Kim, J.K.; Kang, L.W.  
Deposited on : 2013-01-29  
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	397	 79% 15% . .
2	B	397	 82% 12% . .
2	C	397	 2% 82% 12% . .
2	D	397	 1% 84% 10% . .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SER	D	402	-	-	X	-

## 2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 13006 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Cystathionine gamma-lyase-like protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	381	2881	1825	506	535	15	0	5	0

- Molecule 2 is a protein called Cystathionine gamma-lyase-like protein, LYS201A modified.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	P				S
2	B	381	2890	1828	506	540	1	15	0	4	0
2	C	381	2890	1828	506	540	1	15	0	4	0
2	D	384	2924	1846	515	547	1	15	0	5	0

- Molecule 3 is SERINE (three-letter code: SER) (formula: C<sub>3</sub>H<sub>7</sub>NO<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			7	3	1	3		
3	B	1	Total	C	N	O	0	0
			7	3	1	3		
3	C	1	Total	C	N	O	0	0
			7	3	1	3		
3	D	1	Total	C	N	O	0	0
			7	3	1	3		
3	D	1	Total	C	N	O	0	0
			7	3	1	3		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	C	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	O	S			
			5	4	1	0	0	

- Molecule 6 is 2-[(E)-{3-hydroxy-2-methyl-5-[(phosphonoxy)methyl]pyridin-4-yl}methylidene]amino}prop-2-enoic acid (three-letter code: 0JO) (formula: C<sub>11</sub>H<sub>13</sub>N<sub>2</sub>O<sub>7</sub>P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	P		
			21	11	2	7	1	0	0

- Molecule 7 is AMINO-ACRYLATE (three-letter code: NAK) (formula: C<sub>3</sub>H<sub>5</sub>NO<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
7	B	1	6	3	1	2	0	0
7	C	1	6	3	1	2	0	0

- Molecule 8 is PYRUVIC ACID (three-letter code: PYR) (formula: C<sub>3</sub>H<sub>4</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
8	B	1	6	3	3	0	0

- Molecule 9 is water.




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	324	Total 324	O 324	0	0
9	B	365	Total 365	O 365	0	0
9	C	300	Total 300	O 300	0	0
9	D	335	Total 335	O 335	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cystathionine gamma-lyase-like protein



- Molecule 2: Cystathionine gamma-lyase-like protein, LYS201A modified





- Molecule 2: Cystathionine gamma-lyase-like protein, LYS201A modified



ASN

- Molecule 2: Cystathionine gamma-lyase-like protein, LYS201A modified

Chain D:  %



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	76.32Å 86.34Å 226.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.82 – 1.80 33.82 – 1.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (33.82-1.80) 100.0 (33.82-1.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.26 (at 1.81Å)	Xtrriage
Refinement program	REFMAC 5.5.0110	Depositor
R, $R_{free}$	0.136 , 0.180 0.137 , 0.181	Depositor DCC
$R_{free}$ test set	6971 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.7	Xtrriage
Anisotropy	0.025	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 48.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	13006	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.91% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NAK, LLP, SO4, GOL, PYR, OJO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.32	12/2954 (0.4%)	1.15	16/4007 (0.4%)
2	B	1.32	7/2935 (0.2%)	1.07	8/3982 (0.2%)
2	C	1.34	10/2935 (0.3%)	1.09	13/3982 (0.3%)
2	D	1.29	10/2972 (0.3%)	1.09	11/4031 (0.3%)
All	All	1.32	39/11796 (0.3%)	1.10	48/16002 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

The worst 5 of 39 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	233	GLU	CG-CD	7.50	1.63	1.51
2	D	364	GLU	CD-OE2	6.79	1.33	1.25
2	D	70	GLU	CB-CG	6.50	1.64	1.52
2	B	69	TYR	CD2-CE2	6.46	1.49	1.39
2	C	344	GLU	CG-CD	6.38	1.61	1.51

The worst 5 of 48 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	176	ARG	NE-CZ-NH1	-12.59	114.00	120.30
2	B	121	ARG	NE-CZ-NH2	-10.16	115.22	120.30
2	C	363	ARG	NE-CZ-NH2	-10.02	115.29	120.30
1	A	176	ARG	NE-CZ-NH2	9.94	125.27	120.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	176	ARG	NE-CZ-NH2	-9.50	115.55	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	206	HIS	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2881	0	2906	48	0
2	B	2890	0	2896	45	0
2	C	2890	0	2897	31	0
2	D	2924	0	2929	38	0
3	A	7	0	4	2	0
3	B	7	0	4	0	0
3	C	7	0	4	0	0
3	D	14	0	8	9	0
4	A	6	0	8	0	0
4	C	6	0	8	1	0
4	D	6	0	8	3	0
5	A	5	0	0	0	0
6	A	21	0	9	4	0
7	B	6	0	0	2	0
7	C	6	0	0	2	0
8	B	6	0	0	0	0
9	A	324	0	0	11	0
9	B	365	0	0	7	0
9	C	300	0	0	15	0
9	D	335	0	0	15	0
All	All	13006	0	11681	163	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 163 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:124:ARG:NH1	1:A:124:ARG:HB3	1.30	1.38
1:A:370:ASP:HB2	9:A:775:HOH:O	1.19	1.36
1:A:124:ARG:HH11	1:A:124:ARG:CB	1.38	1.36
2:D:210:LLP:C4'	3:D:402:SER:N	1.96	1.28
2:C:210:LLP:C4'	7:C:402:NAK:N	2.02	1.21

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	384/397 (97%)	373 (97%)	10 (3%)	1 (0%)	41 27
2	B	382/397 (96%)	374 (98%)	8 (2%)	0	100 100
2	C	382/397 (96%)	374 (98%)	7 (2%)	1 (0%)	41 27
2	D	386/397 (97%)	380 (98%)	6 (2%)	0	100 100
All	All	1534/1588 (97%)	1501 (98%)	31 (2%)	2 (0%)	51 36

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	339	SER
1	A	339	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	303/313 (97%)	291 (96%)	12 (4%)	31	16
2	B	301/312 (96%)	294 (98%)	7 (2%)	50	37
2	C	301/312 (96%)	294 (98%)	7 (2%)	50	37
2	D	305/312 (98%)	301 (99%)	4 (1%)	69	62
All	All	1210/1249 (97%)	1180 (98%)	30 (2%)	49	34

5 of 30 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	147	ARG
2	D	173	VAL
2	B	291	PRO
2	D	386	ARG
2	C	326	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	228	ASN
2	D	240	ASN
2	D	349	HIS
2	B	246	GLN
2	B	240	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the



expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	LLP	C	210	2	23,24,25	2.62	9 (39%)	25,32,34	2.18	7 (28%)
2	LLP	B	210	2	23,24,25	2.74	11 (47%)	25,32,34	2.16	9 (36%)
2	LLP	D	210	2	23,24,25	2.35	8 (34%)	25,32,34	1.97	8 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LLP	C	210	2	-	6/16/17/19	0/1/1/1
2	LLP	B	210	2	-	4/16/17/19	0/1/1/1
2	LLP	D	210	2	-	7/16/17/19	0/1/1/1

The worst 5 of 28 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	210	LLP	C3-C2	7.67	1.48	1.40
2	C	210	LLP	C3-C2	6.67	1.47	1.40
2	C	210	LLP	C4-C5	5.44	1.48	1.42
2	B	210	LLP	C4-C5	5.33	1.48	1.42
2	D	210	LLP	C3-C2	5.26	1.46	1.40

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	210	LLP	OP4-C5'-C5	6.14	121.05	109.35
2	C	210	LLP	OP4-C5'-C5	5.33	119.50	109.35
2	C	210	LLP	C3-C4-C5	-4.53	114.78	118.26
2	D	210	LLP	CE-NZ-C4'	4.32	132.16	118.90
2	B	210	LLP	OP2-P-OP4	-4.24	95.46	106.73

There are no chirality outliers.

5 of 17 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	210	LLP	O-C-CA-CB

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	C	210	LLP	O-C-CA-CB
2	D	210	LLP	O-C-CA-CB
2	B	210	LLP	CG-CD-CE-NZ
2	C	210	LLP	CG-CD-CE-NZ

There are no ring outliers.

3 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	210	LLP	2	0
2	B	210	LLP	2	0
2	D	210	LLP	7	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	GOL	D	403	-	5,5,5	0.41	0	5,5,5	0.82	0
3	SER	B	401	-	5,6,6	0.99	0	5,7,7	1.57	1 (20%)
3	SER	D	402	-	5,6,6	1.55	1 (20%)	5,7,7	0.72	0
7	NAK	C	402	-	4,5,5	1.23	1 (25%)	4,6,6	3.69	2 (50%)
8	PYR	B	403	-	5,5,5	2.74	2 (40%)	3,6,6	2.20	2 (66%)
6	OJO	A	404	-	20,21,21	2.72	7 (35%)	23,30,30	2.00	6 (26%)
3	SER	C	401	-	5,6,6	1.10	1 (20%)	5,7,7	1.97	2 (40%)
4	GOL	C	403	-	5,5,5	0.45	0	5,5,5	1.21	0
3	SER	D	401	-	5,6,6	0.77	0	5,7,7	1.24	1 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	SO4	A	403	-	4,4,4	0.43	0	6,6,6	0.35	0
4	GOL	A	402	-	5,5,5	0.66	0	5,5,5	0.25	0
3	SER	A	401	-	5,6,6	1.07	1 (20%)	5,7,7	1.23	1 (20%)
7	NAK	B	402	-	4,5,5	1.05	0	4,6,6	3.07	3 (75%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	D	403	-	-	4/4/4/4	-
3	SER	B	401	-	-	0/6/6/6	-
3	SER	D	402	-	-	2/6/6/6	-
7	NAK	C	402	-	-	2/2/4/4	-
8	PYR	B	403	-	-	0/4/4/4	-
6	OJO	A	404	-	-	0/10/15/15	0/1/1/1
3	SER	C	401	-	-	0/6/6/6	-
4	GOL	C	403	-	-	2/4/4/4	-
3	SER	D	401	-	-	1/6/6/6	-
4	GOL	A	402	-	-	0/4/4/4	-
3	SER	A	401	-	-	2/6/6/6	-
7	NAK	B	402	-	-	2/2/4/4	-

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	404	OJO	C4-C3	6.70	1.50	1.40
6	A	404	OJO	C3-C2	5.87	1.46	1.40
6	A	404	OJO	C4A-N	4.85	1.34	1.28
8	B	403	PYR	O-C	4.65	1.35	1.22
6	A	404	OJO	C2A-C2	3.52	1.56	1.50

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	C	402	NAK	CAE-CA-C	6.35	124.34	118.17
6	A	404	OJO	C3-C4-C5	-5.73	113.86	118.26
7	B	402	NAK	CAE-CA-C	4.50	122.54	118.17
3	C	401	SER	OXT-C-O	-3.63	115.85	124.09

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	C	402	NAK	O-C-CA	-3.53	116.87	121.38

There are no chirality outliers.

5 of 15 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	D	402	SER	N-CA-CB-OG
4	C	403	GOL	C1-C2-C3-O3
4	D	403	GOL	O1-C1-C2-C3
7	B	402	NAK	OXT-C-CA-CAE
7	B	402	NAK	O-C-CA-CAE

There are no ring outliers.

8 monomers are involved in 23 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	403	GOL	3	0
3	D	402	SER	7	0
7	C	402	NAK	2	0
6	A	404	OJO	4	0
4	C	403	GOL	1	0
3	D	401	SER	2	0
3	A	401	SER	2	0
7	B	402	NAK	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	381/397 (95%)	-0.63	1 (0%) 94 92	9, 14, 26, 37	0
2	B	380/397 (95%)	-0.66	1 (0%) 94 92	9, 14, 25, 39	0
2	C	380/397 (95%)	-0.33	8 (2%) 63 59	10, 17, 34, 49	0
2	D	383/397 (96%)	-0.57	2 (0%) 91 89	10, 15, 26, 55	0
All	All	1524/1588 (95%)	-0.55	12 (0%) 86 84	9, 15, 29, 55	0

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	14	ALA	4.7
2	B	14	ALA	4.4
2	C	361	ALA	3.7
2	C	394	VAL	3.5
2	C	365	GLN	3.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	LLP	D	210	24/25	0.96	0.12	10,16,28,30	0
2	LLP	C	210	24/25	0.97	0.13	11,17,26,27	0
2	LLP	B	210	24/25	0.97	0.12	9,17,24,25	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	D	403	6/6	0.92	0.25	17,41,41,44	0
4	GOL	C	403	6/6	0.93	0.33	26,34,41,53	0
3	SER	A	401	7/7	0.93	0.10	26,30,30,31	0
7	NAK	C	402	6/6	0.93	0.18	27,33,38,38	0
8	PYR	B	403	6/6	0.94	0.11	18,23,30,34	0
3	SER	C	401	7/7	0.95	0.08	30,31,32,33	0
3	SER	D	401	7/7	0.95	0.08	21,23,26,27	0
3	SER	D	402	7/7	0.96	0.11	18,21,24,30	0
4	GOL	A	402	6/6	0.96	0.06	17,18,20,20	0
3	SER	B	401	7/7	0.96	0.08	19,20,23,26	0
7	NAK	B	402	6/6	0.97	0.13	20,24,31,32	0
6	OJO	A	404	21/21	0.98	0.09	11,19,25,26	0
5	SO4	A	403	5/5	0.99	0.14	32,37,41,41	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.