



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 19, 2024 – 03:08 AM EST

PDB ID : 4J8N  
Title : Aurora A Kinase Apo  
Authors : Meyerowitz, J.G.; Gustafson, W.C.; Shokat, K.M.; Weiss, W.A.  
Deposited on : 2013-02-14  
Resolution : 3.13 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1626 (3.18-3.10)
Clashscore	141614	1735 (3.18-3.10)
Ramachandran outliers	138981	1677 (3.18-3.10)
Sidechain outliers	138945	1677 (3.18-3.10)
RSRZ outliers	127900	1588 (3.18-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	279	 76% 17% • 5%
1	B	279	 80% 14% • 5%
1	C	279	 82% 13% • 5%
1	D	279	 81% 14% • 5%

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 17546 atoms, of which 8807 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aurora kinase A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	266	4390	1403	2205	387	389	6	0	0	0
1	B	266	4390	1403	2205	387	389	6	0	0	0
1	C	266	4390	1403	2205	387	389	6	0	0	0
1	D	265	4368	1397	2192	385	388	6	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	287	ASP	THR	conflict	UNP O14965
B	287	ASP	THR	conflict	UNP O14965
C	287	ASP	THR	conflict	UNP O14965
D	287	ASP	THR	conflict	UNP O14965


- Molecule 2 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	4	Total 4 O 4	0	0
2	C	2	Total 2 O 2	0	0
2	D	2	Total 2 O 2	0	0

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Aurora kinase A

Chain A: 





- Molecule 1: Aurora kinase A

Chain B: 





- Molecule 1: Aurora kinase A

Chain C: 



- Molecule 1: Aurora kinase A

Chain D: 





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	83.78Å 83.78Å 171.78Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	44.95 – 3.13 44.95 – 3.13	Depositor EDS
% Data completeness (in resolution range)	99.8 (44.95-3.13) 97.1 (44.95-3.13)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.28 (at 3.12Å)	Xtrriage
Refinement program	PHENIX 1.8_1069	Depositor
R, $R_{free}$	0.184 , 0.234 0.196 , 0.243	Depositor DCC
$R_{free}$ test set	2033 reflections (8.59%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	93.6	Xtrriage
Anisotropy	0.012	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 64.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtrriage
Estimated twinning fraction	0.480 for -h,-k,l 0.487 for h,-h-k,-l 0.488 for -k,-h,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	17546	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	114.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.32% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.29	0/2240	0.52	0/3027
1	B	0.26	0/2240	0.48	0/3027
1	C	0.26	0/2240	0.48	0/3027
1	D	0.26	0/2231	0.49	0/3016
All	All	0.27	0/8951	0.49	0/12097

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2185	2205	2195	39	0
1	B	2185	2205	2195	25	0
1	C	2185	2205	2195	25	0
1	D	2176	2192	2182	29	0
2	A	4	0	0	0	0
2	C	2	0	0	2	0
2	D	2	0	0	0	0
All	All	8739	8807	8767	111	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 111 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:173:GLY:O	1:A:174:VAL:HG22	1.53	1.09
1:C:367:ASN:OD1	1:C:369:SER:OG	1.99	0.80
1:B:367:ASN:OD1	1:B:369:SER:OG	2.01	0.79
1:A:173:GLY:O	1:A:174:VAL:CG2	2.33	0.75
1:D:367:ASN:OD1	1:D:369:SER:OG	2.04	0.74

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	264/279 (95%)	245 (93%)	14 (5%)	5 (2%)	8	31
1	B	264/279 (95%)	248 (94%)	14 (5%)	2 (1%)	19	53
1	C	264/279 (95%)	248 (94%)	14 (5%)	2 (1%)	19	53
1	D	263/279 (94%)	248 (94%)	14 (5%)	1 (0%)	34	67
All	All	1055/1116 (94%)	989 (94%)	56 (5%)	10 (1%)	17	50

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	174	VAL
1	B	292	THR
1	A	291	GLY
1	A	292	THR
1	B	287	ASP



### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	235/247 (95%)	228 (97%)	7 (3%)	41	70
1	B	235/247 (95%)	231 (98%)	4 (2%)	60	82
1	C	235/247 (95%)	231 (98%)	4 (2%)	60	82
1	D	234/247 (95%)	231 (99%)	3 (1%)	69	86
All	All	939/988 (95%)	921 (98%)	18 (2%)	57	80

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	293	LEU
1	D	277	TRP
1	D	271	LYS
1	B	221	GLU
1	C	277	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	366	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	266/279 (95%)	0.11	2 (0%) 86 74	69, 98, 152, 168	0
1	B	266/279 (95%)	0.14	5 (1%) 66 48	69, 98, 151, 181	0
1	C	266/279 (95%)	0.11	3 (1%) 80 66	69, 98, 150, 183	0
1	D	265/279 (94%)	0.14	4 (1%) 73 56	68, 96, 150, 169	0
All	All	1063/1116 (95%)	0.13	14 (1%) 77 61	68, 98, 152, 183	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	185	GLN	4.3
1	C	161	LEU	3.0
1	D	279	VAL	2.9
1	A	161	LEU	2.7
1	A	185	GLN	2.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.