



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 13, 2024 – 10:09 AM EDT

PDB ID : 4JAR
Title : Crystal structure of mycobacterium tuberculosis pks11 in complex with polyketide intermediates and evidence that it synthesizes ALKYL PYRONES
Authors : Gokulan, K.; Sacchettini, J.C.; Mycobacterium Tuberculosis Structural Proteomics Project (XMTB)
Deposited on : 2013-02-19
Resolution : 1.98 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.20.1
EDS : 2.36.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	11647 (2.00-1.96)
Clashscore	141614	1014 (1.98-1.98)
Ramachandran outliers	138981	1006 (1.98-1.98)
Sidechain outliers	138945	1006 (1.98-1.98)
RSRZ outliers	127900	11410 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	353	
1	B	353	
1	C	353	
1	D	353	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 10576 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Alpha-pyrone synthesis polyketide synthase-like Pks11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	352	Total 2642	C 1671	N 469	O 494	S 8	0	0	0
1	C	352	Total 2642	C 1671	N 469	O 494	S 8	0	0	0
1	B	353	Total 2650	C 1676	N 470	O 495	S 9	0	0	0
1	D	352	Total 2642	C 1671	N 469	O 494	S 8	0	0	0


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alpha-pyrone synthesis polyketide synthase-like Pks11




- Molecule 1: Alpha-pyrone synthesis polyketide synthase-like Pks11






- Molecule 1: Alpha-pyrone synthesis polyketide synthase-like Pks11



- Molecule 1: Alpha-pyrone synthesis polyketide synthase-like Pks11



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	72.60Å 48.52Å 195.23Å 90.00° 98.41° 90.00°	Depositor
Resolution (Å)	38.24 – 1.98 38.24 – 1.98	Depositor EDS
% Data completeness (in resolution range)	70.0 (38.24-1.98) 70.3 (38.24-1.98)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.22 (at 1.98Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE: 1.6_289)	Depositor
R, R_{free}	0.268 , 0.314 0.306 , 0.320	Depositor DCC
R_{free} test set	3377 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	48.7	Xtrriage
Anisotropy	0.288	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 46.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.046 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	10576	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.62	1/2695 (0.0%)	0.70	2/3671 (0.1%)
1	B	0.54	1/2703 (0.0%)	0.70	5/3681 (0.1%)
1	C	0.67	1/2695 (0.0%)	0.73	3/3671 (0.1%)
1	D	0.60	3/2695 (0.1%)	0.71	4/3671 (0.1%)
All	All	0.61	6/10788 (0.1%)	0.71	14/14694 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
All	All	0	3

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	295	PRO	N-CD	5.52	1.55	1.47
1	D	328	PRO	N-CD	5.48	1.55	1.47
1	D	294	PRO	N-CD	5.41	1.55	1.47
1	A	29	PRO	N-CD	5.34	1.55	1.47
1	B	328	PRO	N-CD	5.19	1.55	1.47

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	328	PRO	C-N-CD	6.54	142.13	128.40
1	C	294	PRO	C-N-CD	6.04	141.08	128.40
1	C	328	PRO	C-N-CD	5.82	140.61	128.40
1	D	328	PRO	C-N-CD	5.78	140.53	128.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	327	ARG	C-N-CD	5.73	140.43	128.40

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	272	GLY	Peptide
1	B	2	SER	Peptide
1	C	344	CYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2642	0	2675	146	0
1	B	2650	0	2687	124	0
1	C	2642	0	2675	121	1
1	D	2642	0	2675	157	1
All	All	10576	0	10712	531	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

The worst 5 of 531 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:28:PHE:HB2	1:B:31:LEU:CD1	1.46	1.44
1:C:251:ARG:HG2	1:C:252:TYR:CE1	1.59	1.38
1:A:300:LEU:CD1	1:A:323:THR:HG22	1.58	1.34
1:B:28:PHE:CB	1:B:31:LEU:HD12	1.70	1.22
1:B:84:LEU:HD23	1:B:122:LEU:CD1	1.70	1.21

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:299:GLU:OE1	1:D:303:ARG:NH1[1_455]	2.07	0.13

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	350/353 (99%)	319 (91%)	27 (8%)	4 (1%)	14	5
1	B	351/353 (99%)	320 (91%)	25 (7%)	6 (2%)	9	2
1	C	350/353 (99%)	304 (87%)	38 (11%)	8 (2%)	6	1
1	D	350/353 (99%)	322 (92%)	21 (6%)	7 (2%)	7	1
All	All	1401/1412 (99%)	1265 (90%)	111 (8%)	25 (2%)	8	1

5 of 25 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	278	PRO
1	B	330	SER
1	D	205	GLN
1	D	206	VAL
1	D	352	TRP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	280/281 (100%)	223 (80%)	57 (20%)	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	281/281 (100%)	236 (84%)	45 (16%)	2	0
1	C	280/281 (100%)	226 (81%)	54 (19%)	1	0
1	D	280/281 (100%)	234 (84%)	46 (16%)	2	0
All	All	1121/1124 (100%)	919 (82%)	202 (18%)	1	0

5 of 202 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	19	SER
1	B	296	GLU
1	D	350	LEU
1	B	100	MET
1	B	221	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	92	ASN
1	D	318	HIS
1	B	69	ASN
1	B	205	GLN
1	B	247	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	352/353 (99%)	0.76	32 (9%) 9 10	24, 63, 79, 96	0
1	B	353/353 (100%)	0.80	46 (13%) 3 3	24, 61, 81, 94	0
1	C	352/353 (99%)	0.71	41 (11%) 4 5	22, 59, 83, 92	0
1	D	352/353 (99%)	0.80	40 (11%) 5 5	24, 63, 87, 96	0
All	All	1409/1412 (99%)	0.77	159 (11%) 5 5	22, 62, 83, 96	0

The worst 5 of 159 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	MET	4.8
1	B	133	LEU	4.7
1	D	271	ILE	4.6
1	C	104	ALA	4.3
1	D	294	PRO	4.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.