

wwPDB X-ray Structure Validation Summary Report (i)

Nov 7, 2023 – 06:10 AM EST

PDB ID : 4JGT

Title : Structure and kinetic analysis of H2S production by human Mercaptopyruvate

Sulfurtransferase

Authors: Koutmos, M.; Yamada, K.; Yadav, P.K.; Chiku, T.; Banerjee, R.

Deposited on : 2013-03-03

Resolution : 2.16 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

 $Mol Probity \quad : \quad 4.02b\text{--}467$

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.36

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	299	89%	•	6%
1	В	299	85 %	7%	7%
1	С	299	84%	8%	7%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 7036 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called 3-mercaptopyruvate sulfurtransferase.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	280	Total	С	N	О	S	0	0	0
1	A	280	2219	1407	398	406	8	0		
1	D	278	Total	С	N	О	S	0	0	0
1	Б	210	2203	1398	393	404	8	0	0	
1	С	277	Total	С	N	О	S	0	0	0
1		C 277		1393	394	403	8	U	U	

There are 60 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-9	MET	-	expression tag	UNP P25325
A	-8	GLY	-	expression tag	UNP P25325
A	-7	SER	-	expression tag	UNP P25325
A	-6	SER	-	expression tag	UNP P25325
A	-5	HIS	-	expression tag	UNP P25325
A	-4	HIS	-	expression tag	UNP P25325
A	-3	HIS	ı	expression tag	UNP P25325
A	-2	HIS	-	expression tag	UNP P25325
A	-1	HIS	-	expression tag	UNP P25325
A	0	HIS	-	expression tag	UNP P25325
A	1	SER	-	expression tag	UNP P25325
A	2	SER	-	expression tag	UNP P25325
A	3	GLY	-	expression tag	UNP P25325
A	4	LEU	-	expression tag	UNP P25325
A	5	VAL	-	expression tag	UNP P25325
A	6	PRO	-	expression tag	UNP P25325
A	7	ARG	-	expression tag	UNP P25325
A	8	GLY	-	expression tag	UNP P25325
A	9	SER	-	expression tag	UNP P25325
A	10	HIS	-	expression tag	UNP P25325
В	-9	MET	=	expression tag	UNP P25325
В	-8	GLY	=	expression tag	UNP P25325
В	-7	SER	-	expression tag	UNP P25325




 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
В	-6	SER	-	expression tag	UNP P25325
В	-5	HIS	-	expression tag	UNP P25325
В	-4	HIS	_	expression tag	UNP P25325
В	-3	HIS	-	expression tag	UNP P25325
В	-2	HIS	-	expression tag	UNP P25325
В	-1	HIS	_	expression tag	UNP P25325
В	0	HIS	_	expression tag	UNP P25325
В	1	SER	-	expression tag	UNP P25325
В	2	SER	-	expression tag	UNP P25325
В	3	GLY	-	expression tag	UNP P25325
В	4	LEU	-	expression tag	UNP P25325
В	5	VAL	-	expression tag	UNP P25325
В	6	PRO	-	expression tag	UNP P25325
В	7	ARG	-	expression tag	UNP P25325
В	8	GLY	-	expression tag	UNP P25325
В	9	SER	-	expression tag	UNP P25325
В	10	HIS	-	expression tag	UNP P25325
С	-9	MET	-	expression tag	UNP P25325
С	-8	GLY	-	expression tag	UNP P25325
С	-7	SER	-	expression tag	UNP P25325
С	-6	SER	-	expression tag	UNP P25325
С	-5	HIS	-	expression tag	UNP P25325
С	-4	HIS	-	expression tag	UNP P25325
С	-3	HIS	-	expression tag	UNP P25325
С	-2	HIS	-	expression tag	UNP P25325
С	-1	HIS	-	expression tag	UNP P25325
С	0	HIS	-	expression tag	UNP P25325
С	1	SER	-	expression tag	UNP P25325
С	2	SER	-	expression tag	UNP P25325
С	3	GLY	-	expression tag	UNP P25325
С	4	LEU	-	expression tag	UNP P25325
С	5	VAL	-	expression tag	UNP P25325
С	6	PRO	-	expression tag	UNP P25325
С	7	ARG	-	expression tag	UNP P25325
С	8	GLY	-	expression tag	UNP P25325
С	9	SER	-	expression tag	UNP P25325
С	10	HIS	-	expression tag	UNP P25325


 \bullet Molecule 2 is PYRUVIC ACID (three-letter code: PYR) (formula: $\mathrm{C_3H_4O_3}).$





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	В	1	Total C O 6 3 3	0	0
2	С	1	Total C O 6 3 3	0	0

 \bullet Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
3	A	1	Total 5	O 4	S 1	0	0




4JGT

 $Continued\ from\ previous\ page...$

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	О	S	0	0
3	A	1	5	4	1	0	U
3	A	1	Total	О	S	0	0
3	A	1	5	4	1	0	U
3	В	1	Total	О	S	0	0
3	Ъ	1	5	4	1	U	
3	С	1	Total	Ο	\mathbf{S}	0	0
		1	5	4	1	U	
3	С	1	Total	O	\mathbf{S}	0	0
		1	5	4	1	U	U
3	\mathbf{C}	1	Total	Ο	S	0	0
		ı ı	5	4	1		U
3		1	Total	О	S	0	0
3		1	5	4	1		U

• Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	С	1	Total C O 6 3 3	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	129	Total O 129 129	0	0



Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	107	Total O 107 107	0	0
5	С	116	Total O 116 116	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 3-mercaptopyruvate sulfurtransferase





4 Data and refinement statistics (i)

Property	Value	Source	
Space group	C 1 2 1	Depositor	
Cell constants	110.74Å 171.86Å 72.98Å	Donagitan	
a, b, c, α , β , γ	90.00° 117.18° 90.00°	Depositor	
Resolution (Å)	85.93 - 2.16	Depositor	
Resolution (A)	49.52 - 2.16	EDS	
% Data completeness	95.6 (85.93-2.16)	Depositor	
(in resolution range)	95.7 (49.52-2.16)	EDS	
R_{merge}	(Not available)	Depositor	
R_{sym}	0.07	Depositor	
$< I/\sigma(I) > 1$	1.83 (at 2.16Å)	Xtriage	
Refinement program	REFMAC 5.6.0117	Depositor	
D.D.	0.185 , 0.220	Depositor	
R, R_{free}	0.190 , 0.222	DCC	
R_{free} test set	3121 reflections (5.04%)	wwPDB-VP	
Wilson B-factor (\mathring{A}^2)	29.2	Xtriage	
Anisotropy	0.298	Xtriage	
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.36, 32.5	EDS	
L-test for twinning ²	$< L > = 0.49, < L^2> = 0.32$	Xtriage	
Estimated twinning fraction	0.023 for 1/2 *h + 1/2 *k + 1,3/2 *h - 1/2 *k + 1,-1	Xtriage	
<u> </u>	0.048 for 1/2*h-1/2*k+l,-3/2*h-1/2*k-l,-l		
F_o, F_c correlation	0.95	EDS	
Total number of atoms	7036	wwPDB-VP	
Average B, all atoms (\mathring{A}^2)	32.0	wwPDB-VP	

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.50% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PYR, SO4, CSS, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Boı	nd lengths	Bond angles		
10101		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.68	0/2278	0.75	1/3095~(0.0%)	
1	В	0.68	$1/2261 \ (0.0\%)$	0.73	2/3071 (0.1%)	
1	С	0.65	$1/2256 \ (0.0\%)$	0.72	1/3065 (0.0%)	
All	All	0.67	2/6795~(0.0%)	0.73	4/9231 (0.0%)	

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$Ideal(\AA)$
1	В	114	TRP	CD2-CE2	5.46	1.48	1.41
1	С	15	TRP	CD2-CE2	5.07	1.47	1.41

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	В	190	ARG	NE-CZ-NH1	7.19	123.90	120.30
1	В	190	ARG	NE-CZ-NH2	-5.48	117.56	120.30
1	С	273	ASP	CB-CG-OD1	5.27	123.04	118.30
1	A	21	ARG	NE-CZ-NH1	5.01	122.80	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2219	0	2127	7	0
1	В	2203	0	2108	9	0
1	С	2198	0	2099	15	0
2	A	6	0	0	2	0
2	В	6	0	0	0	0
2	С	6	0	0	1	0
3	A	15	0	0	0	0
3	В	5	0	0	0	0
3	С	20	0	0	1	0
4	С	6	0	8	0	0
5	A	129	0	0	0	0
5	В	107	0	0	0	0
5	С	116	0	0	1	0
All	All	7036	0	6342	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 30 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap} & (ext{Å}) \end{aligned}$	
1:C:188:ARG:NH2	2:C:401:PYR:OXT	2.23	0.71	
1:B:76:LEU:HD22	1:B:163:ILE:HD11	1.78	0.66	
1:B:76:LEU:CD2	1:B:163:ILE:HD11	2.31	0.61	
1:C:107:LEU:HD12	5:C:581:HOH:O	2.00	0.60	
1:A:62:ILE:HD11	1:A:113:VAL:HG23	1.85	0.57	

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed Favoured Allowed		Allowed	Outliers	Percentiles	
1	A	$277/299 \ (93\%)$	267 (96%)	10 (4%)	0	100	100
1	В	273/299 (91%)	265 (97%)	8 (3%)	0	100	100
1	С	272/299 (91%)	263 (97%)	9 (3%)	0	100	100
All	All	822/897 (92%)	795 (97%)	27 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	d Rotameric Outliers		Percentiles		
1	A	228/244 (93%)	225 (99%)	3 (1%)	69 74		
1	В	227/244 (93%)	217 (96%)	10 (4%)	28 25		
1	С	226/244 (93%)	224 (99%)	2 (1%)	78 83		
All	All	681/732 (93%)	666 (98%)	15 (2%)	52 55		

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	47	ARG
1	С	147	SER
1	В	52	ARG
1	С	275	SER
1	В	190	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type Chair	Chain	Chain Res	Res Link	Bond lengths			Bond angles		
MIOI		Chain			Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
1	CSS	С	248	1	4,6,7	0.70	0	1,6,8	0.51	0
1	CSS	В	248	1	4,6,7	0.97	0	1,6,8	0.44	0
1	CSS	A	248	1	4,6,7	0.68	0	1,6,8	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSS	С	248	1	-	0/1/5/7	-
1	CSS	В	248	1	-	0/1/5/7	-
1	CSS	A	248	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	С	248	CSS	1	0

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



5.6 Ligand geometry (i)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

NASI	Т	Clasica	Das	T : 1-	В	ond leng	$_{ m gths}$	В	ond ang	gles
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PYR	С	401	-	5,5,5	2.05	2 (40%)	3,6,6	2.09	1 (33%)
2	PYR	В	401	-	5,5,5	1.71	2 (40%)	3,6,6	2.06	2 (66%)
2	PYR	A	401	-	5,5,5	1.91	2 (40%)	3,6,6	2.18	1 (33%)
3	SO4	С	404	-	4,4,4	0.29	0	6,6,6	0.22	0
3	SO4	С	402	-	4,4,4	0.26	0	6,6,6	0.42	0
3	SO4	A	404	-	4,4,4	0.36	0	6,6,6	0.16	0
3	SO4	A	403	-	4,4,4	0.40	0	6,6,6	0.15	0
4	GOL	С	406	-	5,5,5	0.20	0	5,5,5	0.82	0
3	SO4	В	402	-	4,4,4	0.32	0	6,6,6	0.42	0
3	SO4	С	405	-	4,4,4	0.43	0	6,6,6	0.24	0
3	SO4	A	402	-	4,4,4	0.19	0	6,6,6	0.31	0
3	SO4	С	403	-	4,4,4	0.44	0	6,6,6	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PYR	A	401	-	-	1/4/4/4	-
2	PYR	С	401	-	-	0/4/4/4	-
4	GOL	С	406	-	-	4/4/4/4	-
2	PYR	В	401	-	-	0/4/4/4	-

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	Ideal(A)
2	С	401	PYR	CA-C	-3.60	1.41	1.54
2	A	401	PYR	CA-C	-3.43	1.41	1.54
2	В	401	PYR	CA-C	-3.16	1.42	1.54



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	$\operatorname{Ideal}(ext{\AA})$
2	С	401	PYR	OXT-C	-2.62	1.22	1.30
2	A	401	PYR	OXT-C	-2.36	1.23	1.30

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	С	401	PYR	O3-CA-CB	3.57	127.65	119.73
2	A	401	PYR	O3-CA-CB	3.21	126.85	119.73
2	В	401	PYR	OXT-C-CA	2.61	121.11	113.97
2	В	401	PYR	O3-CA-CB	2.41	125.06	119.73

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	С	406	GOL	C1-C2-C3-O3
4	С	406	GOL	O1-C1-C2-C3
4	С	406	GOL	O2-C2-C3-O3
2	A	401	PYR	OXT-C-CA-O3
4	С	406	GOL	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	С	401	PYR	1	0
2	A	401	PYR	2	0
3	С	403	SO4	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$		$OWAB(A^2)$	Q < 0.9	
1	A	279/299~(93%)	-0.25	1 (0%)	92	94	18, 28, 50, 63	0
1	В	277/299 (92%)	-0.19	2 (0%)	87	91	18, 30, 56, 71	0
1	С	276/299 (92%)	-0.23	3 (1%)	80	85	19, 31, 54, 83	0
All	All	832/897 (92%)	-0.22	6 (0%)	87	91	18, 29, 54, 83	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	231	HIS	2.7
1	С	150	ALA	2.6
1	С	148	GLN	2.3
1	A	288	VAL	2.2
1	С	155	ARG	2.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B\text{-factors}}({f A}^2)$	Q<0.9
1	CSS	A	248	7/8	0.98	0.09	19,21,22,29	0
1	CSS	В	248	7/8	0.98	0.09	19,20,22,32	0
1	CSS	С	248	7/8	0.98	0.08	18,20,23,32	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q<0.9
4	GOL	С	406	6/6	0.83	0.18	39,40,42,49	0
2	PYR	A	401	6/6	0.91	0.22	28,28,29,31	6
2	PYR	С	401	6/6	0.92	0.23	35,36,39,40	6
2	PYR	В	401	6/6	0.92	0.24	40,42,44,50	0
3	SO4	С	403	5/5	0.96	0.15	49,54,56,60	0
3	SO4	С	405	5/5	0.97	0.11	56,59,60,63	0
3	SO4	A	404	5/5	0.97	0.14	63,67,68,68	0
3	SO4	A	403	5/5	0.98	0.07	57,63,68,70	0
3	SO4	С	404	5/5	0.98	0.17	54,55,57,58	0
3	SO4	A	402	5/5	0.99	0.11	29,29,32,32	0
3	SO4	С	402	5/5	0.99	0.09	28,29,30,32	0
3	SO4	В	402	5/5	1.00	0.11	30,32,32,33	0

6.5 Other polymers (i)

There are no such residues in this entry.

