

Full wwPDB X-ray Structure Validation Report (i)

Sep 10, 2023 – 07:32 PM EDT

PDB ID : 4KZ5

Title: Crystal structure of AmpC beta-lactamase in complex with fragment 5 (N-{[

3-(2-chlorophenyl)-5-methyl-1,2-oxazol-4-yl|carbonyl}glycine)

Authors: Eidam, O.; Barelier, S.; Fish, I.; Shoichet, B.K.

Deposited on : 2013-05-29

Resolution : 1.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# ext{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	1509 (1.38-1.34)
Clashscore	141614	1551 (1.38-1.34)
Ramachandran outliers	138981	1530 (1.38-1.34)
Sidechain outliers	138945	1530 (1.38-1.34)
RSRZ outliers	127900	1487 (1.38-1.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	358	90%	10%
1	В	358	96%	•



2 Entry composition (i)


There are 4 unique types of molecules in this entry. The entry contains 6645 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Beta-lactamase.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	Λ	250	Total	С	N	О	S	0	25	0
1	A 358	390	2879	1870	480	521	8	0	20	
1	R	358	Total	С	N	О	S	0	19	0
1	В	390	2829	1836	475	510	8		12	U


• Molecule 2 is N-{[3-(2-chlorophenyl)-5-methyl-1,2-oxazol-4-yl]carbonyl}glycine (three-letter code: 1U3) (formula: C₁₃H₁₁ClN₂O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
2	Δ	1	Total	С	Cl	N	О	0	0	
	11	1	20	13	1	2	4			
9	D	1	Total	С	Cl	N	О	0	0	
	Б	1	20	13	1	2	4	U		

• Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
3	Λ	1	Total O P	0	0	
3	Λ	1	5 4 1		0	
3	Λ	1	Total O P	0	0	
3	Λ	1	5 4 1		U	
3	Λ	1	Total O P		0	
3	A	1	5 4 1	0	0	
3	В	1	Total O P	0	0	
3	Ъ	1	5 4 1		U	

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	440	Total O 440 440	0	17
4	В	437	Total O 437 437	0	10



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Beta-lactamase

Chain A:

90%

10%

Note that the state of the state



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	118.08Å 76.86Å 97.71Å	Donositor
a, b, c, α , β , γ	90.00° 116.26° 90.00°	Depositor
Resolution (Å)	45.45 - 1.35	Depositor
rtesolution (A)	45.45 - 1.35	EDS
% Data completeness	97.1 (45.45-1.35)	Depositor
(in resolution range)	97.2 (45.45-1.35)	EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	3.33 (at 1.35Å)	Xtriage
Refinement program	PHENIX 1.8_1069	Depositor
D.D.	0.135 , 0.156	Depositor
R, R_{free}	0.135 , 0.155	DCC
R_{free} test set	6366 reflections (3.82%)	wwPDB-VP
Wilson B-factor (Å ²)	13.7	Xtriage
Anisotropy	0.358	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.37, 49.6	EDS
L-test for twinning ²	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	6645	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 8.18% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, 1U3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.46	0/3009	0.64	0/4114	
1	В	0.47	0/2947	0.64	0/4030	
All	All	0.46	0/5956	0.64	0/8144	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2879	0	2873	21	0
1	В	2829	0	2804	7	0
2	A	20	0	10	0	0
2	В	20	0	10	0	0
3	A	15	0	0	0	0
3	В	5	0	0	0	0
4	A	440	0	0	3	0
4	В	437	0	0	1	0
All	All	6645	0	5697	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.



All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${f distance}({ m \AA})$	overlap (Å)
1:B:87:ASP:OD2	1:B:92:TYR:OH	2.05	0.72
1:A:240:PRO:HA	1:A:243:ILE:HD12	1.74	0.67
1:A:130:ASP:OD1	1:A:133[B]:ARG:NH2	2.34	0.59
1:A:296:ARG:NE	4:A:800:HOH:O	2.36	0.58
1:B:182:LEU:O	1:B:232:ARG:NH1	2.39	0.55
1:A:4:ALA:N	4:A:827:HOH:O	2.41	0.52
1:A:264[B]:ASP:OD2	1:A:282:SER:OG	2.24	0.48
1:A:276:TRP:CD2	1:A:277:PRO:HA	2.49	0.48
1:A:334[A]:LEU:HG	1:A:357:LEU:HD22	1.95	0.48
1:A:258:ARG:NH1	4:B:798:HOH:O	2.46	0.46
1:B:344:TYR:CZ	1:B:349:ARG:HG2	2.51	0.46
1:B:344:TYR:CE1	1:B:349:ARG:HG2	2.52	0.45
1:B:276:TRP:CD2	1:B:277:PRO:HA	2.51	0.45
1:A:7:GLN:HG2	1:A:11[B]:ILE:HD12	2.00	0.44
1:A:344:TYR:CZ	1:A:349:ARG:HG2	2.52	0.44
1:A:119:LEU:HA	1:A:151:ALA:HA	1.98	0.44
1:A:262:THR:O	1:A:265[A]:MET:HG3	2.18	0.44
1:A:256:GLN:HG3	4:A:567:HOH:O	2.17	0.43
1:A:87:ASP:OD2	1:A:92:TYR:OH	2.17	0.43
1:A:326[B]:VAL:HG12	1:A:337:VAL:HG22	2.01	0.43
1:B:33:ILE:HD12	1:B:33:ILE:N	2.34	0.42
1:A:243:ILE:HD13	1:A:252:ILE:HD12	2.02	0.41
1:A:287:SER:CB	1:A:350:VAL:HG21	2.51	0.41
1:A:316[A]:THR:HG22	1:A:325:TYR:CD1	2.56	0.41
1:A:17:THR:HB	1:A:18:PRO:HD3	2.01	0.41
1:A:33:ILE:HD12	1:A:33:ILE:N	2.35	0.41
1:B:334:LEU:HG	1:B:357:LEU:HD22	2.02	0.40
1:A:124:GLU:CD	1:A:124:GLU:H	2.25	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	ain Analysed Fav		Allowed	Outliers	Perce	entiles
1	A	377/358 (105%)	372 (99%)	5 (1%)	0	100	100
1	В	368/358 (103%)	362 (98%)	6 (2%)	0	100	100
All	All	745/716 (104%)	734 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	$300/292 \ (103\%)$	298 (99%)	2 (1%)	84	64
1	В	291/292 (100%)	289 (99%)	2 (1%)	84	64
All	All	591/584 (101%)	587 (99%)	4 (1%)	84	64

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	124	GLU
1	A	314	HIS
1	В	314	HIS
1	В	361	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mal Type Chain Des Lin		Link	Bond lengths			Bond angles			
MIOI	Type	Chain	Res	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PO4	В	402	-	4,4,4	0.91	0	6,6,6	0.39	0
3	PO4	A	404	-	4,4,4	0.57	0	6,6,6	0.64	0
2	1U3	A	401	-	18,21,21	2.77	2 (11%)	20,29,29	1.33	3 (15%)
3	PO4	A	402	-	4,4,4	0.83	0	6,6,6	0.80	0
3	PO4	A	403	-	4,4,4	0.97	0	6,6,6	0.54	0
2	1U3	В	401	-	18,21,21	2.26	2 (11%)	20,29,29	1.31	2 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1U3	В	401	-	-	3/7/13/13	0/2/2/2
2	1U3	A	401	-		3/7/13/13	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	$Ideal(\AA)$
2	A	401	1U3	C07-C04	-10.32	1.39	1.49
2	В	401	1U3	C07-C04	-8.20	1.41	1.49

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	Ideal(A)
2	A	401	1U3	C14-N16	4.46	1.43	1.33
2	В	401	1U3	C14-N16	3.87	1.42	1.33

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	A	401	1U3	C08-C07-C12	3.06	121.02	117.63
2	В	401	1U3	C04-C07-C12	-2.71	117.10	121.16
2	A	401	1U3	C04-C07-C12	-2.45	117.49	121.16
2	В	401	1U3	C08-C07-C04	2.25	123.64	119.46
2	A	401	1U3	O20-C18-C17	2.12	120.12	112.74

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401	1U3	N05-C04-C07-C08
2	В	401	1U3	N05-C04-C07-C08
2	В	401	1U3	C03-C04-C07-C08
2	A	401	1U3	C03-C04-C07-C08
2	A	401	1U3	C18-C17-N16-C14
2	В	401	1U3	C18-C17-N16-C14

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	358/358 (100%)	-0.16	16 (4%) 33 38	6, 13, 29, 47	7 (1%)
1	В	358/358 (100%)	-0.23	8 (2%) 62 68	6, 14, 26, 34	0
All	All	716/716 (100%)	-0.20	24 (3%) 45 51	6, 14, 28, 47	7 (0%)

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	293[A]	LEU	6.4
1	A	291[A]	ILE	5.4
1	A	290[B]	LYS	5.0
1	A	126	LYS	4.6
1	A	123	ASP	3.4
1	A	292[A]	ALA	3.4
1	A	241	LEU	3.4
1	В	206	GLY	3.3
1	A	289[B]	ASN	3.2
1	В	241	LEU	3.2
1	В	201	TRP	2.9
1	A	244	ASN	2.8
1	A	206	GLY	2.7
1	A	124	GLU	2.7
1	A	36	GLY	2.6
1	В	49	ALA	2.6
1	В	123	ASP	2.5
1	В	48	ILE	2.4
1	В	244	ASN	2.3
1	A	205	GLU	2.2
1	A	242	ASP	2.2
1	В	36	GLY	2.1
1	A	102	ASN	2.1
1	A	34	TYR	2.0



6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
3	PO4	A	404	5/5	0.71	0.26	28,28,29,31	5
2	1U3	A	401	20/20	0.86	0.14	26,31,36,37	20
3	PO4	В	402	5/5	0.88	0.14	51,52,55,56	5
2	1U3	В	401	20/20	0.92	0.08	26,30,36,37	0
3	PO4	A	403	5/5	0.93	0.16	51,52,54,57	5
3	PO4	A	402	5/5	0.95	0.08	23,26,27,34	5

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around 1U3 A 401: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)





6.5 Other polymers (i)

There are no such residues in this entry.

