



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 8, 2023 – 01:57 AM EDT

PDB ID : 4QO2  
Title : Crystal structure of rhomboid intramembrane protease GlpG in complex with peptide derived inhibitor Ac-IATA-cmk  
Authors : Zoll, S.; Strisovsky, K.  
Deposited on : 2014-06-19  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	200	 2% (red), 78% (green), 12% (yellow), 9% (grey)
2	B	6	 67% (green), 33% (yellow)

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 1664 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Rhomboid protease GlpG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	182	1462	985	235	230	12	7	1	0


There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	277	ALA	-	expression tag	UNP U6NA71
A	278	SER	-	expression tag	UNP U6NA71
A	279	LEU	-	expression tag	UNP U6NA71
A	280	GLU	-	expression tag	UNP U6NA71
A	281	ARG	-	expression tag	UNP U6NA71
A	282	GLU	-	expression tag	UNP U6NA71
A	283	ASN	-	expression tag	UNP U6NA71
A	284	LEU	-	expression tag	UNP U6NA71
A	285	TYR	-	expression tag	UNP U6NA71
A	286	PHE	-	expression tag	UNP U6NA71

- Molecule 2 is a protein called 6-AMINO-2-METHYL-1,7-DIHYDRO-8H-IMIDAZO[4,5-G]QUINAZOLIN-8-ONE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	6	29	19	4	6	0	0	1

- Molecule 3 is nonyl beta-D-glucopyranoside (three-letter code: BNG) (formula: C<sub>15</sub>H<sub>30</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			20	15	5		
3	A	1	Total	C	O	0	0
			11	10	1		
3	A	1	Total	C		0	0
			9	9			
3	A	1	Total	C		0	0
			6	6			
3	A	1	Total	C		0	0
			7	7			
3	A	1	Total	C		0	0
			7	7			
3	A	1	Total	C	O	0	0
			11	10	1		
3	A	1	Total	C	O	0	0
			11	10	1		
3	A	1	Total	C		0	0
			6	6			
3	A	1	Total	C		0	0
			5	5			
3	A	1	Total	C		0	0
			5	5			
3	A	1	Total	C		0	0
			8	8			
3	A	1	Total	C	O	0	0
			7	6	1		
3	A	1	Total	C	O	0	0
			9	8	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Cl 1 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	47	Total O 47 47	0	0
5	B	3	Total O 3 3	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	97.99Å 97.99Å 66.40Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.99 – 2.10 48.99 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.99-2.10) 99.9 (48.99-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.69 (at 2.10Å)	Xtrriage
Refinement program	REFMAC 5.8.0069	Depositor
R, $R_{free}$	0.163 , 0.180 0.181 , 0.196	Depositor DCC
$R_{free}$ test set	1066 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.6	Xtrriage
Anisotropy	0.388	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 76.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.084 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	1664	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.42% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BNG, 0QE, CL, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.56	0/1514	0.45	0/2058
2	B	2.88	1/25 (4.0%)	0.95	0/34
All	All	0.66	1/1539 (0.1%)	0.46	0/2092

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	5	ALA	C-O	13.92	1.49	1.23

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	246	GLY	Peptide
1	A	270	LEU	Peptide

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen



atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1462	0	1471	35	0
2	B	29	0	31	2	0
3	A	122	0	197	9	0
4	A	1	0	0	1	0
5	A	47	0	0	2	0
5	B	3	0	0	0	0
All	All	1664	0	1699	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:172:GLY:H	3:A:307:BNG:H7'1	1.33	0.91
1:A:171:SER:H	3:A:307:BNG:H6'2	1.42	0.83
1:A:246:GLY:N	1:A:247:MET:HE2	1.95	0.81
1:A:157:TRP:CZ3	1:A:208[B]:MET:SD	2.74	0.80
1:A:171:SER:N	3:A:307:BNG:H6'2	1.99	0.78
1:A:214:ARG:HB3	1:A:223:ILE:HD11	1.68	0.75
1:A:245:PHE:HB3	1:A:247:MET:CE	2.23	0.69
1:A:246:GLY:N	1:A:247:MET:CE	2.59	0.66
1:A:172:GLY:N	3:A:307:BNG:H7'1	2.10	0.60
1:A:146:PHE:CZ	2:B:2:ILE:HG21	2.37	0.59
1:A:245:PHE:HB3	1:A:247:MET:HE2	1.84	0.58
1:A:205:TYR:CE1	1:A:237:ILE:HG13	2.38	0.58
1:A:98:TRP:CE2	1:A:102:ILE:HD11	2.39	0.58
1:A:208[B]:MET:CE	1:A:225:LEU:HD21	2.34	0.58
1:A:145:HIS:CE1	1:A:154:ASN:HD22	2.22	0.57
1:A:246:GLY:H	1:A:247:MET:CE	2.18	0.56
1:A:145:HIS:HE1	1:A:154:ASN:HD22	1.55	0.54
3:A:307:BNG:H7'2	5:A:417:HOH:O	2.07	0.54
1:A:245:PHE:HB3	1:A:247:MET:HE3	1.89	0.54
3:A:307:BNG:C1	3:A:307:BNG:H3'1	2.39	0.53
1:A:171:SER:OG	3:A:307:BNG:H2'2	2.08	0.52
1:A:205:TYR:CD1	1:A:208[A]:MET:SD	3.03	0.52
1:A:145:HIS:CE1	1:A:200:LEU:H	2.29	0.49
1:A:208[B]:MET:HE3	1:A:225:LEU:HD21	1.93	0.49
1:A:172:GLY:H	3:A:307:BNG:C7'	2.15	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:208[B]:MET:HE1	1:A:225:LEU:HD21	1.96	0.47
1:A:208[B]:MET:CE	1:A:233:ALA:CB	2.93	0.47
1:A:208[B]:MET:CE	1:A:233:ALA:HB2	2.46	0.45
1:A:214:ARG:CB	1:A:223:ILE:HD11	2.43	0.45
1:A:145:HIS:HE1	1:A:200:LEU:H	1.65	0.44
1:A:145:HIS:HD2	5:A:404:HOH:O	2.00	0.43
3:A:306:BNG:H9'1	3:A:314:BNG:H9'1	2.01	0.42
1:A:208[B]:MET:HE3	1:A:225:LEU:CD2	2.50	0.41
1:A:208[B]:MET:HE2	1:A:233:ALA:CB	2.50	0.41
1:A:150:HIS:HB2	4:A:315:CL:CL	2.58	0.41
1:A:211:VAL:HA	1:A:223:ILE:HD12	2.02	0.41
1:A:160:TYR:CE2	1:A:229:LEU:CD1	3.04	0.40
1:A:146:PHE:CE1	2:B:2:ILE:HG21	2.55	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	181/200 (90%)	177 (98%)	4 (2%)	0	100	100
2	B	3/6 (50%)	3 (100%)	0	0	100	100
All	All	184/206 (89%)	180 (98%)	4 (2%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	147/160 (92%)	143 (97%)	4 (3%)	44	48
2	B	2/2 (100%)	2 (100%)	0	100	100
All	All	149/162 (92%)	145 (97%)	4 (3%)	44	48

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	92	ARG
1	A	232	PHE
1	A	247	MET
1	A	248	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	145	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 1 is monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	BNG	A	303	-	8,8,21	0.22	0	7,7,26	0.44	0
3	BNG	A	314	-	7,7,21	0.28	0	6,6,26	0.45	0
3	BNG	A	312	-	7,7,21	0.33	0	6,6,26	0.45	0
3	BNG	A	301	-	20,20,21	0.50	0	23,24,26	0.69	0
3	BNG	A	313	-	5,5,21	0.26	0	4,4,26	0.40	0
3	BNG	A	309	-	5,5,21	0.27	0	4,4,26	0.40	0
3	BNG	A	306	-	6,6,21	0.24	0	5,5,26	0.46	0
3	BNG	A	308	-	10,10,21	0.42	0	9,9,26	0.46	0
3	BNG	A	311	-	4,4,21	0.25	0	3,3,26	0.41	0
3	BNG	A	302	-	10,10,21	0.36	0	9,9,26	0.45	0
3	BNG	A	310	-	4,4,21	0.28	0	3,3,26	0.40	0
3	BNG	A	305	-	6,6,21	0.32	0	5,5,26	0.41	0
3	BNG	A	304	-	5,5,21	0.34	0	4,4,26	0.35	0
3	BNG	A	307	-	10,10,21	0.43	0	9,9,26	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BNG	A	303	-	-	5/6/6/32	-
3	BNG	A	314	-	-	1/5/5/32	-
3	BNG	A	312	-	-	3/5/5/32	-
3	BNG	A	301	-	-	6/12/28/32	0/1/1/1
3	BNG	A	313	-	-	0/3/3/32	-
3	BNG	A	309	-	-	3/3/3/32	-
3	BNG	A	306	-	-	3/4/4/32	-
3	BNG	A	308	-	-	3/8/8/32	-
3	BNG	A	311	-	-	0/2/2/32	-
3	BNG	A	302	-	-	4/8/8/32	-
3	BNG	A	310	-	-	0/2/2/32	-
3	BNG	A	305	-	-	1/4/4/32	-
3	BNG	A	304	-	-	0/3/3/32	-
3	BNG	A	307	-	-	7/8/8/32	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (36) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	307	BNG	C3'-C4'-C5'-C6'
3	A	307	BNG	C2'-C3'-C4'-C5'
3	A	303	BNG	C4'-C5'-C6'-C7'
3	A	308	BNG	C2'-C3'-C4'-C5'
3	A	312	BNG	C3'-C4'-C5'-C6'
3	A	308	BNG	C4'-C5'-C6'-C7'
3	A	302	BNG	O1-C1'-C2'-C3'
3	A	303	BNG	C2'-C3'-C4'-C5'
3	A	308	BNG	C3'-C4'-C5'-C6'
3	A	314	BNG	C6'-C7'-C8'-C9'
3	A	306	BNG	C6'-C7'-C8'-C9'
3	A	312	BNG	C4'-C5'-C6'-C7'
3	A	306	BNG	C4'-C5'-C6'-C7'
3	A	301	BNG	C5'-C6'-C7'-C8'
3	A	303	BNG	C3'-C4'-C5'-C6'
3	A	303	BNG	C1'-C2'-C3'-C4'
3	A	307	BNG	C2'-C1'-O1-C1
3	A	301	BNG	C3'-C4'-C5'-C6'
3	A	309	BNG	C2'-C3'-C4'-C5'
3	A	305	BNG	C1'-C2'-C3'-C4'
3	A	301	BNG	C4-C5-C6-O6
3	A	309	BNG	C1'-C2'-C3'-C4'
3	A	307	BNG	C6'-C7'-C8'-C9'
3	A	302	BNG	C2'-C1'-O1-C1
3	A	302	BNG	C1'-C2'-C3'-C4'
3	A	301	BNG	O1-C1'-C2'-C3'
3	A	301	BNG	O5-C5-C6-O6
3	A	307	BNG	C1'-C2'-C3'-C4'
3	A	309	BNG	C3'-C4'-C5'-C6'
3	A	307	BNG	C5'-C6'-C7'-C8'
3	A	301	BNG	C6'-C7'-C8'-C9'
3	A	312	BNG	C1'-C2'-C3'-C4'
3	A	302	BNG	C6'-C7'-C8'-C9'
3	A	306	BNG	C3'-C4'-C5'-C6'
3	A	307	BNG	O1-C1'-C2'-C3'
3	A	303	BNG	C5'-C6'-C7'-C8'



There are no ring outliers.



3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	314	BNG	1	0
3	A	306	BNG	1	0
3	A	307	BNG	8	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


















## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	182/200 (91%)	-0.11	4 (2%) 62 66	23, 33, 54, 88	1 (0%)
2	B	4/6 (66%)	-0.04	0 100 100	34, 35, 39, 46	0
All	All	186/206 (90%)	-0.11	4 (2%) 62 66	23, 33, 54, 88	1 (0%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	156	LEU	3.1
1	A	133	PHE	2.9
1	A	192	PHE	2.6
1	A	135	PHE	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	BNG	A	312	8/21	0.73	0.37	68,77,83,84	0

*Continued on next page...*

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	BNG	A	301	20/21	0.78	0.18	50,58,67,75	0
3	BNG	A	305	7/21	0.80	0.29	66,73,76,77	0
3	BNG	A	313	7/21	0.80	0.32	67,70,78,80	0
3	BNG	A	302	11/21	0.82	0.52	42,51,74,75	0
3	BNG	A	308	11/21	0.82	0.51	46,57,72,75	0
3	BNG	A	314	9/21	0.83	0.21	67,68,76,82	0
3	BNG	A	307	11/21	0.86	0.28	23,41,67,67	0
3	BNG	A	304	6/21	0.86	0.12	42,50,55,55	0
3	BNG	A	310	5/21	0.86	0.32	66,67,70,74	0
3	BNG	A	303	9/21	0.90	0.21	47,50,63,64	0
3	BNG	A	309	6/21	0.90	0.14	58,61,63,68	0
3	BNG	A	311	5/21	0.92	0.27	53,59,66,67	0
3	BNG	A	306	7/21	0.92	0.26	59,63,66,69	0
4	CL	A	315	1/1	0.99	0.06	37,37,37,37	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around BNG A 301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around BNG A 305:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around BNG A 302:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around BNG A 308:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)
















**Electron density around BNG A 303:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around BNG A 309:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.