



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 17, 2023 – 11:10 AM EST

PDB ID : 4W1V  
Title : Crystal structure of 7,8-diaminopelargonic acid synthase (BioA) from Mycobacterium tuberculosis, complexed with a thiazole inhibitor  
Authors : Finzel, B.C.; Dai, R.  
Deposited on : 2014-08-13  
Resolution : 2.24 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2391 (2.26-2.22)
Clashscore	141614	2539 (2.26-2.22)
Ramachandran outliers	138981	2489 (2.26-2.22)
Sidechain outliers	138945	2490 (2.26-2.22)
RSRZ outliers	127900	2353 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	429	 89% 10% ..
1	B	429	 89% 10% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	EDO	B	503	-	-	X	-
5	EDO	B	506	-	-	-	X

## 2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 6938 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adenosylmethionine-8-amino-7-oxononanoate aminotransferase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	425	Total 3222	C 2052	N 568	O 579	S 23	0	6	0
1	B	429	Total 3231	C 2056	N 567	O 586	S 22	0	6	0

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C<sub>8</sub>H<sub>10</sub>NO<sub>6</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total 15	C 8	N 1	O 5	P 1	0	0
2	B	1	Total 15	C 8	N 1	O 5	P 1	0	0

- Molecule 3 is dimethyl (2R)-5-(3-fluorophenyl)-1H-pyrrolo[1,2-c][1,3]thiazole-6,7-dicarboxylate 2-oxide (three-letter code: 3GS) (formula: C<sub>16</sub>H<sub>14</sub>FNO<sub>5</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	F	N	O			S
3	A	1	Total	C	F	N	O	S	0	0
			24	16	1	1	5	1		
3	B	1	Total	C	F	N	O	S	0	0
			24	16	1	1	5	1		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula:  $C_4H_{10}O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total Cl 1 1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	196	Total O 196 196	0	0
7	B	187	Total O 187 187	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Adenosylmethionine-8-amino-7-oxonanoate aminotransferase



- Molecule 1: Adenosylmethionine-8-amino-7-oxonanoate aminotransferase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.04Å 66.24Å 204.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	102.12 – 2.24 28.19 – 2.24	Depositor EDS
% Data completeness (in resolution range)	96.0 (102.12-2.24) 96.1 (28.19-2.24)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.72 (at 2.24Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.170 , 0.210 0.170 , 0.206	Depositor DCC
$R_{free}$ test set	2023 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.8	Xtrriage
Anisotropy	0.120	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 30.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.036 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6938	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.45% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 3GS, PEG, CL, PLP, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.61	5/3315 (0.2%)	0.65	0/4530
1	B	0.58	4/3326 (0.1%)	0.62	1/4546 (0.0%)
All	All	0.60	9/6641 (0.1%)	0.64	1/9076 (0.0%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	65	TRP	CD2-CE2	5.70	1.48	1.41
1	B	22	TRP	CD2-CE2	5.67	1.48	1.41
1	A	398	TRP	CD2-CE2	5.61	1.48	1.41
1	A	22	TRP	CD2-CE2	5.41	1.47	1.41
1	A	139	TRP	CD2-CE2	5.19	1.47	1.41
1	B	45	TRP	CD2-CE2	5.15	1.47	1.41
1	A	178	TRP	CD2-CE2	5.13	1.47	1.41
1	B	64	TRP	CD2-CE2	5.08	1.47	1.41
1	A	341	TRP	CD2-CE2	5.01	1.47	1.41

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	399	LEU	CA-CB-CG	5.43	127.80	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3222	0	3201	43	0
1	B	3231	0	3214	37	0
2	A	15	0	6	3	0
2	B	15	0	6	2	0
3	A	24	0	14	1	0
3	B	24	0	14	2	0
4	A	7	0	10	0	0
5	B	16	0	24	10	0
6	B	1	0	0	0	0
7	A	196	0	0	5	0
7	B	187	0	0	2	0
All	All	6938	0	6489	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:283:LYS:HZ2	2:B:501:PLP:C4A	1.29	1.32
1:A:283:LYS:HZ1	2:A:501:PLP:C4A	1.41	1.32
1:A:283:LYS:HZ2	2:A:501:PLP:C4A	1.69	1.01
1:B:283:LYS:HZ1	2:B:501:PLP:C4A	1.60	0.91
1:A:65[A]:TRP:HH2	1:A:407:TYR:HH	0.91	0.88
1:B:357:ASP:O	1:B:360:ARG:HG3	1.80	0.81
1:A:25:TYR:OH	1:A:160:ASP:OD2	2.01	0.77
1:A:65[A]:TRP:HB2	1:A:283:LYS:HD3	1.70	0.74
1:A:65[A]:TRP:HH2	1:A:407:TYR:OH	1.71	0.73
1:B:65:TRP:HB2	1:B:283:LYS:HD3	1.71	0.71
1:A:34:SER:N	7:A:756:HOH:O	2.24	0.69
1:A:61:MET:HA	1:A:409[A]:MET:CE	2.24	0.67
1:A:65[A]:TRP:N	1:A:65[A]:TRP:HE3	1.93	0.66
1:A:64:TRP:HA	1:A:65[A]:TRP:CE3	2.31	0.66
1:A:409[A]:MET:HE3	1:A:409[A]:MET:H	1.61	0.66
1:B:286:THR:HG22	1:B:329[A]:SER:OG	1.97	0.65
1:A:409[A]:MET:HE3	1:A:409[A]:MET:N	2.12	0.64
1:B:156:GLY:CA	5:B:503:EDO:H12	2.27	0.63
1:B:246:ARG:HD2	5:B:507:EDO:H12	1.80	0.62
1:A:74:PRO:O	1:A:78:GLN:HG2	2.00	0.61

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:219:VAL:HA	5:B:503:EDO:H22	1.82	0.60
1:B:193:ARG:NH2	1:B:369:ARG:HB3	2.16	0.59
1:A:64:TRP:HA	1:A:65[A]:TRP:HE3	1.69	0.58
1:A:176:SER:HA	1:A:178:TRP:H	1.67	0.58
1:A:65[A]:TRP:N	1:A:65[A]:TRP:CE3	2.72	0.58
1:B:156:GLY:HA3	5:B:503:EDO:H12	1.84	0.58
1:A:61:MET:HA	1:A:409[A]:MET:HE1	1.86	0.56
1:A:65[B]:TRP:N	1:A:65[B]:TRP:CD1	2.75	0.55
1:A:64:TRP:C	1:A:65[A]:TRP:HE3	2.11	0.54
1:B:400:ARG:NH1	7:B:741:HOH:O	2.41	0.54
1:B:395:ARG:CZ	1:B:428:GLU:HG3	2.38	0.54
1:A:303:HIS:HB3	7:A:755:HOH:O	2.09	0.53
1:B:389:THR:HB	1:B:390:PRO:HD3	1.92	0.51
1:A:318:THR:HG21	1:B:283:LYS:HG2	1.92	0.51
1:B:356:LEU:HD22	1:B:377:ILE:HD11	1.92	0.51
1:A:64:TRP:C	1:A:65[A]:TRP:CE3	2.85	0.50
1:B:156:GLY:HA2	5:B:503:EDO:H12	1.93	0.50
1:B:153:TRP:CD1	5:B:503:EDO:H21	2.46	0.50
1:B:153:TRP:HE1	5:B:503:EDO:HO1	1.59	0.50
1:A:9:THR:OG1	1:A:12:GLN:HG3	2.12	0.49
1:B:246:ARG:HB3	5:B:507:EDO:H12	1.95	0.49
1:B:57:VAL:HG12	1:B:396:GLY:HA2	1.95	0.48
1:A:176:SER:HA	1:A:178:TRP:N	2.27	0.48
1:B:66:THR:HB	1:B:288:GLY:HA2	1.95	0.48
1:B:395:ARG:NH1	1:B:428:GLU:HG3	2.27	0.48
1:A:23:HIS:HB3	1:A:24:PRO:HD2	1.95	0.47
1:A:409[A]:MET:CE	1:A:409[A]:MET:N	2.77	0.47
1:A:123:SER:HA	7:A:683:HOH:O	2.14	0.47
1:A:286:THR:HG22	1:A:329[A]:SER:OG	2.14	0.47
1:B:286:THR:HG22	1:B:329[A]:SER:HG	1.78	0.47
1:B:246:ARG:HH11	5:B:507:EDO:H12	1.80	0.47
1:A:65[B]:TRP:HB3	1:A:409[B]:MET:CE	2.46	0.46
1:A:65[B]:TRP:HB3	1:A:409[B]:MET:HE2	1.97	0.46
1:B:193:ARG:HH21	1:B:369:ARG:HB3	1.79	0.46
1:A:156:GLY:HA3	7:A:685:HOH:O	2.15	0.46
1:A:64:TRP:CA	1:A:65[A]:TRP:HE3	2.29	0.46
1:B:156:GLY:HA3	5:B:503:EDO:C1	2.46	0.45
1:A:50:ARG:NH1	1:A:394:ASP:OD1	2.43	0.45
1:A:306:SER:HB3	1:B:27:SER:HB2	1.98	0.45
1:B:381:ARG:HD2	7:B:782:HOH:O	2.17	0.44
1:B:244:CYS:HB3	1:B:249:VAL:O	2.17	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61:MET:HA	1:A:409[A]:MET:HE3	1.98	0.44
1:A:283:LYS:CE	2:A:501:PLP:C4A	2.89	0.44
1:A:160:ASP:HB3	1:B:315:HIS:CE1	2.53	0.43
1:A:27:SER:OG	1:A:30:ARG:HA	2.19	0.43
1:B:312:ALA:HB3	1:B:314:MET:CE	2.48	0.43
3:B:502:3GS:CAI	3:B:502:3GS:CAL	2.97	0.43
1:B:282:GLY:O	1:B:284:ALA:N	2.51	0.42
1:A:282:GLY:O	1:A:284:ALA:N	2.52	0.42
1:A:21:LEU:HD21	1:B:104:ALA:HB2	2.01	0.42
1:A:314:MET:HA	1:A:314:MET:CE	2.50	0.42
1:B:25:TYR:HH	1:B:160[A]:ASP:CG	2.23	0.42
1:A:163:LEU:C	1:A:163:LEU:HD13	2.40	0.42
1:B:25:TYR:OH	1:B:160[A]:ASP:OD2	2.38	0.41
3:B:502:3GS:CAL	3:B:502:3GS:H15	2.50	0.41
1:B:403:ARG:HB3	1:B:404:ASN:H	1.56	0.41
1:A:400:ARG:NH1	7:A:760:HOH:O	2.38	0.41
1:B:429:VAL:O	1:B:433:VAL:HG23	2.21	0.41
3:A:502:3GS:CAI	3:A:502:3GS:CAL	2.99	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	427/429 (100%)	412 (96%)	11 (3%)	4 (1%)	17	13
1	B	433/429 (101%)	420 (97%)	11 (2%)	2 (0%)	29	28
All	All	860/858 (100%)	832 (97%)	22 (3%)	6 (1%)	22	20

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	283	LYS
1	A	174	MET
1	A	179	THR
1	B	283	LYS
1	B	316	GLY
1	A	316	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	329/325 (101%)	321 (98%)	8 (2%)	49	55
1	B	331/325 (102%)	326 (98%)	5 (2%)	65	72
All	All	660/650 (102%)	647 (98%)	13 (2%)	60	62

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	65[A]	TRP
1	A	65[B]	TRP
1	A	153	TRP
1	A	177	LEU
1	A	291	SER
1	A	348	LEU
1	A	409[A]	MET
1	A	409[B]	MET
1	B	122	ASP
1	B	142	ARG
1	B	153	TRP
1	B	357	ASP
1	B	435	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	53	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 1 is monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	PLP	B	501	1	15,15,16	2.80	3 (20%)	20,22,23	1.55	3 (15%)
2	PLP	A	501	1	15,15,16	2.95	3 (20%)	20,22,23	1.64	6 (30%)
5	EDO	B	506	-	3,3,3	0.49	0	2,2,2	0.36	0
5	EDO	B	503	-	3,3,3	0.38	0	2,2,2	0.41	0
3	3GS	B	502	-	22,26,26	3.37	6 (27%)	26,38,38	1.44	5 (19%)
3	3GS	A	502	-	22,26,26	3.38	7 (31%)	26,38,38	1.43	4 (15%)
5	EDO	B	507	-	3,3,3	0.33	0	2,2,2	0.69	0
5	EDO	B	505	-	3,3,3	0.50	0	2,2,2	0.33	0
4	PEG	A	503	-	6,6,6	0.46	0	5,5,5	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	B	501	1	-	0/6/6/8	0/1/1/1
2	PLP	A	501	1	-	0/6/6/8	0/1/1/1
5	EDO	B	506	-	-	0/1/1/1	-
5	EDO	B	503	-	-	0/1/1/1	-
3	3GS	B	502	-	-	4/16/24/24	0/3/3/3
3	3GS	A	502	-	-	2/16/24/24	0/3/3/3
5	EDO	B	507	-	-	1/1/1/1	-
5	EDO	B	505	-	-	1/1/1/1	-
4	PEG	A	503	-	-	3/4/4/4	-

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	502	3GS	CAK-CAS	-10.09	1.35	1.51
3	A	502	3GS	CAK-CAS	-9.75	1.36	1.51
2	A	501	PLP	C5-C4	7.63	1.48	1.40
2	B	501	PLP	C5-C4	7.31	1.48	1.40
2	A	501	PLP	C3-C2	6.73	1.47	1.40
2	B	501	PLP	C3-C2	6.39	1.47	1.40
3	B	502	3GS	CAU-CAP	-6.28	1.35	1.50
3	A	502	3GS	CAU-CAP	-6.03	1.36	1.50
3	A	502	3GS	CAU-CAV	-5.93	1.34	1.40
3	A	502	3GS	CAR-CAV	-5.51	1.35	1.48
3	B	502	3GS	CAT-CAO	-5.36	1.37	1.50
3	B	502	3GS	CAU-CAV	-5.29	1.35	1.40
3	A	502	3GS	CAT-CAO	-5.15	1.38	1.50
3	B	502	3GS	CAR-CAV	-5.12	1.36	1.48
2	A	501	PLP	C3-C4	4.32	1.49	1.40
2	B	501	PLP	C3-C4	4.00	1.48	1.40
3	B	502	3GS	CAT-CAS	-2.73	1.34	1.40
3	A	502	3GS	CAT-CAS	-2.48	1.35	1.40
3	A	502	3GS	FAF-CAQ	-2.36	1.30	1.36

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	502	3GS	CAJ-CAR-CAV	3.66	124.10	119.59
2	A	501	PLP	O4P-C5A-C5	3.56	116.14	109.35
3	A	502	3GS	CAJ-CAR-CAV	3.36	123.72	119.59
3	B	502	3GS	OAE-SAX-CAK	-3.08	103.10	106.89
3	A	502	3GS	OAE-SAX-CAK	-3.02	103.17	106.89

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	PLP	C6-N1-C2	2.99	124.70	119.17
2	A	501	PLP	C6-N1-C2	2.88	124.51	119.17
2	B	501	PLP	C2A-C2-N1	2.60	122.75	117.67
2	B	501	PLP	O4P-C5A-C5	2.55	114.21	109.35
2	A	501	PLP	O2P-P-O4P	-2.28	100.68	106.73
2	A	501	PLP	C2A-C2-N1	2.25	122.07	117.67
3	B	502	3GS	CAG-CAI-CAR	2.16	123.28	120.56
3	A	502	3GS	OAN-CAP-CAU	2.15	117.43	112.27
3	B	502	3GS	CAS-CAT-CAO	-2.15	122.56	127.19
2	A	501	PLP	O3P-P-O2P	2.13	115.77	107.64
2	A	501	PLP	O3-C3-C4	2.08	123.58	118.10
3	B	502	3GS	OAN-CAP-CAU	2.07	117.24	112.27
3	A	502	3GS	CAS-CAT-CAO	-2.05	122.78	127.19

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	502	3GS	CAT-CAO-OAM-CAA
4	A	503	PEG	O1-C1-C2-O2
3	A	502	3GS	CAT-CAO-OAM-CAA
5	B	505	EDO	O1-C1-C2-O2
5	B	507	EDO	O1-C1-C2-O2
3	B	502	3GS	CAU-CAP-OAN-CAB
4	A	503	PEG	C4-C3-O2-C2
3	A	502	3GS	OAD-CAP-CAU-CAV
3	B	502	3GS	OAD-CAP-CAU-CAV
3	B	502	3GS	OAC-CAO-OAM-CAA
4	A	503	PEG	O2-C3-C4-O4


There are no ring outliers.


6 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	PLP	2	0
2	A	501	PLP	3	0
5	B	503	EDO	7	0
3	B	502	3GS	2	0
3	A	502	3GS	1	0
5	B	507	EDO	3	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	425/429 (99%)	-0.51	5 (1%) 79 80	11, 17, 33, 56	2 (0%)
1	B	429/429 (100%)	-0.47	5 (1%) 79 80	12, 19, 38, 54	3 (0%)
All	All	854/858 (99%)	-0.49	10 (1%) 79 80	11, 18, 37, 56	5 (0%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	177	LEU	4.2
1	B	32	ALA	2.8
1	A	51	ASP	2.7
1	B	177	LEU	2.4
1	A	176	SER	2.3
1	B	30	ARG	2.3
1	B	435	SER	2.1
1	A	180	ASP	2.1
1	B	394	ASP	2.1
1	A	65[A]	TRP	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	EDO	B	506	4/4	0.78	0.40	48,48,50,51	0
6	CL	B	504	1/1	0.80	0.07	55,55,55,55	0
5	EDO	B	505	4/4	0.84	0.17	30,31,32,32	0
4	PEG	A	503	7/7	0.87	0.15	48,52,54,54	0
5	EDO	B	507	4/4	0.91	0.11	31,33,33,34	0
3	3GS	A	502	24/24	0.92	0.14	18,21,22,25	0
3	3GS	B	502	24/24	0.94	0.12	18,21,24,24	0
2	PLP	B	501	15/16	0.97	0.10	13,13,14,15	0
5	EDO	B	503	4/4	0.97	0.24	18,19,19,19	0
2	PLP	A	501	15/16	0.98	0.10	11,12,13,13	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around 3GS A 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.