

Full wwPDB X-ray Structure Validation Report (i)

Dec 17, 2023 – 05:07 PM EST

PDB ID	:	4WQP
Title	:	Crystal structure of RORc in complex with a benzyl sulfonamide inverse ago-
		nist
Authors	:	Boenig, G.; Hymowitz, S.G.
Deposited on		
Resolution	:	1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity Mogul Xtriage (Phenix) EDS	:	4.02b-467 1.8.5 (274361), CSD as541be (2020) 1.13 2.36
buster-report Percentile statistics Refmac	: : :	1.1.7 (2018) 20191225.v01 (using entries in the PDB archive December 25th 2019) 5.8.0158 7.0.044 (Gargrove)
Ideal geometry (DNA, RNA) Validation Pipeline (wwPDB-VP)		Parkinson et al. (1996) 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	265	82%	6%	12%
1	В	265	76%	8%	17%
2	Р	13	46%		15%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 4205 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms		ZeroOcc	AltConf	Trace	
1	Δ	234	Total	С	Ν	0	S	0	4	0
	A	234	1924	1222	345	342	15	0	4	0
1	р	221	Total	С	Ν	0	S	0	7	0
	D	221	1837	1170	328	325	14	0	1	0

• Molecule 1 is a protein called Nuclear receptor ROR-gamma.

Chain	Residue	Modelled	Actual	Comment	Reference
А	246	MET	-	initiating methionine	UNP P51449
А	247	HIS	-	expression tag	UNP P51449
А	248	HIS	-	expression tag	UNP P51449
А	249	HIS	-	expression tag	UNP P51449
А	250	HIS	-	expression tag	UNP P51449
А	251	HIS	-	expression tag	UNP P51449
А	252	HIS	-	expression tag	UNP P51449
А	253	GLY	-	expression tag	UNP P51449
А	254	GLU	-	expression tag	UNP P51449
А	255	ASN	-	expression tag	UNP P51449
A	256	LEU	-	expression tag	UNP P51449
А	257	TYR	-	expression tag	UNP P51449
А	258	PHE	-	expression tag	UNP P51449
А	259	GLN	-	expression tag	UNP P51449
А	260	GLY	-	expression tag	UNP P51449
А	261	SER	-	expression tag	UNP P51449
А	508	GLY	-	expression tag	UNP P51449
А	509	ASN	-	expression tag	UNP P51449
А	510	SER	-	expression tag	UNP P51449
В	246	MET	-	initiating methionine	UNP P51449
В	247	HIS	-	expression tag	UNP P51449
В	248	HIS	-	expression tag	UNP P51449
В	249	HIS	-	expression tag	UNP P51449
В	250	HIS	-	expression tag	UNP P51449
В	251	HIS	-	expression tag	UNP P51449

There are 38 discrepancies between the modelled and reference sequences:

Continued on next page...




Chain	Residue	Modelled	Actual	Comment	Reference
В	252	HIS	-	expression tag	UNP P51449
В	253	GLY	-	expression tag	UNP P51449
В	254	GLU	-	expression tag	UNP P51449
В	255	ASN	-	expression tag	UNP P51449
В	256	LEU	-	expression tag	UNP P51449
В	257	TYR	-	expression tag	UNP P51449
В	258	PHE	-	expression tag	UNP P51449
В	259	GLN	-	expression tag	UNP P51449
В	260	GLY	-	expression tag	UNP P51449
В	261	SER	-	expression tag	UNP P51449
В	508	GLY	-	expression tag	UNP P51449
В	509	ASN	-	expression tag	UNP P51449
В	510	SER	-	expression tag	UNP P51449

Continued from previous page...

• Molecule 2 is a protein called VAL-GLU-ARG-LEU-GLN-ILE-PHE-GLN-HIS-LEU-HIS-P RO-ILE.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace		
2	Р	13	Total 121	C 80	N 23	O 18	0	1	0

• Molecule 3 is N-[4-(4-acetylpiperazin-1-yl)benzyl]-N-(2-methylpropyl)-1-phenylmethanesulf on amide (three-letter code: 3SX) (formula: $C_{24}H_{33}N_3O_3S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	А	1	Total 31	C 24	N 3	0 3	S 1	0	0


Continued on next page...





Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	D	1	Total	С	Ν	0	S	0	0
0	D	1	31	24	3	3	1	0	0



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
4	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0


• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	129	Total O 129 129	0	0
5	В	116	Total O 116 116	0	0
5	Р	6	Total O 6 6	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Nuclear receptor ROR-gamma



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 61	Depositor
Cell constants	99.73Å 99.73Å 129.61Å	Denesiten
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.97 - 1.99	Depositor
Resolution (A)	40.97 - 1.99	EDS
% Data completeness	98.9 (40.97-1.99)	Depositor
(in resolution range)	98.9(40.97 - 1.99)	EDS
R _{merge}	(Not available)	Depositor
R _{sym}	0.06	Depositor
$< I/\sigma(I) > 1$	$3.16 (at 1.98 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
D D	0.150 , 0.180	Depositor
R, R_{free}	0.149 , 0.172	DCC
R_{free} test set	2534 reflections $(5.09%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	32.3	Xtriage
Anisotropy	0.220	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.35, 31.6	EDS
L-test for twinning ²	$< L >=0.41, < L^2>=0.24$	Xtriage
Estimated twinning fraction	0.235 for h,-h-k,-l	Xtriage
Dependent of training frontion	0.743 for H, K, L	Deperitor
Reported twinning fraction	0.257 for K, H, -L	Depositor
Outliers	0 of 49779 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4205	wwPDB-VP
Average B, all atoms $(Å^2)$	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.63% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 3SX, $\mathrm{SO4}$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.36	0/1975	0.50	0/2658	
1	В	0.38	0/1894	0.52	0/2546	
2	Р	0.38	0/127	0.47	0/171	
All	All	0.37	0/3996	0.51	0/5375	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1924	0	1933	8	0
1	В	1837	0	1864	18	0
2	Р	121	0	125	3	0
3	А	31	0	33	0	0
3	В	31	0	33	1	0
4	А	5	0	0	0	0
4	В	5	0	0	0	0
5	А	129	0	0	0	0
5	В	116	0	0	1	0
5	Р	6	0	0	0	0
All	All	4205	0	3988	27	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
		distance (Å)	overlap (Å)
1:B:324[A]:LEU:HD11	1:B:358:MET:CE	1.73	1.18
1:B:324[A]:LEU:HD11	1:B:358:MET:HE3	1.35	1.03
1:B:324[A]:LEU:HD11	1:B:358:MET:HE1	1.48	0.95
1:B:362[B]:LEU:HD21	1:B:475:LEU:HD23	1.51	0.92
1:B:362[B]:LEU:HD23	1:B:472:LEU:CD1	2.20	0.71
1:B:317:TRP:CD2	1:B:486:PHE:HB3	2.25	0.71
1:B:362[B]:LEU:HD23	1:B:472:LEU:HD12	1.72	0.70
1:B:358:MET:HB3	1:B:476:CYS:SG	2.37	0.65
1:B:324[A]:LEU:CD1	1:B:358:MET:HE1	2.28	0.59
1:A:317:TRP:CD2	1:A:486:PHE:HB3	2.38	0.59
1:B:332:VAL:HG21	2:P:488:HIS:HB3	1.88	0.55
1:A:325:THR:HG23	2:P:491:PRO:HB2	1.92	0.52
1:A:484:GLN:HA	1:A:487:GLN:HE21	1.76	0.50
1:A:369:TYR:OH	1:A:405:HIS:HD2	1.94	0.50
1:B:317:TRP:CE2	1:B:486:PHE:HB3	2.46	0.50
1:B:412:PHE:HB3	1:B:417:ILE:HG13	1.93	0.50
1:B:332:VAL:HG21	2:P:488:HIS:CB	2.43	0.48
1:A:407:LEU:HD22	1:A:412:PHE:HE2	1.78	0.47
1:A:291:ASP:O	1:A:295:GLN:HG3	2.15	0.46
1:B:427:ASN:O	1:B:430:ARG:HG2	2.16	0.45
3:B:601:3SX:H4	3:B:601:3SX:H7	1.69	0.45
1:B:452:HIS:NE2	1:B:456:LYS:HD2	2.31	0.45
1:A:426:ILE:HG21	1:A:442:LEU:HG	2.01	0.43
1:B:393:CYS:O	1:B:397:ILE:HG12	2.20	0.41
1:B:426:ILE:HG21	1:B:442[A]:LEU:HG	2.02	0.41
1:A:437:ARG:HA	1:A:437:ARG:HD2	1.86	0.41
1:B:354:LYS:HG2	5:B:771:HOH:O	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	236/265~(89%)	234~(99%)	2(1%)	0	100	100
1	В	226/265~(85%)	223~(99%)	3 (1%)	0	100	100
2	Р	12/13~(92%)	12 (100%)	0	0	100	100
All	All	474/543~(87%)	469 (99%)	5 (1%)	0	100	100

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	А	212/235~(90%)	209~(99%)	3~(1%)	67 72		
1	В	204/235~(87%)	204 (100%)	0	100 100		
2	Р	14/13~(108%)	14 (100%)	0	100 100		
All	All	430/483~(89%)	427~(99%)	3 (1%)	84 88		

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	315	GLU
1	А	495	GLN
1	А	498	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	405	HIS
1	А	487	GLN
1	В	478	GLN



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain	Res	Link	Bo	ond leng	$_{\rm ths}$	B	ond ang	gles
NIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
3	3SX	В	601	-	33,33,33	1.04	2 (6%)	44,46,46	1.48	10 (22%)
4	SO4	В	602	-	4,4,4	0.42	0	6,6,6	0.16	0
4	SO4	А	602	-	4,4,4	0.40	0	6,6,6	0.22	0
3	3SX	А	601	-	33,33,33	1.10	2 (6%)	44,46,46	1.48	7 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	3SX	В	601	-	-	3/27/37/37	0/3/3/3
3	3SX	А	601	-	-	4/27/37/37	0/3/3/3

All (4) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Ζ	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
3	А	601	3SX	C25-S22	4.56	1.83	1.79
3	В	601	3SX	C25-S22	4.52	1.83	1.79
3	А	601	3SX	S22-N5	3.04	1.67	1.63
3	В	601	3SX	S22-N5	2.86	1.66	1.63

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
3	В	601	3SX	C6-N5-S22	-4.13	111.43	119.99
3	А	601	3SX	C2-C4-N5	3.92	116.96	112.41
3	А	601	3SX	O23-S22-N5	-3.30	104.43	107.23
3	В	601	3SX	C6-N5-C4	3.19	123.41	117.61
3	А	601	3SX	C21-C19-N16	3.16	121.93	118.26
3	А	601	3SX	C6-N5-S22	-3.00	113.77	119.99
3	В	601	3SX	C2-C4-N5	2.79	115.64	112.41
3	В	601	3SX	O24-S22-C25	2.56	111.37	107.81
3	В	601	3SX	C21-C19-N16	2.56	121.24	118.26
3	В	601	3SX	C11-C10-N13	-2.35	118.14	121.38
3	В	601	3SX	C17-N16-C15	2.34	117.12	112.62
3	А	601	3SX	C9-C10-N13	-2.25	118.28	121.38
3	В	601	3SX	C14-C15-N16	-2.24	105.64	110.44
3	В	601	3SX	C14-N13-C10	-2.23	112.08	118.09
3	В	601	3SX	O23-S22-N5	-2.21	105.35	107.23
3	А	601	3SX	C26-C25-S22	2.18	116.97	112.20
3	А	601	3SX	C17-N16-C15	2.06	116.59	112.62

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	А	601	3SX	C3-C2-C4-N5
3	А	601	3SX	C1-C2-C4-N5
3	В	601	3SX	C3-C2-C4-N5
3	В	601	3SX	C1-C2-C4-N5
3	А	601	3SX	C6-N5-S22-O24
3	А	601	3SX	C6-N5-S22-O23
3	В	601	3SX	C6-N5-S22-O24


There are no ring outliers.

1 monomer is involved in 1 short contact:




Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	В	601	3SX	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$ $#$ RSRZ $>$ 2		$OWAB(Å^2)$	Q<0.9
1	А	234/265~(88%)	0.30	10 (4%) 35 34	21, 35, 72, 93	0
1	В	221/265~(83%)	0.26	7 (3%) 47 46	20, 34, 57, 83	0
2	Р	13/13~(100%)	1.82	6 (46%) 0 0	42, 43, 63, 75	0
All	All	468/543~(86%)	0.32	23 (4%) 29 28	20, 35, 68, 93	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	483	LEU	5.5
1	В	485	ILE	4.8
1	А	492	ILE	4.5
1	В	317	TRP	4.0
1	А	488	HIS	3.9
1	А	489	LEU	3.7
1	А	498	PHE	3.7
1	А	494	VAL	3.6
2	Р	485	ILE	3.5
1	А	493	VAL	3.4
1	В	469	LYS	3.2
1	В	479	HIS	3.1
2	Р	480	VAL	3.1
1	А	495	GLN	2.7
1	А	485	ILE	2.7
2	Р	483	LEU	2.6
1	В	486	PHE	2.5
2	Р	481	GLU	2.5
1	А	490	HIS	2.5
1	В	294	ARG	2.3
1	А	487	GLN	2.3
2	Р	490	HIS	2.1
2	Р	492	ILE	2.0



6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B} ext{-factors}({ m \AA}^2)$	Q<0.9
4	SO4	А	602	5/5	0.90	0.29	34,48,78,94	0
3	3SX	В	601	31/31	0.93	0.16	29,40,60,65	0
3	3SX	А	601	31/31	0.94	0.16	29,39,54,70	0
4	SO4	В	602	5/5	0.94	0.34	45,50,111,114	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers (i)

There are no such residues in this entry.

