



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 8, 2020 – 09:30 PM BST

PDB ID : 5DIY
Title : Thermobaculum terrenum O-GlcNAc hydrolase mutant - D120N
Authors : Ostrowski, A.; Gundogdu, M.; Ferenbach, A.T.; Lebedev, A.; van Aalten, D.M.F.
Deposited on : 2015-09-01
Resolution : 2.06 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2684 (2.08-2.04)
Clashscore	141614	2801 (2.08-2.04)
Ramachandran outliers	138981	2768 (2.08-2.04)
Sidechain outliers	138945	2768 (2.08-2.04)
RSRZ outliers	127900	2646 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	P	7	 71% 29%
1	Q	7	 14% 71% 29%
2	A	474	 % 94% . . .
2	B	474	 % 93% . . .

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8034 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TGF-beta-activated kinase 1 and MAP3K7-binding protein 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
1	P	5	38	25	5	8	0	0	0
1	Q	5	38	25	5	8	0	0	0

- Molecule 2 is a protein called Hyaluronidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	A	463	3818	2445	635	721	17	25	0	0
2	B	462	3815	2444	637	717	17	54	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	120	ASN	ASP	conflict	UNP D1CDN2
B	120	ASN	ASP	conflict	UNP D1CDN2

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	P	1	14	8	1	5	0	0
3	Q	1	14	8	1	5	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	147	147	147	0	0
4	B	150	150	150	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: TGF-beta-activated kinase 1 and MAP3K7-binding protein 1

Chain P: 



- Molecule 1: TGF-beta-activated kinase 1 and MAP3K7-binding protein 1

Chain Q: 





- Molecule 2: Hyaluronidase

Chain A: 



- Molecule 2: Hyaluronidase

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	161.18Å 52.55Å 131.51Å 90.00° 122.35° 90.00°	Depositor
Resolution (Å)	49.03 – 2.06 49.03 – 2.06	Depositor EDS
% Data completeness (in resolution range)	99.5 (49.03-2.06) 99.5 (49.03-2.06)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.78 (at 2.07Å)	Xtrriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.201 , 0.236 0.208 , 0.241	Depositor DCC
R_{free} test set	2923 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	29.6	Xtrriage
Anisotropy	0.060	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 34.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8034	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	P	0.56	0/39	0.49	0/53
1	Q	0.53	0/39	0.70	0/53
2	A	0.51	5/3919 (0.1%)	0.65	7/5320 (0.1%)
2	B	0.44	3/3916 (0.1%)	0.66	6/5315 (0.1%)
All	All	0.48	8/7913 (0.1%)	0.66	13/10741 (0.1%)

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	58	GLU	CD-OE1	-10.74	1.13	1.25
2	A	65	LYS	CE-NZ	-9.95	1.24	1.49
2	A	66	GLU	CB-CG	-9.15	1.34	1.52
2	A	57	LYS	CD-CE	-7.25	1.33	1.51
2	B	410	GLU	CG-CD	-7.22	1.41	1.51

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	58	GLU	OE1-CD-OE2	-7.21	114.65	123.30
2	A	449	LEU	CB-CG-CD2	7.12	123.11	111.00
2	B	226	VAL	CB-CA-C	-5.88	100.22	111.40
2	A	65	LYS	CD-CE-NZ	5.82	125.09	111.70
2	A	212	LYS	CB-CG-CD	-5.64	96.94	111.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	P	38	0	36	0	0
1	Q	38	0	36	0	0
2	A	3818	0	3685	2	0
2	B	3815	0	3689	4	0
3	P	14	0	13	0	0
3	Q	14	0	13	0	0
4	A	147	0	0	0	0
4	B	150	0	0	0	0
All	All	8034	0	7472	6	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

The worst 5 of 6 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:347:LEU:HD23	2:B:390:MET:HB3	1.95	0.48
2:A:388:GLN:HG3	2:A:421:TYR:CE1	2.49	0.48
2:A:393:PHE:CE2	2:A:448:VAL:HG22	2.51	0.45
2:B:191:PHE:CD1	2:B:221:TRP:HB2	2.53	0.44
2:B:196:ASN:HD22	2:B:199:SER:HA	1.84	0.42

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	P	3/7 (43%)	3 (100%)	0	0	100	100
1	Q	3/7 (43%)	3 (100%)	0	0	100	100
2	A	459/474 (97%)	449 (98%)	8 (2%)	2 (0%)	34	25
2	B	458/474 (97%)	443 (97%)	13 (3%)	2 (0%)	34	25
All	All	923/962 (96%)	898 (97%)	21 (2%)	4 (0%)	34	25

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	170	ASN
2	B	170	ASN
2	A	226	VAL
2	B	226	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	P	5/6 (83%)	5 (100%)	0	100	100
1	Q	5/6 (83%)	5 (100%)	0	100	100
2	A	420/429 (98%)	412 (98%)	8 (2%)	57	53
2	B	419/429 (98%)	408 (97%)	11 (3%)	46	40
All	All	849/870 (98%)	830 (98%)	19 (2%)	52	46

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	44	GLN
2	B	212	LYS
2	B	409	ARG
2	A	435	ARG
2	B	435	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such

sidechains are listed below:

Mol	Chain	Res	Type
2	B	132	ASN
2	B	447	GLN
2	B	257	ASN
2	A	274	GLN
2	B	196	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	P	101	1	14,14,15	0.49	0	17,19,21	1.84	3 (17%)
3	NAG	Q	101	1	14,14,15	0.59	0	17,19,21	1.63	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	P	101	1	-	1/6/23/26	0/1/1/1
3	NAG	Q	101	1	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	P	101	NAG	C1-O5-C5	6.30	120.72	112.19
3	Q	101	NAG	C1-O5-C5	5.49	119.63	112.19
3	Q	101	NAG	O5-C1-C2	2.99	116.01	111.29
3	P	101	NAG	O5-C1-C2	2.93	115.92	111.29
3	P	101	NAG	O5-C5-C6	2.06	110.44	107.20

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	Q	101	NAG	C3-C2-N2-C7
3	P	101	NAG	C3-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	P	5/7 (71%)	1.41	0 100 100	39, 47, 51, 52	2 (40%)
1	Q	5/7 (71%)	0.81	1 (20%) 1 0	32, 36, 39, 42	2 (40%)
2	A	463/474 (97%)	-0.26	5 (1%) 80 81	20, 31, 48, 67	10 (2%)
2	B	462/474 (97%)	-0.11	6 (1%) 77 78	17, 31, 58, 71	12 (2%)
All	All	935/962 (97%)	-0.17	12 (1%) 77 78	17, 31, 54, 71	26 (2%)

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	464	TRP	3.7
2	B	133	SER	3.0
2	B	459	ALA	2.7
2	A	365	ASP	2.6
2	A	413	GLN	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NAG	P	101	14/15	0.96	0.08	26,28,29,30	0
3	NAG	Q	101	14/15	0.96	0.10	22,23,25,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.