



# Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 09:24 am BST

PDB ID : 5DX1  
Title : Crystal structure of CARM1, sinefungin, and PABP1 peptide (R455)  
Authors : Boriack-Sjodin, P.A.  
Deposited on : 2015-09-23  
Resolution : 1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4310 (1.96-1.92)
Clashscore	141614	1023 (1.94-1.94)
Ramachandran outliers	138981	1007 (1.94-1.94)
Sidechain outliers	138945	1007 (1.94-1.94)
RSRZ outliers	127900	4250 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	349	
1	B	349	
1	C	349	
1	D	349	
2	F	19	
2	G	19	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	H	19	 <p>5% 42% 58%</p>
2	I	19	 <p>11% 58% 42%</p>

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 12231 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone-arginine methyltransferase CARM1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	344	Total 2844	C 1833	N 475	O 520	S 16	0	11	0
1	B	344	Total 2843	C 1834	N 470	O 524	S 15	0	12	0
1	C	344	Total 2821	C 1818	N 471	O 515	S 17	0	9	0
1	D	344	Total 2836	C 1824	N 473	O 524	S 15	0	10	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	131	SER	-	expression tag	UNP Q86X55
A	132	ILE	-	expression tag	UNP Q86X55
A	133	ALA	-	expression tag	UNP Q86X55
B	131	SER	-	expression tag	UNP Q86X55
B	132	ILE	-	expression tag	UNP Q86X55
B	133	ALA	-	expression tag	UNP Q86X55
C	131	SER	-	expression tag	UNP Q86X55
C	132	ILE	-	expression tag	UNP Q86X55
C	133	ALA	-	expression tag	UNP Q86X55
D	131	SER	-	expression tag	UNP Q86X55
D	132	ILE	-	expression tag	UNP Q86X55
D	133	ALA	-	expression tag	UNP Q86X55

- Molecule 2 is a protein called PABP1 peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	F	9	Total 64	C 40	N 15	O 9	0	0	0
2	G	10	Total 71	C 45	N 16	O 10	0	0	0

*Continued on next page...*


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	H	8	Total	C	N	O	0	0	0
			52	33	11	8			
2	I	11	Total	C	N	O	0	0	0
			78	50	17	11			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	19	NH2	-	amidation	UNP P11940
G	19	NH2	-	amidation	UNP P11940
H	19	NH2	-	amidation	UNP P11940
I	19	NH2	-	amidation	UNP P11940

- Molecule 3 is SINEFUNGIN (three-letter code: SFG) (formula: C<sub>15</sub>H<sub>23</sub>N<sub>7</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	Total	C	N	O	0	0
			27	15	7	5		
3	B	1	Total	C	N	O	0	0
			27	15	7	5		
3	C	1	Total	C	N	O	0	0
			27	15	7	5		
3	D	1	Total	C	N	O	0	0
			27	15	7	5		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0
4	C	1	Total C O 6 3 3	0	0



- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	135	Total O 135 135	0	0
5	B	144	Total O 144 144	0	0
5	C	102	Total O 102 102	0	0
5	D	107	Total O 107 107	0	0
5	F	1	Total O 1 1	0	0
5	G	1	Total O 1 1	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Histone-arginine methyltransferase CARM1





- Molecule 1: Histone-arginine methyltransferase CARM1




- Molecule 1: Histone-arginine methyltransferase CARM1




- Molecule 1: Histone-arginine methyltransferase CARM1




- Molecule 2: PABP1 peptide







- Molecule 2: PABP1 peptide



- Molecule 2: PABP1 peptide



- Molecule 2: PABP1 peptide





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.00Å 98.32Å 208.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.99 – 1.93 45.99 – 1.93	Depositor EDS
% Data completeness (in resolution range)	99.1 (45.99-1.93) 99.1 (45.99-1.93)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.62 (at 1.94Å)	Xtrriage
Refinement program	REFMAC 5.8.0073	Depositor
R, $R_{free}$	0.196 , 0.238 0.202 , 0.243	Depositor DCC
$R_{free}$ test set	5771 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.2	Xtrriage
Anisotropy	0.088	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 41.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	12231	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 37.69 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.1281e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NMM, SFG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.53	0/2930	0.69	2/3965 (0.1%)
1	B	0.55	0/2925	0.68	0/3964
1	C	0.50	0/2907	0.66	0/3935
1	D	0.51	0/2909	0.66	0/3940
2	F	0.54	0/53	0.59	0/72
2	G	0.57	0/60	0.78	0/80
2	H	0.56	0/53	0.83	0/72
2	I	0.55	0/68	0.78	0/92
All	All	0.52	0/11905	0.68	2/16120 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	190	ASP	CB-CG-OD1	5.73	123.46	118.30
1	A	190	ASP	CB-CG-OD2	-5.02	113.78	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2844	0	2803	14	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2843	0	2800	14	0
1	C	2821	0	2780	24	0
1	D	2836	0	2781	16	0
2	F	64	0	68	1	0
2	G	71	0	74	1	0
2	H	52	0	55	0	0
2	I	78	0	82	0	0
3	A	27	0	22	0	0
3	B	27	0	22	0	0
3	C	27	0	22	0	0
3	D	27	0	22	0	0
4	A	6	0	8	0	0
4	B	12	0	16	1	0
4	C	6	0	8	0	0
5	A	135	0	0	4	0
5	B	144	0	0	3	0
5	C	102	0	0	3	0
5	D	107	0	0	2	0
5	F	1	0	0	0	0
5	G	1	0	0	0	0
All	All	12231	0	11563	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:464[B]:SER:C	1:D:465[B]:ASN:ND2	2.40	0.74
1:D:422:PHE:HA	1:D:465[B]:ASN:OD1	1.90	0.72
1:C:311[A]:ASN:ND2	1:C:327[A]:ARG:HH21	1.87	0.71
1:D:465[B]:ASN:ND2	1:D:465[B]:ASN:N	2.42	0.67
1:B:353:VAL:HG13	1:B:376:PHE:CE1	2.30	0.67
1:D:227:LYS:HD2	1:D:232:THR:HG21	1.77	0.66
1:B:309:LYS:NZ	5:B:601:HOH:O	2.29	0.66
1:A:174[B]:ARG:NH2	5:A:601:HOH:O	2.27	0.65
1:C:424:SER:HB3	5:C:658:HOH:O	1.96	0.65
1:B:383:LEU:HG	1:B:427:PHE:CE1	2.33	0.64
1:C:294:LEU:HD12	1:C:376:PHE:HZ	1.63	0.64
1:C:262:MET:HE3	1:C:418:VAL:HG11	1.80	0.63
1:B:440:LEU:HD22	1:B:450:ILE:HG12	1.82	0.62

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:336:ARG:HG2	1:C:466:LEU:O	1.99	0.62
1:D:423:GLN:N	1:D:465[B]:ASN:OD1	2.33	0.60
1:C:458:GLN:H	1:C:458:GLN:NE2	1.99	0.60
1:C:311[A]:ASN:ND2	1:C:327[A]:ARG:HD3	2.18	0.58
1:C:294:LEU:CD1	1:C:376:PHE:HZ	2.17	0.58
1:D:465[B]:ASN:HD22	1:D:465[B]:ASN:N	2.03	0.56
1:B:473:PHE:CD1	4:B:503:GOL:H2	2.41	0.56
1:B:383:LEU:HD22	1:B:425:PRO:HB2	1.90	0.54
1:A:174[A]:ARG:HG3	5:A:611:HOH:O	2.07	0.53
1:B:276:LYS:HD3	1:B:285[A]:MET:SD	2.48	0.53
1:C:148:GLN:NE2	1:D:148:GLN:HG3	2.25	0.51
1:B:422:PHE:HA	1:B:465:ASN:OD1	2.10	0.51
1:D:137:PHE:HB2	1:D:243:GLU:CG	2.42	0.50
1:B:336:ARG:HD3	5:B:736:HOH:O	2.12	0.50
1:B:180:HIS:ND1	5:B:603:HOH:O	2.35	0.49
1:B:376:PHE:O	1:B:433[B]:THR:HA	2.12	0.48
1:C:311[A]:ASN:HD21	1:C:327[A]:ARG:HD3	1.78	0.48
1:C:264:PHE:HB3	1:C:448:TYR:CE1	2.49	0.48
1:A:187:ILE:HG22	1:A:251:VAL:HG12	1.95	0.48
1:D:464[B]:SER:CA	1:D:465[B]:ASN:ND2	2.77	0.48
1:C:294:LEU:HD12	1:C:376:PHE:CZ	2.48	0.47
1:D:309:LYS:NZ	5:D:608:HOH:O	2.48	0.47
1:A:327[A]:ARG:HD2	1:C:327[A]:ARG:NH1	2.30	0.46
1:C:276:LYS:HD3	1:C:285[A]:MET:SD	2.55	0.46
1:B:254:ILE:HG13	1:B:279:LEU:HD13	1.97	0.46
1:C:363:LYS:NZ	5:C:604:HOH:O	2.49	0.46
1:D:207:ALA:HB3	1:D:210:ILE:HD11	1.98	0.46
1:C:458:GLN:H	1:C:458:GLN:HE21	1.61	0.46
1:C:433:THR:H	1:C:458:GLN:HE22	1.63	0.46
1:A:267:ARG:CZ	1:A:442:ALA:HB1	2.47	0.45
1:C:279:LEU:HD11	1:C:283:GLY:HA3	1.99	0.45
1:C:439:LEU:HB3	1:C:451[A]:SER:OG	2.17	0.45
1:A:252:ASP:OD1	1:A:280:LYS:HE2	2.16	0.45
1:A:276:LYS:HD3	1:A:285[B]:MET:SD	2.56	0.45
1:A:294:LEU:HG	1:A:389:PHE:CE2	2.51	0.44
1:D:262:MET:HB2	1:D:418:VAL:HG21	1.99	0.44
1:B:440:LEU:CD2	1:B:450:ILE:HG12	2.48	0.44
2:G:12:NMM:HD3	2:G:12:NMM:HA	1.85	0.44
1:D:258:PRO:HD2	1:D:259:MET:SD	2.58	0.44
1:A:314:TYR:HD1	1:A:327[B]:ARG:HG3	1.82	0.43
1:A:474:PHE:CE2	2:F:5:ALA:HB1	2.54	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:314:TYR:HB2	1:C:327[A]:ARG:HG3	2.01	0.42
1:D:227:LYS:HD2	1:D:232:THR:CG2	2.48	0.42
1:D:350:ALA:HA	1:D:379:LEU:HG	2.02	0.42
1:A:178:GLN:HG3	5:A:723:HOH:O	2.18	0.42
1:A:277:LYS:HE2	1:A:278:TYR:CE2	2.55	0.41
1:C:311[A]:ASN:HD21	1:C:327[A]:ARG:HH21	1.66	0.41
1:C:349[A]:MET:SD	1:C:384:VAL:HG22	2.60	0.41
1:A:262:MET:HA	5:A:710:HOH:O	2.19	0.41
1:C:436:GLY:HA3	1:C:453:VAL:O	2.21	0.41
1:A:360[A]:LEU:HA	1:A:360[A]:LEU:HD12	1.75	0.40
1:C:368:HIS:HE1	5:C:602:HOH:O	2.03	0.40
1:D:412[B]:LEU:CD2	5:D:633:HOH:O	2.69	0.40
1:B:264:PHE:HB3	1:B:448:TYR:CE1	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	353/349 (101%)	341 (97%)	12 (3%)	0	100	100
1	B	354/349 (101%)	344 (97%)	9 (2%)	1 (0%)	41	32
1	C	351/349 (101%)	339 (97%)	12 (3%)	0	100	100
1	D	352/349 (101%)	343 (97%)	8 (2%)	1 (0%)	41	32
2	F	7/19 (37%)	7 (100%)	0	0	100	100
2	G	7/19 (37%)	7 (100%)	0	0	100	100
2	H	6/19 (32%)	6 (100%)	0	0	100	100
2	I	8/19 (42%)	8 (100%)	0	0	100	100
All	All	1438/1472 (98%)	1395 (97%)	41 (3%)	2 (0%)	51	43

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	258	PRO
1	D	258	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	311/303 (103%)	307 (99%)	4 (1%)	69	62
1	B	312/303 (103%)	306 (98%)	6 (2%)	57	45
1	C	309/303 (102%)	308 (100%)	1 (0%)	92	93
1	D	310/303 (102%)	304 (98%)	6 (2%)	57	45
2	F	4/13 (31%)	4 (100%)	0	100	100
2	G	5/13 (38%)	5 (100%)	0	100	100
2	H	4/13 (31%)	4 (100%)	0	100	100
2	I	6/13 (46%)	6 (100%)	0	100	100
All	All	1261/1264 (100%)	1244 (99%)	17 (1%)	73	62

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	140[A]	ARG
1	A	140[B]	ARG
1	A	360[A]	LEU
1	A	360[B]	LEU
1	B	257	GLU
1	B	281	PRO
1	B	286	PHE
1	B	294[A]	LEU
1	B	294[B]	LEU
1	B	353	VAL
1	C	257	GLU
1	D	139	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	243	GLU
1	D	247	LEU
1	D	257	GLU
1	D	286	PHE
1	D	424	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	180	HIS
1	A	417	GLN
1	B	178	GLN
1	B	417	GLN
1	C	148	GLN
1	C	158	GLN
1	C	160	GLN
1	C	164	GLN
1	C	178	GLN
1	C	446	GLN
1	C	458	GLN
1	D	151	GLN
1	D	158	GLN
1	D	178	GLN
1	D	306	GLN
1	D	455	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NMM	I	12	2	9,11,12	0.53	0	6,12,14	0.37	0
2	NMM	F	12	2	9,11,12	0.56	0	6,12,14	1.78	2 (33%)
2	NMM	G	12	2	9,11,12	0.65	0	6,12,14	1.40	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NMM	I	12	2	-	2/9/11/13	-
2	NMM	F	12	2	-	3/9/11/13	-
2	NMM	G	12	2	-	4/9/11/13	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	12	NMM	CG-CD-NE	-3.36	102.61	112.21
2	G	12	NMM	NE-CZ-NH2	2.70	121.96	119.48
2	F	12	NMM	CD-NE-CZ	2.16	127.45	123.50

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	F	12	NMM	O-C-CA-CB
2	G	12	NMM	C-CA-CB-CG
2	G	12	NMM	N-CA-CB-CG
2	G	12	NMM	O-C-CA-CB
2	G	12	NMM	CA-CB-CG-CD
2	I	12	NMM	CA-CB-CG-CD
2	I	12	NMM	NE-CD-CG-CB
2	F	12	NMM	C-CA-CB-CG
2	F	12	NMM	CG-CD-NE-CZ

There are no ring outliers.

1 monomer is involved in 1 short contact:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	12	NMM	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GOL	B	502	-	5,5,5	0.31	0	5,5,5	0.24	0
3	SFG	B	501	-	22,29,29	0.78	0	18,42,42	1.62	4 (22%)
3	SFG	D	501	-	22,29,29	0.92	1 (4%)	18,42,42	1.59	3 (16%)
4	GOL	B	503	-	5,5,5	0.52	0	5,5,5	0.57	0
3	SFG	A	501	-	22,29,29	0.97	1 (4%)	18,42,42	1.62	4 (22%)
3	SFG	C	501	-	22,29,29	1.09	2 (9%)	18,42,42	1.59	5 (27%)
4	GOL	C	502	-	5,5,5	0.30	0	5,5,5	0.25	0
4	GOL	A	502	-	5,5,5	0.23	0	5,5,5	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	502	-	-	2/4/4/4	-
3	SFG	B	501	-	-	2/9/33/33	0/3/3/3
3	SFG	D	501	-	-	2/9/33/33	0/3/3/3
4	GOL	B	503	-	-	4/4/4/4	-
3	SFG	A	501	-	-	2/9/33/33	0/3/3/3

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SFG	C	501	-	-	3/9/33/33	0/3/3/3
4	GOL	C	502	-	-	4/4/4/4	-
4	GOL	A	502	-	-	1/4/4/4	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	501	SFG	C2-N3	2.70	1.36	1.32
3	A	501	SFG	C2-N3	2.59	1.36	1.32
3	C	501	SFG	C5-C4	2.12	1.46	1.40
3	D	501	SFG	C5-C4	2.07	1.46	1.40

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	SFG	N3-C2-N1	-4.34	121.90	128.68
3	D	501	SFG	N3-C2-N1	-4.21	122.11	128.68
3	B	501	SFG	N3-C2-N1	-4.09	122.29	128.68
3	C	501	SFG	N3-C2-N1	-3.31	123.51	128.68
3	D	501	SFG	C2-N1-C6	2.68	123.33	118.75
3	C	501	SFG	N6-C6-N1	2.65	124.08	118.57
3	C	501	SFG	O4'-C4'-C5'	-2.65	104.78	109.35
3	A	501	SFG	O4'-C4'-C5'	-2.56	104.92	109.35
3	A	501	SFG	C2-N1-C6	2.34	122.75	118.75
3	B	501	SFG	C1'-N9-C4	-2.28	122.63	126.64
3	B	501	SFG	N6-C6-N1	2.25	123.24	118.57
3	B	501	SFG	O4'-C4'-C5'	-2.23	105.50	109.35
3	A	501	SFG	C1'-N9-C4	-2.14	122.88	126.64
3	C	501	SFG	C2'-C3'-C4'	2.13	106.78	102.64
3	D	501	SFG	N6-C6-N1	2.07	122.88	118.57
3	C	501	SFG	C1'-N9-C4	-2.02	123.10	126.64

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	502	GOL	O1-C1-C2-C3
3	B	501	SFG	NE-CD-CG-CB
3	B	501	SFG	C5'-CD-CG-CB
3	D	501	SFG	NE-CD-CG-CB
3	D	501	SFG	C5'-CD-CG-CB

*Continued on next page...*

*Continued from previous page...*




Mol	Chain	Res	Type	Atoms
4	B	503	GOL	O1-C1-C2-C3
3	A	501	SFG	NE-CD-CG-CB
3	A	501	SFG	C5'-CD-CG-CB
3	C	501	SFG	NE-CD-CG-CB
3	C	501	SFG	C5'-CD-CG-CB
4	C	502	GOL	O1-C1-C2-C3
4	C	502	GOL	C1-C2-C3-O3
4	B	503	GOL	C1-C2-C3-O3
4	B	503	GOL	O1-C1-C2-O2
4	B	502	GOL	O1-C1-C2-O2
4	C	502	GOL	O1-C1-C2-O2
4	C	502	GOL	O2-C2-C3-O3
4	B	503	GOL	O2-C2-C3-O3
4	A	502	GOL	O2-C2-C3-O3
3	C	501	SFG	N-CA-CB-CG


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	503	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	344/349 (98%)	-0.07	5 (1%) 73 79	17, 27, 39, 63	0
1	B	344/349 (98%)	-0.12	3 (0%) 84 87	16, 25, 38, 71	0
1	C	344/349 (98%)	0.01	8 (2%) 60 67	22, 31, 42, 74	0
1	D	344/349 (98%)	0.07	7 (2%) 65 71	21, 32, 47, 78	0
2	F	8/19 (42%)	0.68	1 (12%) 3 5	29, 31, 56, 67	0
2	G	9/19 (47%)	0.89	2 (22%) 0 0	27, 31, 58, 63	0
2	H	8/19 (42%)	0.61	1 (12%) 3 5	32, 36, 64, 69	0
2	I	10/19 (52%)	1.26	2 (20%) 1 1	33, 42, 65, 76	0
All	All	1411/1472 (95%)	-0.00	29 (2%) 63 70	16, 29, 44, 78	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	I	14	PRO	5.3
2	H	4	GLY	5.0
1	C	477	THR	4.4
1	D	134	ARG	4.1
1	B	134	ARG	3.6
1	B	477	THR	3.3
1	C	314	TYR	3.2
2	I	4	GLY	3.1
1	D	282	SER	3.0
1	C	134	ARG	3.0
1	A	345	ILE	3.0
2	F	4	GLY	3.0
1	A	314	TYR	3.0
1	C	346	ARG	2.9
1	C	377[A]	HIS	2.8
1	C	139	GLU	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	477	THR	2.6
2	G	4	GLY	2.5
1	D	283	GLY	2.4
2	G	13	PRO	2.4
1	D	246	SER	2.3
1	D	428	ALA	2.3
1	A	377[A]	HIS	2.3
1	C	349[A]	MET	2.2
1	C	345	ILE	2.2
1	B	282	SER	2.1
1	D	233	ASP	2.1
1	D	177	LEU	2.1
1	A	346	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NMM	F	12	12/13	0.78	0.31	46,58,70,73	0
2	NMM	I	12	12/13	0.80	0.34	55,66,74,76	0
2	NMM	G	12	12/13	0.80	0.32	48,67,74,76	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	B	503	6/6	0.81	0.14	44,45,47,51	0
4	GOL	A	502	6/6	0.85	0.12	48,49,50,52	0

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	GOL	C	502	6/6	0.87	0.12	48,48,49,51	0
4	GOL	B	502	6/6	0.87	0.12	50,55,57,57	0
3	SFG	C	501	27/27	0.94	0.11	21,25,30,30	0
3	SFG	D	501	27/27	0.96	0.08	23,25,33,34	0
3	SFG	B	501	27/27	0.96	0.08	16,18,26,27	0
3	SFG	A	501	27/27	0.96	0.11	16,18,23,26	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









**Electron density around SFG D 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around SFG B 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.