



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 20, 2023 – 06:16 AM EDT

PDB ID : 5ERP
Title : Crystal structure of human Desmocollin-2 ectodomain fragment EC2-5
Authors : Harrison, O.J.; Brasch, J.; Shapiro, L.
Deposited on : 2015-11-14
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	451	 14% 87% 10% 5%
1	B	451	 5% 85% 12% 5%
2	C	4	 75% 25%
2	D	4	 50% 50%
2	E	4	 50% 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	CA	B	621	-	-	-	X
7	EDO	A	822	-	-	-	X

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 14287 atoms, of which 6991 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Desmocollin-2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	443	6802	2140	3358	575	710	19	0	0	0
1	B	438	6733	2121	3323	569	701	19	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	546	HIS	-	expression tag	UNP Q02487
A	547	HIS	-	expression tag	UNP Q02487
A	548	HIS	-	expression tag	UNP Q02487
A	549	HIS	-	expression tag	UNP Q02487
A	550	HIS	-	expression tag	UNP Q02487
A	551	HIS	-	expression tag	UNP Q02487
B	546	HIS	-	expression tag	UNP Q02487
B	547	HIS	-	expression tag	UNP Q02487
B	548	HIS	-	expression tag	UNP Q02487
B	549	HIS	-	expression tag	UNP Q02487
B	550	HIS	-	expression tag	UNP Q02487
B	551	HIS	-	expression tag	UNP Q02487

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
2	C	4	100	28	50	2	20	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
2	D	4	Total	C	H	N	O	0	0	0
			100	28	50	2	20			
2	E	4	Total	C	H	N	O	0	0	0
			100	28	50	2	20			

- Molecule 3 is alpha-D-mannopyranose (three-letter code: MAN) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			22	6	11	5		
3	A	1	Total	C	H	O	0	0
			22	6	11	5		
3	A	1	Total	C	H	O	0	0
			22	6	11	5		
3	A	1	Total	C	H	O	0	0
			22	6	11	5		
3	B	1	Total	C	H	O	0	0
			22	6	11	5		
3	B	1	Total	C	H	O	0	0
			22	6	11	5		
3	B	1	Total	C	H	O	0	0
			22	6	11	5		
3	B	1	Total	C	H	O	0	0
			22	6	11	5		

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
			Total	C	H	N			O	
4	A	1	Total	28	8	14	1	5	0	0
4	A	1	Total	28	8	14	1	5	0	0
4	B	1	Total	28	8	14	1	5	0	0

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
			Total	Ca			
5	A	10	Total	10	10	0	0
5	B	11	Total	11	11	0	0

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		
6	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	H	O	0	0
			10	2	6	2		
7	A	1	Total	C	H	O	0	0
			10	2	6	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	H	O	0	0
			10	2	6	2		
7	B	1	Total	C	H	O	0	0
			10	2	6	2		
7	B	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	59	Total	O	0	0
			59	59		
8	B	52	Total	O	0	0
			52	52		

Chain D:  50% 50%

MAG1
MAG2
BGA3
MAN4

- Molecule 2: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  50% 50%

MAG1
MAG2
BGA3
MAN4

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	146.22Å 94.17Å 126.31Å 90.00° 93.66° 90.00°	Depositor
Resolution (Å)	79.12 – 2.70 79.12 – 2.70	Depositor EDS
% Data completeness (in resolution range)	97.4 (79.12-2.70) 86.7 (79.12-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.48 (at 2.69Å)	Xtrriage
Refinement program	PHENIX dev_1810	Depositor
R, R_{free}	0.224 , 0.262 0.226 , 0.264	Depositor DCC
R_{free} test set	1951 reflections (4.23%)	wwPDB-VP
Wilson B-factor (Å ²)	67.3	Xtrriage
Anisotropy	0.402	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 55.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	14287	wwPDB-VP
Average B, all atoms (Å ²)	110.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, EDO, CA, BMA, NAG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.29	1/3505 (0.0%)	0.52	0/4778
1	B	0.25	0/3470	0.50	0/4729
All	All	0.27	1/6975 (0.0%)	0.51	0/9507

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	497	ALA	CA-CB	5.16	1.63	1.52

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3444	3358	3355	28	0
1	B	3410	3323	3321	27	1
2	C	50	50	43	3	0
2	D	50	50	43	0	0
2	E	50	50	43	2	0
3	A	44	44	39	1	0
3	B	44	44	40	2	0
4	A	28	28	26	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	14	14	13	0	0
5	A	10	0	0	0	0
5	B	11	0	0	0	0
6	A	5	0	0	0	0
6	B	5	0	0	0	0
7	A	12	18	18	0	0
7	B	8	12	11	1	0
8	A	59	0	0	0	0
8	B	52	0	0	0	0
All	All	7296	6991	6952	62	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 62 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:801:MAN:O6	3:A:801:MAN:C6	1.65	1.43
2:E:3:BMA:O2	2:E:4:MAN:O2	2.00	0.76
1:B:336:CYS:N	1:B:424:CYS:SG	2.64	0.71
1:B:192:ASP:OD1	1:B:200:LEU:N	2.29	0.65
1:A:454:MET:O	1:A:502:TYR:OH	2.13	0.63

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:228:SER:O	1:B:489:ARG:NH2[3_555]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	441/451 (98%)	416 (94%)	25 (6%)	0	100	100
1	B	434/451 (96%)	410 (94%)	24 (6%)	0	100	100
All	All	875/902 (97%)	826 (94%)	49 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	402/410 (98%)	393 (98%)	9 (2%)	52	79
1	B	398/410 (97%)	385 (97%)	13 (3%)	38	67
All	All	800/820 (98%)	778 (97%)	22 (3%)	43	73

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	272	VAL
1	B	324	THR
1	B	323	VAL
1	B	366	ARG
1	A	255	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates i

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	1	2,1	14,14,15	0.19	0	17,19,21	0.59	0
2	NAG	C	2	2	14,14,15	0.30	0	17,19,21	1.07	2 (11%)
2	BMA	C	3	2	11,11,12	0.69	0	15,15,17	0.76	0
2	MAN	C	4	2	11,11,12	0.66	0	15,15,17	0.94	1 (6%)
2	NAG	D	1	2,1	14,14,15	1.89	3 (21%)	17,19,21	1.11	2 (11%)
2	NAG	D	2	2	14,14,15	0.41	0	17,19,21	0.60	0
2	BMA	D	3	2	11,11,12	0.81	0	15,15,17	0.85	0
2	MAN	D	4	2	11,11,12	0.78	0	15,15,17	1.00	2 (13%)
2	NAG	E	1	2,1	14,14,15	0.30	0	17,19,21	0.61	0
2	NAG	E	2	2	14,14,15	0.25	0	17,19,21	0.54	0
2	BMA	E	3	2	11,11,12	1.53	3 (27%)	15,15,17	1.50	4 (26%)
2	MAN	E	4	2	11,11,12	1.03	1 (9%)	15,15,17	2.25	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	3/6/23/26	0/1/1/1
2	NAG	C	2	2	-	6/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	MAN	C	4	2	-	1/2/19/22	0/1/1/1
2	NAG	D	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	D	2	2	-	0/6/23/26	0/1/1/1
2	BMA	D	3	2	-	2/2/19/22	0/1/1/1
2	MAN	D	4	2	-	1/2/19/22	0/1/1/1
2	NAG	E	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	E	2	2	-	2/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BMA	E	3	2	-	2/2/19/22	0/1/1/1
2	MAN	E	4	2	-	2/2/19/22	0/1/1/1

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1	NAG	O5-C1	5.89	1.53	1.43
2	E	3	BMA	O3-C3	3.16	1.50	1.43
2	E	4	MAN	C1-C2	2.67	1.58	1.52
2	D	1	NAG	C1-C2	2.59	1.56	1.52
2	E	3	BMA	C1-C2	-2.15	1.47	1.52

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	4	MAN	C1-C2-C3	5.43	116.34	109.67
2	E	3	BMA	C1-O5-C5	3.99	117.59	112.19
2	E	4	MAN	C1-O5-C5	3.94	117.54	112.19
2	E	4	MAN	O2-C2-C3	-3.94	102.24	110.14
2	C	2	NAG	C2-N2-C7	3.39	127.73	122.90

There are no chirality outliers.

5 of 19 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	E	4	MAN	C4-C5-C6-O6
2	E	2	NAG	C4-C5-C6-O6
2	C	2	NAG	O5-C5-C6-O6
2	E	2	NAG	O5-C5-C6-O6
2	C	1	NAG	O5-C5-C6-O6


There are no ring outliers.

5 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	NAG	1	0
2	E	3	BMA	2	0
2	C	3	BMA	1	0
2	E	4	MAN	2	0
2	C	2	NAG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 39 ligands modelled in this entry, 21 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
7	EDO	A	822	-	3,3,3	0.44	0	2,2,2	0.44	0
6	SO4	B	625	-	4,4,4	0.14	0	6,6,6	0.06	0
3	MAN	A	801	1	11,11,12	2.08	3 (27%)	15,15,17	2.82	4 (26%)
4	NAG	A	805	1	14,14,15	1.08	2 (14%)	17,19,21	1.00	1 (5%)
4	NAG	A	810	1	14,14,15	1.70	1 (7%)	17,19,21	1.56	4 (23%)
3	MAN	A	802	1	11,11,12	0.88	1 (9%)	15,15,17	1.56	3 (20%)
3	MAN	B	612	1	11,11,12	0.92	1 (9%)	15,15,17	1.28	2 (13%)
4	NAG	B	601	1	14,14,15	0.50	0	17,19,21	0.66	1 (5%)
7	EDO	B	626	-	3,3,3	0.44	0	2,2,2	0.29	0
7	EDO	A	824	-	3,3,3	0.45	0	2,2,2	0.31	0
3	MAN	B	610	1	11,11,12	0.68	0	15,15,17	1.28	2 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MAN	A	803	1	11,11,12	0.94	1 (9%)	15,15,17	1.09	1 (6%)
7	EDO	A	823	-	3,3,3	0.44	0	2,2,2	0.33	0
7	EDO	B	627	-	3,3,3	0.69	0	2,2,2	2.10	1 (50%)
3	MAN	B	611	1	11,11,12	0.69	0	15,15,17	1.33	3 (20%)
3	MAN	B	613	1	11,11,12	0.72	0	15,15,17	1.99	4 (26%)
6	SO4	A	821	-	4,4,4	0.14	0	6,6,6	0.05	0
3	MAN	A	804	1	11,11,12	0.93	1 (9%)	15,15,17	1.01	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	EDO	A	822	-	-	1/1/1/1	-
3	MAN	A	801	1	-	0/2/19/22	0/1/1/1
4	NAG	A	805	1	-	1/6/23/26	0/1/1/1
4	NAG	A	810	1	-	5/6/23/26	0/1/1/1
3	MAN	A	802	1	-	0/2/19/22	1/1/1/1
3	MAN	B	612	1	-	1/2/19/22	0/1/1/1
4	NAG	B	601	1	-	2/6/23/26	0/1/1/1
7	EDO	B	626	-	-	1/1/1/1	-
7	EDO	A	824	-	-	0/1/1/1	-
3	MAN	B	610	1	-	0/2/19/22	0/1/1/1
3	MAN	A	803	1	-	0/2/19/22	0/1/1/1
7	EDO	A	823	-	-	0/1/1/1	-
7	EDO	B	627	-	-	0/1/1/1	-
3	MAN	B	611	1	-	0/2/19/22	1/1/1/1
3	MAN	B	613	1	-	0/2/19/22	0/1/1/1
3	MAN	A	804	1	-	1/2/19/22	0/1/1/1

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	810	NAG	O5-C1	6.13	1.53	1.43
3	A	801	MAN	O6-C6	5.57	1.65	1.42
4	A	805	NAG	O5-C1	3.29	1.49	1.43
3	A	801	MAN	O5-C1	-2.48	1.39	1.43
3	B	612	MAN	O5-C5	2.43	1.48	1.43

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	801	MAN	C1-O5-C5	9.37	124.89	112.19
3	B	613	MAN	C1-O5-C5	4.64	118.48	112.19
3	A	802	MAN	C1-O5-C5	4.53	118.33	112.19
3	B	613	MAN	O5-C1-C2	4.37	117.52	110.77
3	A	801	MAN	O5-C1-C2	3.75	116.56	110.77

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	601	NAG	O5-C5-C6-O6
4	A	810	NAG	C8-C7-N2-C2
4	A	810	NAG	O7-C7-N2-C2
4	B	601	NAG	C4-C5-C6-O6
4	A	810	NAG	O5-C5-C6-O6

All (2) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	802	MAN	C1-C2-C3-C4-C5-O5
3	B	611	MAN	C1-C2-C3-C4-C5-O5

5 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	801	MAN	1	0
4	A	810	NAG	1	0
3	B	610	MAN	1	0
7	B	627	EDO	1	0
3	B	611	MAN	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	443/451 (98%)	0.90	64 (14%) 2 1	63, 89, 199, 224	0
1	B	438/451 (97%)	0.54	21 (4%) 30 28	61, 84, 124, 165	0
All	All	881/902 (97%)	0.72	85 (9%) 8 6	61, 86, 187, 224	0

The worst 5 of 85 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	511	TYR	12.6
1	A	531	VAL	9.7
1	A	529	LEU	9.6
1	A	513	VAL	9.1
1	A	533	LEU	8.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BMA	C	3	11/12	0.73	0.17	177,195,231,234	0
2	MAN	E	4	11/12	0.79	0.15	164,186,218,222	0
2	MAN	D	4	11/12	0.81	0.14	150,172,202,207	0
2	MAN	C	4	11/12	0.82	0.17	181,206,242,247	0
2	BMA	E	3	11/12	0.84	0.12	158,185,213,222	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	BMA	D	3	11/12	0.87	0.10	145,170,196,200	0
2	NAG	C	2	14/15	0.88	0.25	138,166,193,202	0
2	NAG	D	1	14/15	0.89	0.15	82,117,141,157	0
2	NAG	D	2	14/15	0.89	0.15	96,124,161,193	0
2	NAG	C	1	14/15	0.91	0.19	94,124,152,159	0
2	NAG	E	2	14/15	0.92	0.15	93,130,163,184	0
2	NAG	E	1	14/15	0.93	0.18	72,100,134,151	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain E:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	CA	B	621	1/1	0.49	0.49	217,217,217,217	0
7	EDO	A	824	4/4	0.51	0.34	98,118,122,133	0
7	EDO	A	823	4/4	0.62	0.21	135,162,168,168	0
7	EDO	A	822	4/4	0.73	0.72	106,128,132,132	0
7	EDO	B	626	4/4	0.80	0.23	92,110,126,126	0
5	CA	A	817	1/1	0.81	0.31	119,119,119,119	0
4	NAG	A	805	14/15	0.81	0.16	121,149,181,193	0
3	MAN	A	801	11/12	0.83	0.22	114,136,159,176	0
4	NAG	A	810	14/15	0.83	0.10	164,197,226,234	0
6	SO4	B	625	5/5	0.84	0.24	139,141,148,165	0
3	MAN	B	611	11/12	0.85	0.19	124,149,179,185	0
5	CA	B	614	1/1	0.86	0.21	103,103,103,103	0
3	MAN	A	804	11/12	0.88	0.25	113,147,171,194	0
4	NAG	B	601	14/15	0.88	0.14	99,123,145,155	0
3	MAN	B	613	11/12	0.89	0.27	124,149,178,179	0
3	MAN	B	612	11/12	0.89	0.33	116,143,183,187	0
7	EDO	B	627	4/4	0.89	0.33	97,132,142,158	0
3	MAN	A	802	11/12	0.90	0.21	108,133,157,166	0
5	CA	A	820	1/1	0.91	0.24	123,123,123,123	0
5	CA	B	624	1/1	0.91	0.11	119,119,119,119	0
6	SO4	A	821	5/5	0.91	0.08	150,154,165,171	0
3	MAN	A	803	11/12	0.92	0.20	94,119,141,144	0
5	CA	B	623	1/1	0.92	0.36	102,102,102,102	0
3	MAN	B	610	11/12	0.94	0.17	117,146,169,179	0
5	CA	B	615	1/1	0.95	0.27	81,81,81,81	0
5	CA	B	616	1/1	0.95	0.22	91,91,91,91	0
5	CA	B	619	1/1	0.95	0.29	77,77,77,77	0
5	CA	B	620	1/1	0.95	0.25	92,92,92,92	0
5	CA	A	811	1/1	0.95	0.30	116,116,116,116	0
5	CA	A	813	1/1	0.95	0.28	76,76,76,76	0
5	CA	A	814	1/1	0.95	0.29	83,83,83,83	0
5	CA	A	819	1/1	0.96	0.25	86,86,86,86	0
5	CA	A	818	1/1	0.96	0.26	110,110,110,110	0
5	CA	B	622	1/1	0.96	0.30	76,76,76,76	0
5	CA	A	816	1/1	0.98	0.25	74,74,74,74	0
5	CA	A	815	1/1	0.99	0.27	72,72,72,72	0
5	CA	B	617	1/1	0.99	0.23	75,75,75,75	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	CA	B	618	1/1	0.99	0.27	72,72,72,72	0
5	CA	A	812	1/1	0.99	0.28	81,81,81,81	0

6.5 Other polymers [i](#)

There are no such residues in this entry.