

wwPDB X-ray Structure Validation Summary Report (i)

Sep 20, 2023 – 08:02 AM EDT

PDB ID : 5FEM

Title : Saccharomyces cerevisiae Acetohydroxyacid Synthase in complex with bensul-

furon methyl

Authors: Guddat, L.W.; Lonhienne, T.

Deposited on : 2015-12-17

Resolution : 2.17 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.17 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	677	85%	•	12%
1	В	677	84%		12%



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 9937 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Acetolactate synthase catalytic subunit, mitochondrial.

\mathbf{Mol}	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	599	Total C N O S 4584 2901 795 866 22		0	4	0			
1	В	599	Total 4578	C 2898	N 794	O 865	S 21	0	2	0

There are 94 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	11	MET	-	expression tag	UNP P07342
A	12	HIS	-	expression tag	UNP P07342
A	13	HIS	-	expression tag	UNP P07342
A	14	HIS	-	expression tag	UNP P07342
A	15	HIS	-	expression tag	UNP P07342
A	16	HIS	-	expression tag	UNP P07342
A	17	HIS	-	expression tag	UNP P07342
A	18	SER	-	expression tag	UNP P07342
A	19	SER	-	expression tag	UNP P07342
A	20	GLY	-	expression tag	UNP P07342
A	21	LEU	-	expression tag	UNP P07342
A	22	VAL	-	expression tag	UNP P07342
A	23	PRO	-	expression tag	UNP P07342
A	24	ARG	-	expression tag	UNP P07342
A	25	GLY	-	expression tag	UNP P07342
A	26	SER	-	expression tag	UNP P07342
A	27	GLY	-	expression tag	UNP P07342
A	28	MET	-	expression tag	UNP P07342
A	29	LYS	-	expression tag	UNP P07342
A	30	GLU	-	expression tag	UNP P07342
A	31	THR	-	expression tag	UNP P07342
A	32	ALA	-	expression tag	UNP P07342
A	33	ALA	-	expression tag	UNP P07342
A	34	ALA	-	expression tag	UNP P07342
A	35	LYS	-	expression tag	UNP P07342



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
A	36	PHE	-	expression tag	UNP P07342
A	37	GLU	_	expression tag	UNP P07342
A	38	ARG	_	expression tag	UNP P07342
A	39	GLN	_	expression tag	UNP P07342
A	40	HIS	_	expression tag	UNP P07342
A	41	MET	-	expression tag	UNP P07342
A	42	ASP	_	expression tag	UNP P07342
A	43	SER	-	expression tag	UNP P07342
A	44	PRO	_	expression tag	UNP P07342
A	45	ASP	-	expression tag	UNP P07342
A	46	LEU	_	expression tag	UNP P07342
A	47	GLY	_	expression tag	UNP P07342
A	48	THR	-	expression tag	UNP P07342
A	49	ASP	_	expression tag	UNP P07342
A	50	ASP	_	expression tag	UNP P07342
A	51	ASP	_	expression tag	UNP P07342
A	52	ASP	-	expression tag	UNP P07342
A	53	LYS	_	expression tag	UNP P07342
A	54	ALA	_	expression tag	UNP P07342
A	55	MET	-	expression tag	UNP P07342
A	56	GLY	_	expression tag	UNP P07342
A	57	SER	-	expression tag	UNP P07342
В	11	MET	-	expression tag	UNP P07342
В	12	HIS	-	expression tag	UNP P07342
В	13	HIS	-	expression tag	UNP P07342
В	14	HIS	-	expression tag	UNP P07342
В	15	HIS	-	expression tag	UNP P07342
В	16	HIS	-	expression tag	UNP P07342
В	17	HIS	-	expression tag	UNP P07342
В	18	SER	-	expression tag	UNP P07342
В	19	SER	-	expression tag	UNP P07342
В	20	GLY	-	expression tag	UNP P07342
В	21	LEU	-	expression tag	UNP P07342
В	22	VAL	-	expression tag	UNP P07342
В	23	PRO	_	expression tag	UNP P07342
В	24	ARG	-	expression tag	UNP P07342
В	25	GLY	-	expression tag	UNP P07342
В	26	SER	-	expression tag	UNP P07342
В	27	GLY	-	expression tag	UNP P07342
В	28	MET	-	expression tag	UNP P07342
В	29	LYS	-	expression tag	UNP P07342
В	30	GLU	-	expression tag	UNP P07342



Continued from previous page...


Chain	Residue	Modelled Modelled	Actual	Comment	Reference
В	31	THR	_	expression tag	UNP P07342
В	32	ALA	_	expression tag	UNP P07342
В	33	ALA	_	expression tag	UNP P07342
В	34	ALA	-	expression tag	UNP P07342
В	35	LYS	-	expression tag	UNP P07342
В	36	PHE	-	expression tag	UNP P07342
В	37	GLU	-	expression tag	UNP P07342
В	38	ARG	-	expression tag	UNP P07342
В	39	GLN	-	expression tag	UNP P07342
В	40	HIS	-	expression tag	UNP P07342
В	41	MET	-	expression tag	UNP P07342
В	42	ASP	-	expression tag	UNP P07342
В	43	SER	-	expression tag	UNP P07342
В	44	PRO	-	expression tag	UNP P07342
В	45	ASP	-	expression tag	UNP P07342
В	46	LEU	-	expression tag	UNP P07342
В	47	GLY	-	expression tag	UNP P07342
В	48	THR	-	expression tag	UNP P07342
В	49	ASP	-	expression tag	UNP P07342
В	50	ASP	-	expression tag	UNP P07342
В	51	ASP	-	expression tag	UNP P07342
В	52	ASP	-	expression tag	UNP P07342
В	53	LYS	-	expression tag	UNP P07342
В	54	ALA	-	expression tag	UNP P07342
В	55	MET	-	expression tag	UNP P07342
В	56	GLY	-	expression tag	UNP P07342
В	57	SER	_	expression tag	UNP P07342

 \bullet Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

\mathbf{N}	lol	Chain	Residues	Atoms	ZeroOcc	AltConf
	2	A	2	Total Mg 2 2	0	0
	2	В	2	Total Mg 2 2	0	0


• Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
2	Λ	1	Total	С	N	О	Р	0	0	
3	A	1	53	27	9	15	2	U		
9	D	1	Total	С	N	О	Р	0	0	
3	Б	1	53	27	9	15	2	U	U	

• Molecule 4 is methyl 2-[(4,6-dimethoxypyrimidin-2-yl)carbamoylsulfamoylmethyl]benzoate (three-letter code: 60G) (formula: $C_{16}H_{18}N_4O_7S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 28	C 16	N 4	O 7	S 1	0	0



 $Continued\ from\ previous\ page...$

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
4	D	1	Total	С	N	О	S	0	0
4	Б	1	28	16	4	7	1	U	

 $\bullet \ \ \mathrm{Molecule} \ 5 \ \mathrm{is} \ \mathrm{THIAMINE} \ \mathrm{DIPHOSPHATE} \ (\mathrm{three-letter} \ \mathrm{code} \colon \ \mathrm{TPP}) \ (\mathrm{formula:} \ \ \mathrm{C}_{12}\mathrm{H}_{19}\mathrm{N}_4\mathrm{O}_7\mathrm{P}_2\mathrm{S}).$



Mol	Chain	Residues		A	tom	ıs			ZeroOcc	AltConf	
5	Λ	1	Total	С	N	О	Р	S	0	0	
5	A	1	26	12	4	7	2	1	U	U	
5	D	1	Total	С	N	О	Р	S	0	0	
	Б	1	26	12	4	7	2	1		U	

• Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	329	Total O 329 329	0	0
6	В	228	Total O 228 228	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Acetolactate synthase catalytic subunit, mitochondrial





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 4 2 2	Depositor
Cell constants	154.67Å 154.67Å 178.38Å	Donogitor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.91 - 2.17	Depositor
Resolution (A)	48.91 - 2.17	EDS
% Data completeness	99.7 (48.91-2.17)	Depositor
(in resolution range)	99.7 (48.91-2.17)	EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.56 (at 2.16Å)	Xtriage
Refinement program	PHENIX 1.8.1_1168	Depositor
D D.	0.150 , 0.173	Depositor
R, R_{free}	0.152 , 0.174	DCC
R_{free} test set	2000 reflections (1.75%)	wwPDB-VP
Wilson B-factor (Å ²)	24.7	Xtriage
Anisotropy	0.160	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.40,60.0	EDS
L-test for twinning ²	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9937	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.67% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, 60G, TPP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Во	ond angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.32	0/4688	0.50	1/6358~(0.0%)
1	В	0.29	0/4679	0.47	$2/6343 \ (0.0\%)$
All	All	0.30	0/9367	0.48	3/12701 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\mathbf{Ideal}(^{o})$
1	A	557	LEU	CA-CB-CG	6.36	129.94	115.30
1	В	557	LEU	CA-CB-CG	6.04	129.20	115.30
1	В	135	LEU	CA-CB-CG	5.16	127.16	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4584	0	4593	13	0
1	В	4578	0	4598	16	0
2	A	2	0	0	0	0
2	В	2	0	0	0	0
3	A	53	0	30	1	0
3	В	53	0	30	1	0



I 'omtamalod	trom	mmonia	maaa
Continued	11 0116	DICUIUUS	Daue
	.,	10	1

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	28	0	0	0	0
4	В	28	0	0	0	0
5	A	26	0	16	3	0
5	В	26	0	16	3	0
6	A	329	0	0	0	0
6	В	228	0	0	1	0
All	All	9937	0	9283	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 35 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:B:681:LYS:NZ	6:B:801:HOH:O	2.27	0.68
1:B:580:GLN:HB3	5:B:705:TPP:H61	1.81	0.63
1:A:379:ASP:OD1	1:A:380:ARG:NE	2.30	0.62
1:B:619:VAL:HG23	1:B:641:LEU:HD11	1.80	0.62
5:B:705:TPP:H2	5:B:705:TPP:HN42	1.65	0.61

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percent	tiles
1	A	599/677 (88%)	592 (99%)	7 (1%)	0	100	100
1	В	597/677 (88%)	590 (99%)	7 (1%)	0	100	100
All	All	1196/1354 (88%)	1182 (99%)	14 (1%)	0	100	100

There are no Ramachandran outliers to report.



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	493/556 (89%)	486 (99%)	7 (1%)	67	72	
1	В	493/556 (89%)	485 (98%)	8 (2%)	62	67	
All	All	986/1112 (89%)	971 (98%)	15 (2%)	67	69	

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	106	ASN
1	В	557	LEU
1	В	135	LEU
1	В	560	LEU
1	В	411	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.



In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Type Chain		Res	Link	В	Bond lengths			Bond angles		
MIOI	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
5	TPP	A	705	2	22,27,27	1.61	2 (9%)	29,40,40	2.01	9 (31%)	
4	60G	A	703	-	28,29,29	1.30	5 (17%)	38,40,40	3.13	16 (42%)	
4	60G	В	703	-	28,29,29	1.38	5 (17%)	38,40,40	3.13	15 (39%)	
3	FAD	В	701	-	53,58,58	2.13	12 (22%)	68,89,89	1.30	10 (14%)	
3	FAD	A	702	-	53,58,58	2.09	12 (22%)	68,89,89	1.36	11 (16%)	
5	TPP	В	705	2	22,27,27	1.71	3 (13%)	29,40,40	2.06	11 (37%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	TPP	A	705	2	-	1/16/17/17	0/2/2/2
4	60G	A	703	ı	-	3/24/24/24	0/2/2/2
4	60G	В	703	-	-	2/24/24/24	0/2/2/2
3	FAD	В	701	-	-	4/30/50/50	0/6/6/6
3	FAD	A	702	-	-	4/30/50/50	0/6/6/6
5	TPP	В	705	2	-	3/16/17/17	0/2/2/2

The worst 5 of 39 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(A)
3	В	701	FAD	O2'-C2'	-9.60	1.23	1.43
3	A	702	FAD	O2'-C2'	-9.30	1.23	1.43
5	A	705	TPP	C4-N3	-6.00	1.34	1.39
5	В	705	TPP	C4-N3	-5.72	1.34	1.39
3	В	701	FAD	C10-N1	5.01	1.43	1.33

The worst 5 of 72 bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
4	В	703	60G	OAG-SBB-OAF	-11.06	103.32	119.35
4	A	703	60G	OAG-SBB-OAF	-11.04	103.35	119.35
4	В	703	60G	C6-C5-C4	6.05	120.71	115.21
4	A	703	60G	C2-N3-C4	5.91	121.94	114.99
4	В	703	60G	C2-N3-C4	5.67	121.65	114.99

There are no chirality outliers.

5 of 17 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	В	705	TPP	PA-O3A-PB-O2B
4	В	703	60G	OAE-CAV-OAR-CAA
5	В	705	TPP	PA-O3A-PB-O1B
4	A	703	60G	CAW-CAM-SBB-OAF
4	В	703	60G	CAW-CAM-SBB-OAF


There are no ring outliers.


4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	705	TPP	3	0
3	В	701	FAD	1	0
3	A	702	FAD	1	0
5	В	705	TPP	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





















5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q<0.9
1	A	599/677 (88%)	-0.22	12 (2%) 65 72	15, 25, 46, 96	0
1	В	599/677 (88%)	0.00	27 (4%) 33 42	18, 34, 62, 104	0
All	All	1198/1354 (88%)	-0.11	39 (3%) 46 55	15, 29, 57, 104	0

The worst 5 of 39 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	83	PRO	7.6
1	A	83	PRO	6.9
1	В	280	ALA	6.0
1	A	277	THR	5.8
1	В	279	ARG	5.8

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
4	60G	В	703	28/28	0.96	0.09	30,36,42,44	0
4	60G	A	703	28/28	0.97	0.11	21,26,32,34	0
3	FAD	В	701	53/53	0.97	0.09	21,28,36,41	0
5	TPP	A	705	26/26	0.97	0.13	16,20,36,50	0
5	TPP	В	705	26/26	0.97	0.12	17,23,48,54	0
2	MG	В	702	1/1	0.98	0.07	19,19,19,19	0
3	FAD	A	702	53/53	0.99	0.11	13,18,22,23	0
2	MG	A	701	1/1	0.99	0.10	17,17,17,17	0
2	MG	В	704	1/1	0.99	0.13	19,19,19,19	0
2	MG	A	704	1/1	1.00	0.19	12,12,12,12	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around 60G A 703:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around FAD B 701:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around TPP A 705: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive) Electron density around TPP B 705: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray mF_o -DF_c (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

