



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 31, 2023 – 04:32 PM JST

PDB ID : 5GQW
Title : Crystal structure of branching enzyme W610N mutant from *Cyanothece* sp. ATCC 51142
Authors : Suzuki, R.; Suzuki, E.
Deposited on : 2016-08-08
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	793	 % 88% 8% 5%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7169 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 1,4-alpha-glucan branching enzyme GlgB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	755	6272	4053	1042	1153	24	0	1	0

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP B1WPM8
A	-18	GLY	-	expression tag	UNP B1WPM8
A	-17	SER	-	expression tag	UNP B1WPM8
A	-16	SER	-	expression tag	UNP B1WPM8
A	-15	HIS	-	expression tag	UNP B1WPM8
A	-14	HIS	-	expression tag	UNP B1WPM8
A	-13	HIS	-	expression tag	UNP B1WPM8
A	-12	HIS	-	expression tag	UNP B1WPM8
A	-11	HIS	-	expression tag	UNP B1WPM8
A	-10	HIS	-	expression tag	UNP B1WPM8
A	-9	SER	-	expression tag	UNP B1WPM8
A	-8	SER	-	expression tag	UNP B1WPM8
A	-7	GLY	-	expression tag	UNP B1WPM8
A	-6	LEU	-	expression tag	UNP B1WPM8
A	-5	VAL	-	expression tag	UNP B1WPM8
A	-4	PRO	-	expression tag	UNP B1WPM8
A	-3	ARG	-	expression tag	UNP B1WPM8
A	-2	GLY	-	expression tag	UNP B1WPM8
A	-1	SER	-	expression tag	UNP B1WPM8
A	0	HIS	-	expression tag	UNP B1WPM8
A	610	ASN	TRP	engineered mutation	UNP B1WPM8

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	4	Total Mg 4 4	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	863	Total O 863 863	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 1,4-alpha-glucan branching enzyme GlgB



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	133.62Å 133.62Å 185.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.24 – 1.80 38.06 – 1.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.24-1.80) 100.0 (38.06-1.80)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.02 (at 1.81Å)	Xtrriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.149 , 0.171 0.163 , 0.182	Depositor DCC
R_{free} test set	7758 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	19.1	Xtrriage
Anisotropy	0.004	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 45.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7169	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.33	30/6493 (0.5%)	1.19	31/8835 (0.4%)

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	58	GLU	CD-OE2	9.90	1.36	1.25
1	A	619	GLN	CG-CD	7.59	1.68	1.51
1	A	412	ARG	CZ-NH2	7.41	1.42	1.33
1	A	135	GLU	CD-OE2	7.39	1.33	1.25
1	A	654	GLU	CD-OE1	-7.20	1.17	1.25
1	A	285	TRP	CB-CG	-6.74	1.38	1.50
1	A	412	ARG	CZ-NH1	6.25	1.41	1.33
1	A	343	ARG	CZ-NH2	-6.17	1.25	1.33
1	A	605	SER	CB-OG	-6.17	1.34	1.42
1	A	531	TYR	CE1-CZ	-5.96	1.30	1.38
1	A	480	PRO	C-N	-5.91	1.22	1.33
1	A	621	GLU	CD-OE1	-5.83	1.19	1.25
1	A	8	ASP	CB-CG	5.82	1.64	1.51
1	A	489	SER	CB-OG	5.78	1.49	1.42
1	A	85	TYR	CB-CG	-5.70	1.43	1.51
1	A	572	GLU	CD-OE1	-5.69	1.19	1.25
1	A	703	TYR	CG-CD1	5.67	1.46	1.39
1	A	654	GLU	CD-OE2	-5.62	1.19	1.25
1	A	135	GLU	CG-CD	5.52	1.60	1.51
1	A	553	SER	CB-OG	5.48	1.49	1.42
1	A	91	GLU	CD-OE1	-5.35	1.19	1.25
1	A	69	HIS	N-CA	-5.31	1.35	1.46
1	A	187	SER	CB-OG	-5.31	1.35	1.42
1	A	432	ARG	CG-CD	-5.30	1.38	1.51
1	A	299	GLU	CD-OE2	5.26	1.31	1.25
1	A	406	TYR	CE1-CZ	-5.25	1.31	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	58	GLU	CG-CD	5.21	1.59	1.51
1	A	203	TYR	CE1-CZ	5.20	1.45	1.38
1	A	460	GLU	CG-CD	5.18	1.59	1.51
1	A	206	SER	CB-OG	5.15	1.49	1.42

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	525	ASP	CB-CG-OD1	11.06	128.25	118.30
1	A	525	ASP	CB-CG-OD2	-9.57	109.68	118.30
1	A	381	ASP	CB-CG-OD1	7.85	125.36	118.30
1	A	412	ARG	NE-CZ-NH1	-7.79	116.40	120.30
1	A	241	ARG	NE-CZ-NH2	-7.37	116.61	120.30
1	A	207	ASP	CB-CG-OD2	-7.25	111.77	118.30
1	A	610	ASN	CB-CA-C	-7.01	96.39	110.40
1	A	612	ASP	CB-CG-OD1	-6.69	112.28	118.30
1	A	658	CYS	CB-CA-C	6.59	123.59	110.40
1	A	575	LYS	CD-CE-NZ	-6.25	97.33	111.70
1	A	671	ARG	NE-CZ-NH1	6.22	123.41	120.30
1	A	725	LYS	CD-CE-NZ	6.00	125.50	111.70
1	A	632	ASP	CB-CG-OD1	5.84	123.56	118.30
1	A	517	ASP	CB-CG-OD2	-5.79	113.09	118.30
1	A	671	ARG	NE-CZ-NH2	-5.78	117.41	120.30
1	A	571	ASP	CB-CG-OD1	5.71	123.44	118.30
1	A	290	ARG	NE-CZ-NH2	5.65	123.12	120.30
1	A	225	ASP	CB-CG-OD1	5.54	123.29	118.30
1	A	345	GLU	OE1-CD-OE2	-5.54	116.65	123.30
1	A	58	GLU	CA-CB-CG	5.47	125.43	113.40
1	A	173	ARG	CA-CB-CG	-5.37	101.59	113.40
1	A	387	GLU	OE1-CD-OE2	-5.34	116.89	123.30
1	A	640	GLU	OE1-CD-OE2	5.29	129.65	123.30
1	A	386	TYR	CB-CG-CD2	5.23	124.14	121.00
1	A	285	TRP	CA-CB-CG	5.18	123.55	113.70
1	A	225	ASP	CB-CG-OD2	-5.18	113.64	118.30
1	A	674	ASP	CB-CG-OD1	5.14	122.92	118.30
1	A	487	GLU	OE1-CD-OE2	5.07	129.39	123.30
1	A	55	ARG	NE-CZ-NH2	-5.06	117.77	120.30
1	A	582	LEU	CB-CG-CD1	5.03	119.55	111.00
1	A	580	ARG	NE-CZ-NH1	5.00	122.80	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6272	0	5857	2	0
2	A	30	0	40	1	0
3	A	4	0	0	0	0
4	A	863	0	0	1	0
All	All	7169	0	5897	3	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (3) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:46:PRO:O	1:A:47:LYS:HB2	2.10	0.51
1:A:302:ILE:HB	1:A:303:PRO:HD3	2.00	0.43
2:A:803:GOL:H32	4:A:1361:HOH:O	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	754/793 (95%)	735 (98%)	19 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	669/700 (96%)	663 (99%)	6 (1%)	78 75

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	216	ARG
1	A	369	HIS
1	A	413	ASN
1	A	434	ASP
1	A	447	LYS
1	A	491	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	A	16	ASN
1	A	162	ASN
1	A	634	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	803	-	5,5,5	0.47	0	5,5,5	0.95	0
2	GOL	A	804	-	5,5,5	0.96	0	5,5,5	1.67	1 (20%)
2	GOL	A	802	-	5,5,5	0.74	0	5,5,5	0.57	0
2	GOL	A	801	-	5,5,5	1.84	1 (20%)	5,5,5	1.27	0
2	GOL	A	805	-	5,5,5	0.35	0	5,5,5	1.72	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	803	-	-	4/4/4/4	-
2	GOL	A	804	-	-	1/4/4/4	-
2	GOL	A	802	-	-	0/4/4/4	-
2	GOL	A	801	-	-	0/4/4/4	-
2	GOL	A	805	-	-	0/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	GOL	O2-C2	3.39	1.53	1.43

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	804	GOL	C3-C2-C1	2.94	123.14	111.70
2	A	805	GOL	O2-C2-C3	2.54	120.31	109.12

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	803	GOL	O1-C1-C2-C3
2	A	803	GOL	C1-C2-C3-O3
2	A	803	GOL	O2-C2-C3-O3
2	A	804	GOL	C1-C2-C3-O3
2	A	803	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	803	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	755/793 (95%)	-0.38	7 (0%) 84 82	12, 19, 39, 74	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	758	VAL	4.9
1	A	285	TRP	3.0
1	A	759	PRO	2.9
1	A	35	ASN	2.9
1	A	456	TYR	2.2
1	A	448	GLU	2.0
1	A	730	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	A	801	6/6	0.87	0.17	28,30,37,43	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	A	804	6/6	0.89	0.18	31,49,56,63	0
2	GOL	A	803	6/6	0.90	0.27	40,45,52,53	0
3	MG	A	809	1/1	0.93	0.27	41,41,41,41	0
3	MG	A	806	1/1	0.94	0.22	36,36,36,36	0
2	GOL	A	802	6/6	0.95	0.08	31,32,33,33	0
2	GOL	A	805	6/6	0.96	0.11	21,30,37,40	0
3	MG	A	808	1/1	0.98	0.19	23,23,23,23	0
3	MG	A	807	1/1	0.98	0.05	17,17,17,17	0

6.5 Other polymers [i](#)

There are no such residues in this entry.