



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 31, 2023 – 07:57 PM JST

PDB ID : 5GZN
Title : Structure of neutralizing antibody bound to Zika envelope protein
Authors : Wang, Q.; Yang, H.; Liu, X.; Dai, L.; Ma, T.; Qi, J.; Wong, G.; Peng, R.; Liu, S.; Li, J.; Li, S.; Song, J.; Liu, J.; He, J.; Yuan, H.; Xiong, Y.; Liao, Y.; Li, J.; Yang, J.; Tong, Z.; Griffin, B.; Bi, Y.; Liang, M.; Xu, X.; Cheng, G.; Wang, P.; Qiu, X.; Kobinger, G.; Shi, Y.; Yan, J.; Gao, G.F.
Deposited on : 2016-09-29
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	409	
1	B	409	
1	E	409	
1	G	409	
2	C	228	
2	H	228	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	D	216	 <p>% 81% 17% **</p>
3	L	216	 <p>87% 11% **</p>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 15537 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Genome polyprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	402	Total 3071	C 1915	N 537	O 594	S 25	0	0	0
1	B	401	Total 3061	C 1909	N 534	O 593	S 25	0	0	0
1	E	184	Total 1431	C 895	N 250	O 274	S 12	0	0	0
1	G	183	Total 1429	C 896	N 249	O 272	S 12	0	0	0

- Molecule 2 is a protein called Antibody Heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	224	Total 1692	C 1070	N 286	O 330	S 6	0	0	0
2	C	223	Total 1686	C 1067	N 285	O 328	S 6	0	0	0

- Molecule 3 is a protein called Antibody light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	L	213	Total 1579	C 991	N 262	O 322	S 4	0	0	0
3	D	214	Total 1588	C 996	N 263	O 325	S 4	0	0	0



- Molecule 3: Antibody light chain

Chain L: ..



- Molecule 3: Antibody light chain

Chain D: ..



4 Data and refinement statistics i

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	124.71Å 129.38Å 428.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.08 – 3.00 42.87 – 2.99	Depositor EDS
% Data completeness (in resolution range)	98.3 (42.08-3.00) 98.2 (42.87-2.99)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.28 (at 3.01Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.223 , 0.258 0.228 , 0.259	Depositor DCC
R_{free} test set	3491 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	88.7	Xtrriage
Anisotropy	0.325	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 53.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.052 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	15537	wwPDB-VP
Average B, all atoms (Å ²)	97.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/3135	0.46	0/4247
1	B	0.26	0/3124	0.58	2/4232 (0.0%)
1	E	0.24	0/1463	0.45	0/1977
1	G	0.23	0/1460	0.45	0/1971
2	C	0.24	0/1730	0.49	1/2359 (0.0%)
2	H	0.26	0/1736	0.45	0/2367
3	D	0.24	0/1628	0.45	0/2226
3	L	0.25	0/1619	0.47	0/2214
All	All	0.25	0/15895	0.49	3/21593 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	146	SER	N-CA-CB	-21.96	77.56	110.50
2	C	160	PHE	C-N-CD	-7.01	105.17	120.60
1	B	144	HIS	CB-CA-C	-5.35	99.69	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3071	0	2996	32	0
1	B	3061	0	2989	37	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1431	0	1370	24	0
1	G	1429	0	1372	20	0
2	C	1686	0	1632	30	0
2	H	1692	0	1637	20	0
3	D	1588	0	1546	19	0
3	L	1579	0	1540	18	0
All	All	15537	0	15082	190	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 190 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:155:ASP:OD2	3:L:192:HIS:HB3	1.67	0.93
3:L:155:ASP:OD2	3:L:192:HIS:CB	2.27	0.82
2:C:160:PHE:HD2	2:C:161:PRO:HD3	1.47	0.80
2:C:160:PHE:CD2	2:C:161:PRO:HD3	2.18	0.78
3:L:155:ASP:CG	3:L:192:HIS:HB3	2.04	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	398/409 (97%)	366 (92%)	32 (8%)	0	100	100
1	B	397/409 (97%)	363 (91%)	34 (9%)	0	100	100
1	E	180/409 (44%)	168 (93%)	12 (7%)	0	100	100
1	G	177/409 (43%)	166 (94%)	11 (6%)	0	100	100
2	C	219/228 (96%)	192 (88%)	26 (12%)	1 (0%)	29	68

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	H	220/228 (96%)	194 (88%)	26 (12%)	0	100	100
3	D	212/216 (98%)	200 (94%)	12 (6%)	0	100	100
3	L	211/216 (98%)	200 (95%)	11 (5%)	0	100	100
All	All	2014/2524 (80%)	1849 (92%)	164 (8%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	161	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	336/341 (98%)	314 (94%)	22 (6%)	17	50
1	B	335/341 (98%)	317 (95%)	18 (5%)	22	57
1	E	152/341 (45%)	140 (92%)	12 (8%)	12	41
1	G	152/341 (45%)	141 (93%)	11 (7%)	14	45
2	C	187/192 (97%)	179 (96%)	8 (4%)	29	66
2	H	188/192 (98%)	177 (94%)	11 (6%)	19	54
3	D	180/182 (99%)	169 (94%)	11 (6%)	18	53
3	L	179/182 (98%)	169 (94%)	10 (6%)	21	56
All	All	1709/2112 (81%)	1606 (94%)	103 (6%)	19	53

5 of 103 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	276	GLU
3	D	64	SER
1	G	237	ASN
1	B	304	SER
2	C	84	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	158	HIS
1	E	226	HIS
1	E	249	HIS
1	G	253	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	402/409 (98%)	-0.17	1 (0%) 95 87	52, 84, 131, 224	0
1	B	401/409 (98%)	-0.10	2 (0%) 91 75	54, 85, 135, 203	0
1	E	184/409 (44%)	0.42	15 (8%) 11 3	76, 114, 186, 242	0
1	G	183/409 (44%)	0.29	18 (9%) 7 2	74, 108, 182, 219	0
2	C	223/228 (97%)	-0.06	3 (1%) 77 51	56, 97, 147, 179	0
2	H	224/228 (98%)	-0.11	3 (1%) 77 51	60, 98, 142, 192	0
3	D	214/216 (99%)	-0.14	3 (1%) 75 49	52, 79, 163, 202	0
3	L	213/216 (98%)	-0.17	0 100 100	48, 73, 144, 212	0
All	All	2044/2524 (80%)	-0.04	45 (2%) 62 33	48, 91, 154, 242	0

The worst 5 of 45 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	137	TYR	7.2
1	E	135	LEU	5.8
2	C	146	SER	5.5
1	G	137	TYR	5.4
1	G	135	LEU	4.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.