



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 1, 2023 – 12:25 PM JST

PDB ID : 5HFN
Title : Crystal structure of a loop truncation variant of *Thermotoga maritima* Acetyl Esterase TM0077 (apo structure) at 2.75 Angstrom resolution
Authors : Manoj, N.; Singh, M.K.
Deposited on : 2016-01-07
Resolution : 2.75 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1235 (2.78-2.74)
Clashscore	141614	1277 (2.78-2.74)
Ramachandran outliers	138981	1257 (2.78-2.74)
Sidechain outliers	138945	1257 (2.78-2.74)
RSRZ outliers	127900	1207 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	311	
1	B	311	
1	C	311	
1	D	311	
1	E	311	
1	F	311	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 13326 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cephalosporin-C deacetylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	280	2187	1425	363	390	9	0	0	0
1	B	278	2196	1436	361	390	9	0	1	0
1	C	279	2195	1436	360	390	9	0	1	0
1	D	283	2201	1444	356	391	10	0	1	0
1	E	279	2222	1450	365	398	9	0	1	0
1	F	279	2240	1461	370	400	9	0	1	0

There are 228 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	initiating methionine	UNP Q9WXT2
A	-10	GLY	-	expression tag	UNP Q9WXT2
A	-9	SER	-	expression tag	UNP Q9WXT2
A	-8	ASP	-	expression tag	UNP Q9WXT2
A	-7	LYS	-	expression tag	UNP Q9WXT2
A	-6	ILE	-	expression tag	UNP Q9WXT2
A	-5	HIS	-	expression tag	UNP Q9WXT2
A	-4	HIS	-	expression tag	UNP Q9WXT2
A	-3	HIS	-	expression tag	UNP Q9WXT2
A	-2	HIS	-	expression tag	UNP Q9WXT2
A	-1	HIS	-	expression tag	UNP Q9WXT2
A	0	HIS	-	expression tag	UNP Q9WXT2
A	?	-	GLN	deletion	UNP Q9WXT2
A	?	-	GLY	deletion	UNP Q9WXT2
A	?	-	SER	deletion	UNP Q9WXT2
A	?	-	GLY	deletion	UNP Q9WXT2
A	?	-	TRP	deletion	UNP Q9WXT2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	LEU	deletion	UNP Q9WXT2
A	?	-	LYS	deletion	UNP Q9WXT2
A	?	-	GLY	deletion	UNP Q9WXT2
A	?	-	ASP	deletion	UNP Q9WXT2
A	?	-	THR	deletion	UNP Q9WXT2
A	?	-	PRO	deletion	UNP Q9WXT2
A	?	-	ASP	deletion	UNP Q9WXT2
A	?	-	TYR	deletion	UNP Q9WXT2
A	?	-	PRO	deletion	UNP Q9WXT2
A	?	-	GLU	deletion	UNP Q9WXT2
A	?	-	GLY	deletion	UNP Q9WXT2
A	?	-	PRO	deletion	UNP Q9WXT2
A	?	-	VAL	deletion	UNP Q9WXT2
A	?	-	ASP	deletion	UNP Q9WXT2
A	?	-	PRO	deletion	UNP Q9WXT2
A	?	-	GLN	deletion	UNP Q9WXT2
A	?	-	TYR	deletion	UNP Q9WXT2
A	?	-	PRO	deletion	UNP Q9WXT2
A	?	-	GLY	deletion	UNP Q9WXT2
A	?	-	PHE	deletion	UNP Q9WXT2
A	?	-	MET	deletion	UNP Q9WXT2
B	-11	MET	-	initiating methionine	UNP Q9WXT2
B	-10	GLY	-	expression tag	UNP Q9WXT2
B	-9	SER	-	expression tag	UNP Q9WXT2
B	-8	ASP	-	expression tag	UNP Q9WXT2
B	-7	LYS	-	expression tag	UNP Q9WXT2
B	-6	ILE	-	expression tag	UNP Q9WXT2
B	-5	HIS	-	expression tag	UNP Q9WXT2
B	-4	HIS	-	expression tag	UNP Q9WXT2
B	-3	HIS	-	expression tag	UNP Q9WXT2
B	-2	HIS	-	expression tag	UNP Q9WXT2
B	-1	HIS	-	expression tag	UNP Q9WXT2
B	0	HIS	-	expression tag	UNP Q9WXT2
B	?	-	GLN	deletion	UNP Q9WXT2
B	?	-	GLY	deletion	UNP Q9WXT2
B	?	-	SER	deletion	UNP Q9WXT2
B	?	-	GLY	deletion	UNP Q9WXT2
B	?	-	TRP	deletion	UNP Q9WXT2
B	?	-	LEU	deletion	UNP Q9WXT2
B	?	-	LYS	deletion	UNP Q9WXT2
B	?	-	GLY	deletion	UNP Q9WXT2
B	?	-	ASP	deletion	UNP Q9WXT2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	THR	deletion	UNP Q9WXT2
B	?	-	PRO	deletion	UNP Q9WXT2
B	?	-	ASP	deletion	UNP Q9WXT2
B	?	-	TYR	deletion	UNP Q9WXT2
B	?	-	PRO	deletion	UNP Q9WXT2
B	?	-	GLU	deletion	UNP Q9WXT2
B	?	-	GLY	deletion	UNP Q9WXT2
B	?	-	PRO	deletion	UNP Q9WXT2
B	?	-	VAL	deletion	UNP Q9WXT2
B	?	-	ASP	deletion	UNP Q9WXT2
B	?	-	PRO	deletion	UNP Q9WXT2
B	?	-	GLN	deletion	UNP Q9WXT2
B	?	-	TYR	deletion	UNP Q9WXT2
B	?	-	PRO	deletion	UNP Q9WXT2
B	?	-	GLY	deletion	UNP Q9WXT2
B	?	-	PHE	deletion	UNP Q9WXT2
B	?	-	MET	deletion	UNP Q9WXT2
C	-11	MET	-	initiating methionine	UNP Q9WXT2
C	-10	GLY	-	expression tag	UNP Q9WXT2
C	-9	SER	-	expression tag	UNP Q9WXT2
C	-8	ASP	-	expression tag	UNP Q9WXT2
C	-7	LYS	-	expression tag	UNP Q9WXT2
C	-6	ILE	-	expression tag	UNP Q9WXT2
C	-5	HIS	-	expression tag	UNP Q9WXT2
C	-4	HIS	-	expression tag	UNP Q9WXT2
C	-3	HIS	-	expression tag	UNP Q9WXT2
C	-2	HIS	-	expression tag	UNP Q9WXT2
C	-1	HIS	-	expression tag	UNP Q9WXT2
C	0	HIS	-	expression tag	UNP Q9WXT2
C	?	-	GLN	deletion	UNP Q9WXT2
C	?	-	GLY	deletion	UNP Q9WXT2
C	?	-	SER	deletion	UNP Q9WXT2
C	?	-	GLY	deletion	UNP Q9WXT2
C	?	-	TRP	deletion	UNP Q9WXT2
C	?	-	LEU	deletion	UNP Q9WXT2
C	?	-	LYS	deletion	UNP Q9WXT2
C	?	-	GLY	deletion	UNP Q9WXT2
C	?	-	ASP	deletion	UNP Q9WXT2
C	?	-	THR	deletion	UNP Q9WXT2
C	?	-	PRO	deletion	UNP Q9WXT2
C	?	-	ASP	deletion	UNP Q9WXT2
C	?	-	TYR	deletion	UNP Q9WXT2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	?	-	PRO	deletion	UNP Q9WXT2
C	?	-	GLU	deletion	UNP Q9WXT2
C	?	-	GLY	deletion	UNP Q9WXT2
C	?	-	PRO	deletion	UNP Q9WXT2
C	?	-	VAL	deletion	UNP Q9WXT2
C	?	-	ASP	deletion	UNP Q9WXT2
C	?	-	PRO	deletion	UNP Q9WXT2
C	?	-	GLN	deletion	UNP Q9WXT2
C	?	-	TYR	deletion	UNP Q9WXT2
C	?	-	PRO	deletion	UNP Q9WXT2
C	?	-	GLY	deletion	UNP Q9WXT2
C	?	-	PHE	deletion	UNP Q9WXT2
C	?	-	MET	deletion	UNP Q9WXT2
D	-11	MET	-	initiating methionine	UNP Q9WXT2
D	-10	GLY	-	expression tag	UNP Q9WXT2
D	-9	SER	-	expression tag	UNP Q9WXT2
D	-8	ASP	-	expression tag	UNP Q9WXT2
D	-7	LYS	-	expression tag	UNP Q9WXT2
D	-6	ILE	-	expression tag	UNP Q9WXT2
D	-5	HIS	-	expression tag	UNP Q9WXT2
D	-4	HIS	-	expression tag	UNP Q9WXT2
D	-3	HIS	-	expression tag	UNP Q9WXT2
D	-2	HIS	-	expression tag	UNP Q9WXT2
D	-1	HIS	-	expression tag	UNP Q9WXT2
D	0	HIS	-	expression tag	UNP Q9WXT2
D	?	-	GLN	deletion	UNP Q9WXT2
D	?	-	GLY	deletion	UNP Q9WXT2
D	?	-	SER	deletion	UNP Q9WXT2
D	?	-	GLY	deletion	UNP Q9WXT2
D	?	-	TRP	deletion	UNP Q9WXT2
D	?	-	LEU	deletion	UNP Q9WXT2
D	?	-	LYS	deletion	UNP Q9WXT2
D	?	-	GLY	deletion	UNP Q9WXT2
D	?	-	ASP	deletion	UNP Q9WXT2
D	?	-	THR	deletion	UNP Q9WXT2
D	?	-	PRO	deletion	UNP Q9WXT2
D	?	-	ASP	deletion	UNP Q9WXT2
D	?	-	TYR	deletion	UNP Q9WXT2
D	?	-	PRO	deletion	UNP Q9WXT2
D	?	-	GLU	deletion	UNP Q9WXT2
D	?	-	GLY	deletion	UNP Q9WXT2
D	?	-	PRO	deletion	UNP Q9WXT2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	?	-	VAL	deletion	UNP Q9WXT2
D	?	-	ASP	deletion	UNP Q9WXT2
D	?	-	PRO	deletion	UNP Q9WXT2
D	?	-	GLN	deletion	UNP Q9WXT2
D	?	-	TYR	deletion	UNP Q9WXT2
D	?	-	PRO	deletion	UNP Q9WXT2
D	?	-	GLY	deletion	UNP Q9WXT2
D	?	-	PHE	deletion	UNP Q9WXT2
D	?	-	MET	deletion	UNP Q9WXT2
E	-11	MET	-	initiating methionine	UNP Q9WXT2
E	-10	GLY	-	expression tag	UNP Q9WXT2
E	-9	SER	-	expression tag	UNP Q9WXT2
E	-8	ASP	-	expression tag	UNP Q9WXT2
E	-7	LYS	-	expression tag	UNP Q9WXT2
E	-6	ILE	-	expression tag	UNP Q9WXT2
E	-5	HIS	-	expression tag	UNP Q9WXT2
E	-4	HIS	-	expression tag	UNP Q9WXT2
E	-3	HIS	-	expression tag	UNP Q9WXT2
E	-2	HIS	-	expression tag	UNP Q9WXT2
E	-1	HIS	-	expression tag	UNP Q9WXT2
E	0	HIS	-	expression tag	UNP Q9WXT2
E	?	-	GLN	deletion	UNP Q9WXT2
E	?	-	GLY	deletion	UNP Q9WXT2
E	?	-	SER	deletion	UNP Q9WXT2
E	?	-	GLY	deletion	UNP Q9WXT2
E	?	-	TRP	deletion	UNP Q9WXT2
E	?	-	LEU	deletion	UNP Q9WXT2
E	?	-	LYS	deletion	UNP Q9WXT2
E	?	-	GLY	deletion	UNP Q9WXT2
E	?	-	ASP	deletion	UNP Q9WXT2
E	?	-	THR	deletion	UNP Q9WXT2
E	?	-	PRO	deletion	UNP Q9WXT2
E	?	-	ASP	deletion	UNP Q9WXT2
E	?	-	TYR	deletion	UNP Q9WXT2
E	?	-	PRO	deletion	UNP Q9WXT2
E	?	-	GLU	deletion	UNP Q9WXT2
E	?	-	GLY	deletion	UNP Q9WXT2
E	?	-	PRO	deletion	UNP Q9WXT2
E	?	-	VAL	deletion	UNP Q9WXT2
E	?	-	ASP	deletion	UNP Q9WXT2
E	?	-	PRO	deletion	UNP Q9WXT2
E	?	-	GLN	deletion	UNP Q9WXT2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	?	-	TYR	deletion	UNP Q9WXT2
E	?	-	PRO	deletion	UNP Q9WXT2
E	?	-	GLY	deletion	UNP Q9WXT2
E	?	-	PHE	deletion	UNP Q9WXT2
E	?	-	MET	deletion	UNP Q9WXT2
F	-11	MET	-	initiating methionine	UNP Q9WXT2
F	-10	GLY	-	expression tag	UNP Q9WXT2
F	-9	SER	-	expression tag	UNP Q9WXT2
F	-8	ASP	-	expression tag	UNP Q9WXT2
F	-7	LYS	-	expression tag	UNP Q9WXT2
F	-6	ILE	-	expression tag	UNP Q9WXT2
F	-5	HIS	-	expression tag	UNP Q9WXT2
F	-4	HIS	-	expression tag	UNP Q9WXT2
F	-3	HIS	-	expression tag	UNP Q9WXT2
F	-2	HIS	-	expression tag	UNP Q9WXT2
F	-1	HIS	-	expression tag	UNP Q9WXT2
F	0	HIS	-	expression tag	UNP Q9WXT2
F	?	-	GLN	deletion	UNP Q9WXT2
F	?	-	GLY	deletion	UNP Q9WXT2
F	?	-	SER	deletion	UNP Q9WXT2
F	?	-	GLY	deletion	UNP Q9WXT2
F	?	-	TRP	deletion	UNP Q9WXT2
F	?	-	LEU	deletion	UNP Q9WXT2
F	?	-	LYS	deletion	UNP Q9WXT2
F	?	-	GLY	deletion	UNP Q9WXT2
F	?	-	ASP	deletion	UNP Q9WXT2
F	?	-	THR	deletion	UNP Q9WXT2
F	?	-	PRO	deletion	UNP Q9WXT2
F	?	-	ASP	deletion	UNP Q9WXT2
F	?	-	TYR	deletion	UNP Q9WXT2
F	?	-	PRO	deletion	UNP Q9WXT2
F	?	-	GLU	deletion	UNP Q9WXT2
F	?	-	GLY	deletion	UNP Q9WXT2
F	?	-	PRO	deletion	UNP Q9WXT2
F	?	-	VAL	deletion	UNP Q9WXT2
F	?	-	ASP	deletion	UNP Q9WXT2
F	?	-	PRO	deletion	UNP Q9WXT2
F	?	-	GLN	deletion	UNP Q9WXT2
F	?	-	TYR	deletion	UNP Q9WXT2
F	?	-	PRO	deletion	UNP Q9WXT2
F	?	-	GLY	deletion	UNP Q9WXT2
F	?	-	PHE	deletion	UNP Q9WXT2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	?	-	MET	deletion	UNP Q9WXT2


- Molecule 2 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	12	Total O 12 12	0	0
2	B	12	Total O 12 12	0	0
2	C	12	Total O 12 12	0	0
2	D	15	Total O 15 15	0	0
2	E	20	Total O 20 20	0	0
2	F	14	Total O 14 14	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cephalosporin-C deacetylase

Chain A: 





- Molecule 1: Cephalosporin-C deacetylase

Chain B: 





- Molecule 1: Cephalosporin-C deacetylase

Chain C: 




- Molecule 1: Cephalosporin-C deacetylase

Chain D: 



E323
LYS
GLY


- Molecule 1: Cephalosporin-C deacetylase

Chain E:  83% 6% 10%

MET
GLY
SER
ASP
LYS
ILE
HIS
HIS
HIS
HIS
HIS
HIS
MET
MET
H2
F3
M47
E55
A56
Y57
L73
E81
I90
GLY
TVR
ASN
GLY
GLY
R96
G97
T117
ARG
GLY
THR
ARG
GLY
ILE
LEU
ASP
PRO
ARG
THR
TYR
Y156
S168
V221
Q222
H227
E231
H239

R240
D241
K242
P265
P292
H303
E323
LYS
GLY

- Molecule 1: Cephalosporin-C deacetylase

Chain F:  83% 7% 10%

MET
GLY
SER
ASP
LYS
ILE
HIS
HIS
HIS
HIS
HIS
MET
MET
ALA
PHE
F4
G67
I96
GLY
TYR
ASN
GLY
G95
R96
G97
R118
GLY
THR
ARG
ARG
GLY
ILE
LEU
ASP
PRO
ARG
THR
TYR
Y156
R159
E180
S188
Q189
R218
V221
Q222
H227
P228
E231
F235

H239
R240
D241
K242
R248
P265
P292
H303
E313
E323
LYS
GLY

4 Data and refinement statistics i

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	155.08Å 155.08Å 165.10Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	52.09 – 2.75 52.09 – 2.75	Depositor EDS
% Data completeness (in resolution range)	99.9 (52.09-2.75) 99.9 (52.09-2.75)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.20 (at 2.77Å)	Xtrriage
Refinement program	REFMAC 5	Depositor
R, R_{free}	0.198 , 0.231 0.205 , 0.236	Depositor DCC
R_{free} test set	3028 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	50.9	Xtrriage
Anisotropy	0.362	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 43.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	0.010 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	13326	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.96	0/2249	0.94	3/3057 (0.1%)
1	B	1.02	3/2263 (0.1%)	0.94	2/3076 (0.1%)
1	C	0.89	1/2262 (0.0%)	0.86	1/3076 (0.0%)
1	D	0.92	0/2269	0.90	2/3088 (0.1%)
1	E	1.04	3/2290 (0.1%)	0.94	4/3111 (0.1%)
1	F	1.00	3/2308 (0.1%)	0.99	11/3132 (0.4%)
All	All	0.97	10/13641 (0.1%)	0.93	23/18540 (0.1%)

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	188	SER	CB-OG	-6.88	1.33	1.42
1	F	313	GLU	CD-OE2	-6.75	1.18	1.25
1	C	188	SER	CB-OG	-6.27	1.34	1.42
1	E	55	GLU	CD-OE1	5.97	1.32	1.25
1	F	180	GLU	CD-OE2	-5.75	1.19	1.25

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	241	ASP	CB-CG-OD1	8.94	126.35	118.30
1	A	241	ASP	CB-CG-OD1	8.83	126.25	118.30
1	D	241	ASP	CB-CG-OD1	8.61	126.05	118.30
1	C	241	ASP	CB-CG-OD1	8.39	125.85	118.30
1	E	241	ASP	CB-CG-OD1	8.23	125.71	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2187	0	2067	6	0
1	B	2196	0	2086	8	0
1	C	2195	0	2079	8	0
1	D	2201	0	2069	7	0
1	E	2222	0	2107	10	0
1	F	2240	0	2150	8	0
2	A	12	0	0	1	0
2	B	12	0	0	0	0
2	C	12	0	0	0	0
2	D	15	0	0	0	0
2	E	20	0	0	0	0
2	F	14	0	0	0	0
All	All	13326	0	12558	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 43 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:3:PHE:CD2	1:E:222:GLN:HB2	2.22	0.75
1:C:118:ARG:HH21	1:C:159:ARG:CB	2.06	0.67
1:D:188:SER:OG	1:D:303:HIS:NE2	2.43	0.51
1:A:188:SER:OG	1:A:303:HIS:NE2	2.45	0.49
1:C:188:SER:OG	1:C:303:HIS:NE2	2.45	0.49

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	274/311 (88%)	264 (96%)	9 (3%)	1 (0%)	34	53
1	B	273/311 (88%)	267 (98%)	6 (2%)	0	100	100
1	C	274/311 (88%)	265 (97%)	8 (3%)	1 (0%)	34	53
1	D	278/311 (89%)	267 (96%)	10 (4%)	1 (0%)	34	53
1	E	274/311 (88%)	266 (97%)	7 (3%)	1 (0%)	34	53
1	F	274/311 (88%)	268 (98%)	5 (2%)	1 (0%)	34	53
All	All	1647/1866 (88%)	1597 (97%)	45 (3%)	5 (0%)	41	60

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	97	GLY
1	F	97	GLY
1	A	97	GLY
1	D	97	GLY
1	C	97	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	217/264 (82%)	216 (100%)	1 (0%)	88	92
1	B	221/264 (84%)	217 (98%)	4 (2%)	59	75
1	C	220/264 (83%)	219 (100%)	1 (0%)	88	92
1	D	217/264 (82%)	216 (100%)	1 (0%)	88	92
1	E	225/264 (85%)	224 (100%)	1 (0%)	91	93
1	F	230/264 (87%)	228 (99%)	2 (1%)	78	87
All	All	1330/1584 (84%)	1320 (99%)	10 (1%)	81	88

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	221	VAL
1	F	118	ARG
1	F	221	VAL
1	B	221	VAL
1	B	238	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	100	HIS
1	F	239	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	280/311 (90%)	-0.07	1 (0%) 92 95	34, 49, 81, 99	0
1	B	278/311 (89%)	-0.11	0 100 100	37, 50, 72, 94	0
1	C	279/311 (89%)	0.13	7 (2%) 57 66	39, 60, 89, 109	0
1	D	283/311 (90%)	-0.04	3 (1%) 80 86	30, 56, 86, 102	0
1	E	279/311 (89%)	-0.08	2 (0%) 87 91	34, 44, 65, 85	0
1	F	279/311 (89%)	0.02	1 (0%) 92 95	34, 49, 77, 93	0
All	All	1678/1866 (89%)	-0.02	14 (0%) 86 90	30, 51, 81, 109	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	157	TYR	3.2
1	E	2	ALA	2.7
1	D	-2	HIS	2.7
1	D	156	TYR	2.6
1	C	157	TYR	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.