



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 19, 2023 – 08:05 PM EDT

PDB ID : 5HP2  
Title : Human Adenosine Deaminase Acting on dsRNA (ADAR2) bound to dsRNA sequence derived from *S. cerevisiae* BDF2 gene with AU basepair at reaction site  
Authors : Matthews, M.M.; Fisher, A.J.; Beal, P.A.  
Deposited on : 2016-01-20  
Resolution : 2.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2754 (3.00-2.96)
Clashscore	141614	3103 (3.00-2.96)
Ramachandran outliers	138981	2993 (3.00-2.96)
Sidechain outliers	138945	2996 (3.00-2.96)
RSRZ outliers	127900	2644 (3.00-2.96)
RNA backbone	3102	1088 (3.26-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	403	
1	D	403	
2	B	23	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	C	23	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into three segments: a red segment on the left labeled '13%', a green segment in the middle labeled '61%', and a yellow segment on the right labeled '35%'. A small black dot is visible at the far right end of the bar.</p>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7216 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Double-stranded RNA-specific editase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	396	Total	C	N	O	S	0	0	0
			3096	1952	562	571	11			
1	D	397	Total	C	N	O	S	0	0	0
			3072	1940	553	568	11			


- Molecule 2 is a RNA chain called RNA (5'-R(\*UP\*UP\*CP\*CP\*CP\*CP\*AP\*CP\*AP\*UP\*UP\*(8AZ)P\*GP\*AP\*CP\*GP\*UP\*UP\*CP\*AP\*GP\*UP\*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	B	23	Total	C	N	O	P	0	0	0
			476	214	78	162	22			

- Molecule 3 is a RNA chain called RNA (5'-R(\*GP\*AP\*CP\*UP\*GP\*AP\*AP\*CP\*GP\*AP\*CP\*UP\*AP\*AP\*UP\*GP\*UP\*GP\*GP\*GP\*GP\*AP\*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	C	23	Total	C	N	O	P	0	0	0
			497	223	97	155	22			

- Molecule 4 is INOSITOL HEXAKISPHOSPHATE (three-letter code: IHP) (formula: C<sub>6</sub>H<sub>18</sub>O<sub>24</sub>P<sub>6</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O P 36 6 24 6	0	0
4	D	1	Total C O P 36 6 24 6	0	0

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Zn 1 1	0	0
5	D	1	Total Zn 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	D	1	Total O 1 1	0	0


### 3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Double-stranded RNA-specific editase 1




- Molecule 1: Double-stranded RNA-specific editase 1



- Molecule 2: RNA (5'-R(\*UP\*UP\*CP\*CP\*CP\*CP\*AP\*CP\*AP\*UP\*UP\*(8AZ)P\*GP\*AP\*CP\*GP\*UP\*UP\*CP\*AP\*GP\*UP\*C)-3')





- Molecule 3: RNA (5'-R(\*GP\*AP\*CP\*UP\*GP\*AP\*AP\*CP\*GP\*AP\*CP\*UP\*AP\*AP\*UP\*GP\*UP\*GP\*GP\*GP\*GP\*AP\*A)-3')



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.32Å 106.68Å 120.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.53 – 2.98 38.53 – 2.98	Depositor EDS
% Data completeness (in resolution range)	97.3 (38.53-2.98) 97.4 (38.53-2.98)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.22 (at 3.01Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.167 , 0.247 0.169 , 0.248	Depositor DCC
$R_{free}$ test set	1086 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	60.7	Xtrriage
Anisotropy	0.669	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 52.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7216	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	64.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.61% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 8AZ, IHP, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.46	0/3160	0.65	0/4274
1	D	0.43	0/3135	0.63	0/4245
2	B	0.57	0/503	1.15	4/777 (0.5%)
3	C	0.52	0/558	1.06	1/870 (0.1%)
All	All	0.46	0/7356	0.74	5/10166 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	14	A	N1-C6-N6	6.10	122.26	118.60
3	C	10	A	C2-N3-C4	-5.79	107.70	110.60
2	B	14	A	C4-C5-N7	5.41	113.41	110.70
2	B	14	A	N9-C4-C5	-5.19	103.72	105.80
2	B	14	A	C5-C6-N6	-5.14	119.59	123.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3096	0	3119	35	1
1	D	3072	0	3071	60	1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	476	0	237	4	0
3	C	497	0	251	5	0
4	A	36	0	6	1	0
4	D	36	0	6	1	0
5	A	1	0	0	0	0
5	D	1	0	0	0	0
6	D	1	0	0	0	0
All	All	7216	0	6690	99	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (99) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:392:ASP:OD2	1:D:483:LYS:NZ	2.18	0.75
1:D:494:ARG:HB2	1:D:497:ALA:HB2	1.69	0.75
1:D:420:ASP:OD1	1:D:423:ARG:NH1	2.25	0.70
1:D:464:LEU:O	1:D:466:GLU:N	2.21	0.69
2:B:14:A:N1	3:C:10:A:H2	1.92	0.68
1:A:519:LYS:NZ	4:A:801:IHP:O31	2.25	0.67
1:D:400:ARG:HD3	1:D:523:TRP:CE2	2.30	0.67
1:D:481:ARG:HE	1:D:490:THR:CG2	2.09	0.66
1:D:434:PHE:HB2	1:D:537:VAL:HG12	1.79	0.62
1:A:392:ASP:OD2	1:A:483:LYS:NZ	2.31	0.62
1:A:652:ILE:HD11	1:A:654:LYS:HE2	1.82	0.62
1:A:461:GLU:HG3	1:A:462:PRO:HD2	1.81	0.61
1:A:400:ARG:NH1	1:A:528:ILE:O	2.33	0.60
1:D:552:HIS:ND1	1:D:554:ASP:OD1	2.33	0.59
1:A:404:LEU:HD22	1:A:533:LEU:HD12	1.84	0.58
1:A:400:ARG:HD3	1:A:523:TRP:CE2	2.38	0.58
1:A:475:LYS:O	1:A:479:GLN:NE2	2.36	0.57
2:B:11:U:H3	3:C:13:A:H61	1.52	0.57
1:A:400:ARG:HD3	1:A:523:TRP:CZ2	2.40	0.57
1:D:319:LEU:HB2	1:D:322:VAL:HG12	1.87	0.56
1:D:458:SER:HB3	1:D:461:GLU:HG3	1.87	0.55
1:D:483:LYS:HE3	1:D:515:SER:HA	1.89	0.54
1:A:350:LYS:NZ	1:A:591:GLN:O	2.37	0.54
1:A:665:ALA:O	1:A:669:GLN:HB2	2.08	0.54
1:A:464:LEU:O	1:A:467:PRO:HD3	2.07	0.53
1:D:400:ARG:HD3	1:D:523:TRP:CZ2	2.44	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:393:CYS:HA	1:D:398:ILE:HD11	1.91	0.53
1:A:396:GLU:OE2	2:B:12:8AZ:O6	2.27	0.53
1:D:377:CYS:SG	1:D:483:LYS:HD3	2.50	0.52
1:D:467:PRO:C	1:D:469:ASP:H	2.13	0.52
1:D:400:ARG:NH1	1:D:528:ILE:O	2.38	0.51
1:D:437:LYS:O	1:D:440:VAL:HG12	2.10	0.51
1:D:312:SER:HB3	1:D:580:LEU:N	2.27	0.50
1:D:514:MET:HG3	1:D:687:TRP:CE2	2.47	0.50
1:D:466:GLU:O	1:D:468:ALA:N	2.45	0.49
1:D:404:LEU:HD22	1:D:533:LEU:HD12	1.95	0.49
1:D:481:ARG:HB3	1:D:490:THR:HG23	1.94	0.48
1:A:451:CYS:HA	1:A:455:ARG:HD2	1.95	0.48
1:A:485:GLU:O	1:A:510:ARG:HD2	2.13	0.48
1:A:516:CYS:O	1:A:520:ILE:HG13	2.13	0.47
1:D:561:TYR:CD1	1:D:578:LYS:HG2	2.50	0.47
1:D:349:ARG:HD2	1:D:352:LEU:HB2	1.96	0.47
1:D:397:ILE:HD12	1:D:519:LYS:HD2	1.97	0.46
1:D:372:SER:OG	1:D:399:SER:OG	2.23	0.46
1:A:306:SER:HB3	1:A:467:PRO:HD2	1.99	0.45
1:A:464:LEU:C	1:A:467:PRO:HD3	2.37	0.45
1:A:430:GLU:H	1:A:430:GLU:HG3	1.46	0.45
1:A:602:TRP:CH2	1:A:604:VAL:HA	2.52	0.45
1:D:516:CYS:O	1:D:520:ILE:HG13	2.17	0.45
1:A:519:LYS:O	1:A:522:ARG:HB3	2.17	0.44
1:D:312:SER:H	1:D:316:GLN:HE22	1.65	0.44
1:A:339:THR:HB	1:A:344:SER:HB3	1.98	0.44
1:D:452:GLY:HA2	1:D:490:THR:HG21	1.99	0.44
1:D:561:TYR:CG	1:D:578:LYS:HG2	2.52	0.44
1:A:439:ASN:OD1	1:A:439:ASN:N	2.50	0.44
1:A:393:CYS:HA	1:A:398:ILE:HD11	1.99	0.44
1:A:459:PRO:HD2	1:A:460:HIS:CD2	2.53	0.44
1:D:494:ARG:HD2	1:D:510:ARG:HH22	1.83	0.44
1:A:372:SER:OG	1:A:399:SER:OG	2.22	0.43
1:D:349:ARG:CD	1:D:352:LEU:HB2	2.47	0.43
1:D:349:ARG:NH1	1:D:588:GLU:OE2	2.48	0.43
1:D:425:ILE:O	1:D:437:LYS:HG3	2.17	0.43
1:A:466:GLU:O	1:A:468:ALA:N	2.43	0.43
1:A:556:LEU:O	1:A:560:MET:HG2	2.18	0.43
1:D:350:LYS:HB3	1:D:375:THR:HG21	2.01	0.43
1:D:453:ASP:CG	1:D:478:GLY:HA2	2.39	0.43
1:D:467:PRO:HG3	1:D:472:PRO:HD3	2.01	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:537:VAL:O	1:D:649:ARG:NH1	2.52	0.42
1:A:514:MET:HG3	1:A:687:TRP:CE2	2.54	0.42
3:C:16:G:H2'	3:C:17:U:C6	2.54	0.42
3:C:18:G:H2'	3:C:19:G:C8	2.54	0.42
1:A:561:TYR:CD2	1:A:578:LYS:HG2	2.54	0.42
1:D:602:TRP:CD1	1:D:606:ASP:HB2	2.55	0.42
1:D:318:HIS:CD2	1:D:363:VAL:HG13	2.54	0.42
1:A:567:ILE:HD11	1:A:570:LEU:HD21	2.02	0.42
1:D:394:HIS:CE1	1:D:483:LYS:HE2	2.55	0.42
1:D:457:PHE:O	1:D:459:PRO:HD3	2.20	0.42
1:D:554:ASP:OD1	1:D:554:ASP:N	2.53	0.42
1:D:602:TRP:CZ2	1:D:604:VAL:HA	2.55	0.42
1:A:483:LYS:N	1:A:513:THR:O	2.51	0.42
1:D:610:GLU:OE1	1:D:626:ARG:HG2	2.20	0.41
1:D:431:ARG:HH22	1:D:435:ARG:CZ	2.32	0.41
1:D:601:ASN:ND2	1:D:610:GLU:HB3	2.35	0.41
1:D:494:ARG:HD2	1:D:510:ARG:NH2	2.35	0.41
2:B:5:C:H2'	2:B:6:C:C6	2.55	0.41
1:D:360:GLY:HA2	1:D:439:ASN:HD22	1.85	0.41
1:D:436:LEU:HB3	1:D:440:VAL:CG1	2.50	0.41
1:D:502:TRP:HE3	1:D:691:PRO:HG2	1.86	0.41
1:A:488:GLU:HG2	3:C:12:U:O2	2.20	0.41
1:D:340:ASP:O	1:D:343:SER:OG	2.30	0.41
1:D:558:ARG:O	1:D:563:ARG:HG2	2.21	0.41
1:D:411:LEU:O	1:D:415:LEU:HG	2.20	0.41
1:D:690:LYS:HG2	4:D:801:IHP:O45	2.20	0.41
1:D:308:GLN:HA	1:D:309:PRO:HD2	1.83	0.40
1:D:319:LEU:O	1:D:322:VAL:HG12	2.21	0.40
1:D:602:TRP:NE1	1:D:606:ASP:HB2	2.37	0.40
1:A:397:ILE:HD12	1:A:519:LYS:HD2	2.03	0.40
1:A:435:ARG:HH12	1:D:616:THR:HG21	1.87	0.40
1:D:456:ILE:O	1:D:457:PHE:HB3	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:344:SER:OG	1:D:469:ASP:OD2[2_445]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	394/403 (98%)	374 (95%)	18 (5%)	2 (0%)	29	66
1	D	395/403 (98%)	366 (93%)	25 (6%)	4 (1%)	15	50
All	All	789/806 (98%)	740 (94%)	43 (5%)	6 (1%)	19	55

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	465	GLU
1	D	470	ARG
1	D	466	GLU
1	D	468	ALA
1	A	497	ALA
1	A	499	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	337/347 (97%)	326 (97%)	11 (3%)	38	71
1	D	330/347 (95%)	319 (97%)	11 (3%)	38	71
All	All	667/694 (96%)	645 (97%)	22 (3%)	38	71

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	363	VAL
1	A	415	LEU
1	A	416	ASN
1	A	430	GLU
1	A	457	PHE
1	A	464	LEU
1	A	479	GLN
1	A	496	ASN
1	A	550	LEU
1	A	652	ILE
1	A	669	GLN
1	D	363	VAL
1	D	415	LEU
1	D	416	ASN
1	D	419	ASP
1	D	477	ARG
1	D	488	GLU
1	D	490	THR
1	D	597	ASN
1	D	633	TYR
1	D	652	ILE
1	D	669	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	318	HIS
1	A	441	GLN
1	A	496	ASN
1	A	562	GLN
1	A	591	GLN
1	D	316	GLN
1	D	318	HIS
1	D	416	ASN
1	D	427	GLN
1	D	496	ASN
1	D	601	ASN

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	21/23 (91%)	1 (4%)	0
3	C	22/23 (95%)	2 (9%)	0
All	All	43/46 (93%)	3 (6%)	0

All (3) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	B	23	C
3	C	6	A
3	C	21	G

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	8AZ	B	12	2,5	17,24,25	0.73	0	14,35,38	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	8AZ	B	12	2,5	-	2/3/35/36	0/3/3/3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	12	8AZ	O4'-C4'-C5'-O5'
2	B	12	8AZ	C3'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	12	8AZ	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	IHP	A	801	-	36,36,36	0.84	2 (5%)	54,60,60	1.22	6 (11%)
4	IHP	D	801	-	36,36,36	0.89	1 (2%)	54,60,60	1.14	2 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	IHP	A	801	-	-	3/30/54/54	0/1/1/1
4	IHP	D	801	-	-	6/30/54/54	0/1/1/1

All (3) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	801	IHP	P2-O12	2.38	1.63	1.59
4	A	801	IHP	P4-O14	2.26	1.63	1.59
4	A	801	IHP	P1-O11	2.19	1.63	1.59

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	801	IHP	O16-P6-O26	-2.78	98.65	109.39
4	A	801	IHP	O16-P6-O26	-2.66	99.14	109.39
4	A	801	IHP	O36-P6-O26	2.58	120.77	110.68
4	D	801	IHP	O46-P6-O26	2.21	119.32	110.68
4	A	801	IHP	O42-P2-O32	2.18	115.97	107.64
4	A	801	IHP	O41-P1-O31	2.14	115.81	107.64
4	A	801	IHP	O14-C4-C5	2.06	113.55	108.69
4	A	801	IHP	O14-C4-C3	-2.03	103.90	108.69

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	801	IHP	C1-O11-P1-O21
4	D	801	IHP	C1-O11-P1-O21
4	D	801	IHP	C3-O13-P3-O23
4	A	801	IHP	C4-O14-P4-O44
4	A	801	IHP	C6-O16-P6-O46
4	D	801	IHP	C1-O11-P1-O31
4	D	801	IHP	C2-O12-P2-O42
4	D	801	IHP	C3-O13-P3-O33
4	D	801	IHP	C4-O14-P4-O34


There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	801	IHP	1	0
4	D	801	IHP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	396/403 (98%)	0.02	7 (1%) 68 48	29, 49, 102, 154	0
1	D	397/403 (98%)	0.44	34 (8%) 10 5	36, 66, 121, 155	0
2	B	22/23 (95%)	0.57	2 (9%) 9 5	39, 64, 89, 100	0
3	C	23/23 (100%)	0.64	3 (13%) 3 1	43, 65, 93, 95	0
All	All	838/852 (98%)	0.25	46 (5%) 25 14	29, 58, 113, 155	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	472	PRO	7.9
1	D	473	ASN	7.4
1	D	505	VAL	5.9
1	D	471	HIS	5.6
1	D	503	ASP	5.5
1	D	655	PRO	5.2
1	D	501	THR	5.2
1	A	465	GLU	4.1
1	D	502	TRP	4.0
1	D	497	ALA	4.0
1	D	653	THR	3.9
1	D	654	LYS	3.8
1	D	498	SER	3.8
1	D	496	ASN	3.7
1	A	468	ALA	3.6
1	D	507	GLN	3.5
1	A	463	ILE	3.4
1	D	510	ARG	3.2
1	A	467	PRO	3.2
1	A	464	LEU	3.2
3	C	10	A	3.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	504	GLY	3.1
1	D	495	SER	3.0
1	D	308	GLN	3.0
1	D	508	GLY	2.9
3	C	9	G	2.9
1	D	509	GLU	2.8
1	D	486	SER	2.7
1	D	666	LYS	2.6
1	D	467	PRO	2.6
1	A	305	PRO	2.6
1	D	319	LEU	2.5
1	D	494	ARG	2.5
1	A	645	SER	2.4
1	D	465	GLU	2.3
2	B	5	C	2.3
1	D	506	LEU	2.3
1	D	464	LEU	2.3
2	B	6	C	2.2
3	C	19	G	2.2
1	D	652	ILE	2.2
1	D	527	GLY	2.1
1	D	499	ILE	2.1
1	D	500	GLN	2.1
1	D	309	PRO	2.0
1	D	339	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	8AZ	B	12	22/23	0.98	0.22	30,43,54,62	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	IHP	D	801	36/36	0.97	0.22	27,48,63,76	0
4	IHP	A	801	36/36	0.99	0.21	21,37,53,58	0
5	ZN	A	802	1/1	0.99	0.13	34,34,34,34	0
5	ZN	D	802	1/1	0.99	0.16	42,42,42,42	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around IHP D 801:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.