



# Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 05:53 pm BST

PDB ID : 5J77  
Title : Mutant glyceraldehyde dehydrogenase (F34M+S405N) from *Thermoplasma acidophilum*  
Authors : Iermak, I.; Mesters, J.R.; Kuta Smatanova, I.  
Deposited on : 2016-04-06  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	508	94%
1	B	508	93%
1	C	508	3% 93%
1	D	508	3% 92%

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 15691 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D-glyceraldehyde dehydrogenase (NADP(+)).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	492	3806	2430	630	730	16	0	3	0
1	B	492	3818	2436	634	732	16	0	5	0
1	C	492	3745	2397	610	724	14	0	2	0
1	D	492	3744	2395	616	718	15	0	2	0

There are 68 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	34	MET	PHE	engineered mutation	UNP Q9HK01
A	405	ASN	SER	engineered mutation	UNP Q9HK01
A	494	SER	-	expression tag	UNP Q9HK01
A	495	GLY	-	expression tag	UNP Q9HK01
A	496	ARG	-	expression tag	UNP Q9HK01
A	497	PRO	-	expression tag	UNP Q9HK01
A	498	VAL	-	expression tag	UNP Q9HK01
A	499	LEU	-	expression tag	UNP Q9HK01
A	500	GLY	-	expression tag	UNP Q9HK01
A	501	SER	-	expression tag	UNP Q9HK01
A	502	SER	-	expression tag	UNP Q9HK01
A	503	HIS	-	expression tag	UNP Q9HK01
A	504	HIS	-	expression tag	UNP Q9HK01
A	505	HIS	-	expression tag	UNP Q9HK01
A	506	HIS	-	expression tag	UNP Q9HK01
A	507	HIS	-	expression tag	UNP Q9HK01
A	508	HIS	-	expression tag	UNP Q9HK01
B	34	MET	PHE	engineered mutation	UNP Q9HK01
B	405	ASN	SER	engineered mutation	UNP Q9HK01
B	494	SER	-	expression tag	UNP Q9HK01
B	495	GLY	-	expression tag	UNP Q9HK01

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	496	ARG	-	expression tag	UNP Q9HK01
B	497	PRO	-	expression tag	UNP Q9HK01
B	498	VAL	-	expression tag	UNP Q9HK01
B	499	LEU	-	expression tag	UNP Q9HK01
B	500	GLY	-	expression tag	UNP Q9HK01
B	501	SER	-	expression tag	UNP Q9HK01
B	502	SER	-	expression tag	UNP Q9HK01
B	503	HIS	-	expression tag	UNP Q9HK01
B	504	HIS	-	expression tag	UNP Q9HK01
B	505	HIS	-	expression tag	UNP Q9HK01
B	506	HIS	-	expression tag	UNP Q9HK01
B	507	HIS	-	expression tag	UNP Q9HK01
B	508	HIS	-	expression tag	UNP Q9HK01
C	34	MET	PHE	engineered mutation	UNP Q9HK01
C	405	ASN	SER	engineered mutation	UNP Q9HK01
C	494	SER	-	expression tag	UNP Q9HK01
C	495	GLY	-	expression tag	UNP Q9HK01
C	496	ARG	-	expression tag	UNP Q9HK01
C	497	PRO	-	expression tag	UNP Q9HK01
C	498	VAL	-	expression tag	UNP Q9HK01
C	499	LEU	-	expression tag	UNP Q9HK01
C	500	GLY	-	expression tag	UNP Q9HK01
C	501	SER	-	expression tag	UNP Q9HK01
C	502	SER	-	expression tag	UNP Q9HK01
C	503	HIS	-	expression tag	UNP Q9HK01
C	504	HIS	-	expression tag	UNP Q9HK01
C	505	HIS	-	expression tag	UNP Q9HK01
C	506	HIS	-	expression tag	UNP Q9HK01
C	507	HIS	-	expression tag	UNP Q9HK01
C	508	HIS	-	expression tag	UNP Q9HK01
D	34	MET	PHE	engineered mutation	UNP Q9HK01
D	405	ASN	SER	engineered mutation	UNP Q9HK01
D	494	SER	-	expression tag	UNP Q9HK01
D	495	GLY	-	expression tag	UNP Q9HK01
D	496	ARG	-	expression tag	UNP Q9HK01
D	497	PRO	-	expression tag	UNP Q9HK01
D	498	VAL	-	expression tag	UNP Q9HK01
D	499	LEU	-	expression tag	UNP Q9HK01
D	500	GLY	-	expression tag	UNP Q9HK01
D	501	SER	-	expression tag	UNP Q9HK01
D	502	SER	-	expression tag	UNP Q9HK01
D	503	HIS	-	expression tag	UNP Q9HK01

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	504	HIS	-	expression tag	UNP Q9HK01
D	505	HIS	-	expression tag	UNP Q9HK01
D	506	HIS	-	expression tag	UNP Q9HK01
D	507	HIS	-	expression tag	UNP Q9HK01
D	508	HIS	-	expression tag	UNP Q9HK01

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	225	Total O 225 225	0	0
2	B	190	Total O 190 190	0	0
2	C	75	Total O 75 75	0	0
2	D	88	Total O 88 88	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: D-glyceraldehyde dehydrogenase (NADP(+))

Chain A:  94%





- Molecule 1: D-glyceraldehyde dehydrogenase (NADP(+))

Chain B:  93%




- Molecule 1: D-glyceraldehyde dehydrogenase (NADP(+))

Chain C:  3% 93%



- Molecule 1: D-glyceraldehyde dehydrogenase (NADP(+))

Chain D:  3% 92%



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	108.07Å 158.42Å 130.05Å 90.00° 91.57° 90.00°	Depositor
Resolution (Å)	44.63 – 2.10 44.63 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.6 (44.63-2.10) 99.6 (44.63-2.10)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.80 (at 2.10Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.176 , 0.226 0.188 , 0.234	Depositor DCC
$R_{free}$ test set	6331 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.0	Xtrriage
Anisotropy	0.488	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 46.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.019 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	15691	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.63% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.93	0/3897	0.90	5/5281 (0.1%)
1	B	0.88	0/3915	0.84	1/5307 (0.0%)
1	C	0.77	0/3833	0.85	6/5208 (0.1%)
1	D	0.80	0/3832	0.85	7/5204 (0.1%)
All	All	0.85	0/15477	0.86	19/21000 (0.1%)

There are no bond length outliers.

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	247[A]	GLU	CA-C-O	8.34	137.60	120.10
1	C	247[B]	GLU	CA-C-O	8.34	137.60	120.10
1	D	247[A]	GLU	CA-C-N	-7.66	100.34	117.20
1	D	247[B]	GLU	CA-C-N	-7.66	100.34	117.20
1	C	247[A]	GLU	CA-C-N	-7.48	100.75	117.20
1	C	247[B]	GLU	CA-C-N	-7.48	100.75	117.20
1	D	247[A]	GLU	CA-C-O	7.35	135.53	120.10
1	D	247[B]	GLU	CA-C-O	7.35	135.53	120.10
1	D	246	LEU	C-N-CA	6.83	138.78	121.70
1	B	102	ASP	CB-CG-OD1	6.31	123.97	118.30
1	A	247[A]	GLU	CA-C-O	6.11	132.94	120.10
1	A	247[B]	GLU	CA-C-O	6.11	132.94	120.10
1	A	247[A]	GLU	CA-C-N	-5.96	104.10	117.20
1	A	247[B]	GLU	CA-C-N	-5.96	104.10	117.20
1	A	405	ASN	N-CA-CB	-5.83	100.10	110.60
1	D	247[A]	GLU	N-CA-C	5.34	125.41	111.00
1	D	247[B]	GLU	N-CA-C	5.34	125.41	111.00
1	C	247[A]	GLU	N-CA-C	5.33	125.40	111.00
1	C	247[B]	GLU	N-CA-C	5.33	125.40	111.00

There are no chirality outliers.

There are no planarity outliers.



## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3806	0	3713	9	0
1	B	3818	0	3716	9	0
1	C	3745	0	3583	12	0
1	D	3744	0	3599	13	0
2	A	225	0	0	0	0
2	B	190	0	0	0	1
2	C	75	0	0	0	0
2	D	88	0	0	0	0
All	All	15691	0	14611	40	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:188:PHE:O	1:C:193:VAL:HG23	1.97	0.64
1:D:247[B]:GLU:OE1	1:D:457:GLY:HA2	1.99	0.63
1:C:384:ALA:HB1	1:C:385:PRO:HD2	1.86	0.56
1:D:273:LYS:HA	1:D:273:LYS:HE3	1.89	0.55
1:D:142:VAL:HG22	1:D:220:LEU:HB3	1.88	0.55
1:A:444:GLN:HB2	1:B:123:VAL:HG11	1.92	0.52
1:D:70:ARG:NH1	1:D:191:ALA:O	2.43	0.51
1:B:172:LYS:HA	1:B:201:ILE:O	2.12	0.50
1:A:384:ALA:HB1	1:A:385:PRO:HD2	1.95	0.49
1:B:247[A]:GLU:HG3	1:B:458:GLU:OE1	2.14	0.48
1:C:291:GLU:CD	1:C:393:SER:HB3	2.34	0.47
1:D:172:LYS:HA	1:D:201:ILE:O	2.15	0.47
1:A:258:LYS:HB3	1:A:290:HIS:CG	2.50	0.47
1:A:274:TYR:CE1	1:A:365:PRO:HB3	2.50	0.47
1:A:257:TRP:HD1	1:A:259:ASP:OD1	1.98	0.46
1:C:273:LYS:HE3	1:C:273:LYS:HA	1.98	0.46
1:C:390:ARG:NH1	1:C:391:LYS:O	2.49	0.45
1:A:347:GLY:HA3	1:A:365:PRO:O	2.16	0.45
1:C:444:GLN:HB2	1:D:123:VAL:HG11	1.98	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:414:PHE:CE2	1:D:436:ASN:HA	2.52	0.45
1:B:303:GLU:OE2	1:B:306:ARG:NH1	2.51	0.44
1:A:483:HIS:ND1	1:A:489:ASP:OD2	2.43	0.44
1:B:390[A]:ARG:NH1	1:B:391:LYS:O	2.50	0.44
1:B:414:PHE:CE2	1:B:436:ASN:HA	2.52	0.44
1:C:473:ILE:N	1:C:473:ILE:HD12	2.33	0.44
1:C:123:VAL:HG11	1:D:444:GLN:HB2	1.99	0.43
1:A:273:LYS:HA	1:A:273:LYS:HE3	2.00	0.43
1:C:49:ALA:HA	1:C:196:GLY:O	2.19	0.43
1:D:384:ALA:HB1	1:D:385:PRO:HD2	2.00	0.42
1:D:15:SER:HB2	1:D:41:ASP:OD1	2.20	0.42
1:C:462:TYR:O	1:C:466:GLU:HG2	2.20	0.42
1:D:414:PHE:HA	1:D:436:ASN:OD1	2.20	0.42
1:B:56:TRP:CE2	1:B:64:ARG:HD2	2.54	0.41
1:C:302:VAL:HG13	1:C:346:PHE:HB2	2.02	0.41
1:B:142:VAL:HG22	1:B:220:LEU:HB3	2.03	0.41
1:B:310:LEU:HD12	1:B:310:LEU:N	2.35	0.41
1:D:449:GLY:HA3	1:D:453:THR:HG21	2.03	0.41
1:D:134:GLN:HA	1:D:471:LYS:O	2.21	0.41
1:C:151:PRO:HB2	1:C:180:SER:OG	2.21	0.41
1:A:247[B]:GLU:HG3	1:A:458:GLU:OE1	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:646:HOH:O	2:B:646:HOH:O[2_555]	2.16	0.04

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	493/508 (97%)	480 (97%)	13 (3%)	0	100	100
1	B	495/508 (97%)	480 (97%)	15 (3%)	0	100	100
1	C	492/508 (97%)	479 (97%)	13 (3%)	0	100	100
1	D	492/508 (97%)	476 (97%)	16 (3%)	0	100	100
All	All	1972/2032 (97%)	1915 (97%)	57 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	392/419 (94%)	388 (99%)	4 (1%)	76	82
1	B	392/419 (94%)	389 (99%)	3 (1%)	81	86
1	C	375/419 (90%)	373 (100%)	2 (0%)	88	92
1	D	376/419 (90%)	372 (99%)	4 (1%)	73	79
All	All	1535/1676 (92%)	1522 (99%)	13 (1%)	84	86

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	247[A]	GLU
1	A	247[B]	GLU
1	A	273	LYS
1	A	383	PHE
1	B	273	LYS
1	B	383	PHE
1	B	409	LEU
1	C	273	LYS
1	C	383	PHE
1	D	247[A]	GLU
1	D	247[B]	GLU
1	D	273	LYS
1	D	383	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	492/508 (96%)	-0.34	1 (0%) 95 95	26, 37, 53, 69	4 (0%)
1	B	492/508 (96%)	-0.28	1 (0%) 95 95	27, 38, 58, 72	2 (0%)
1	C	492/508 (96%)	0.06	14 (2%) 53 59	41, 53, 69, 83	4 (0%)
1	D	492/508 (96%)	0.01	14 (2%) 53 59	40, 52, 65, 76	4 (0%)
All	All	1968/2032 (96%)	-0.14	30 (1%) 73 77	26, 47, 64, 83	14 (0%)

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	492	PHE	3.4
1	C	341	GLY	3.0
1	C	40	ASP	3.0
1	D	203	GLY	3.0
1	D	294	TYR	2.9
1	C	15	SER	2.8
1	D	470	LEU	2.8
1	A	341	GLY	2.7
1	D	491	LEU	2.7
1	C	12	VAL	2.6
1	D	341	GLY	2.5
1	D	135	TYR	2.5
1	C	211	TYR	2.5
1	D	489	ASP	2.5
1	D	490	ASP	2.5
1	C	48	ALA	2.4
1	C	339	GLU	2.4
1	D	11	TRP	2.4
1	C	206	SER	2.3
1	C	342	ALA	2.3
1	C	11	TRP	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	473	ILE	2.3
1	D	483	HIS	2.2
1	B	356	PRO	2.2
1	C	492	PHE	2.2
1	C	340	SER	2.1
1	C	304	LEU	2.1
1	C	44	ARG	2.1
1	D	15	SER	2.0
1	D	119[A]	ASN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.