



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 20, 2023 – 06:33 PM EDT

PDB ID : 5J7U
Title : Faustovirus major capsid protein
Authors : Klose, T.; Rossmann, M.G.
Deposited on : 2016-04-06
Resolution : 2.44 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






















Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1564 (2.46-2.42)
Clashscore	141614	1631 (2.46-2.42)
Ramachandran outliers	138981	1617 (2.46-2.42)
Sidechain outliers	138945	1617 (2.46-2.42)
RSRZ outliers	127900	1547 (2.46-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	645	 83% 11% 6%
1	B	645	 83% 13% .
1	C	645	 82% 14% .
1	D	645	 82% 12% 6%
1	E	645	 82% 13% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	645	 % 83% 11% 5%
1	G	645	 % 81% 13% 5%
1	H	645	 % 85% 10% 5%
1	I	645	 2% 82% 13% 5%
1	J	645	 % 84% 11% 5%
1	K	645	 % 82% 13% 5%
1	L	645	 % 82% 12% 5%
2	M	21	 100%
2	N	21	 95% 5%
2	O	21	 100%
2	P	21	 95% 5%
2	Q	21	 76% 24%
2	R	21	 71% 19% 10%
2	S	21	 86% 14%
2	T	21	 100%
2	U	21	 95% 5%
2	V	21	 86% 14%
2	W	21	 90% 10%
2	X	21	 67% 24% 10%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 61035 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called major capsid protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	609	Total 4785	C 3060	N 803	O 908	S 14	0	0	0
1	B	616	Total 4837	C 3092	N 812	O 919	S 14	0	0	0
1	C	616	Total 4842	C 3098	N 812	O 918	S 14	0	0	0
1	D	609	Total 4785	C 3060	N 803	O 908	S 14	0	0	0
1	E	616	Total 4841	C 3097	N 812	O 918	S 14	0	0	0
1	F	610	Total 4793	C 3064	N 804	O 911	S 14	0	0	0
1	G	611	Total 4799	C 3067	N 805	O 913	S 14	0	0	0
1	H	615	Total 4831	C 3089	N 811	O 917	S 14	0	0	0
1	I	616	Total 4842	C 3098	N 812	O 918	S 14	0	0	0
1	J	610	Total 4793	C 3064	N 804	O 911	S 14	0	0	0
1	K	614	Total 4823	C 3083	N 810	O 916	S 14	0	0	0
1	L	610	Total 4793	C 3064	N 804	O 911	S 14	0	0	0

- Molecule 2 is a protein called unknown.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	M	21	Total 105	C 63	N 21	O 21	0	0	0
2	N	21	Total 105	C 63	N 21	O 21	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	O	21	Total 105	C 63	N 21	O 21	0	0	0
2	P	21	Total 105	C 63	N 21	O 21	0	0	0
2	Q	21	Total 105	C 63	N 21	O 21	0	0	0
2	R	19	Total 95	C 57	N 19	O 19	0	0	0
2	S	21	Total 105	C 63	N 21	O 21	0	0	0
2	T	21	Total 105	C 63	N 21	O 21	0	0	0
2	U	21	Total 105	C 63	N 21	O 21	0	0	0
2	V	21	Total 105	C 63	N 21	O 21	0	0	0
2	W	21	Total 105	C 63	N 21	O 21	0	0	0
2	X	19	Total 95	C 57	N 19	O 19	0	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	173	Total 173	O 173	0	0
3	B	170	Total 170	O 170	0	0
3	C	164	Total 164	O 164	0	0
3	D	191	Total 191	O 191	0	0
3	E	168	Total 168	O 168	0	0
3	F	172	Total 172	O 172	0	0
3	N	2	Total 2	O 2	0	0
3	O	2	Total 2	O 2	0	0
3	P	5	Total 5	O 5	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	Q	2	Total O 2 2	0	0
3	R	2	Total O 2 2	0	0
3	G	176	Total O 176 176	0	0
3	H	165	Total O 165 165	0	0
3	I	133	Total O 133 133	0	0
3	J	163	Total O 163 163	0	0
3	K	156	Total O 156 156	0	0
3	L	171	Total O 171 171	0	0
3	S	1	Total O 1 1	0	0
3	U	7	Total O 7 7	0	0
3	V	3	Total O 3 3	0	0
3	W	2	Total O 2 2	0	0
3	X	3	Total O 3 3	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: major capsid protein



- Molecule 1: major capsid protein





- Molecule 1: major capsid protein









• Molecule 1: major capsid protein



• Molecule 1: major capsid protein




• Molecule 1: major capsid protein








ILE

- Molecule 1: major capsid protein






TYR
SER
THR
LYS
PHE
TYR
LEU
GLN
CYS
LEU
ILE
LEU
ARG
CYS
ILE

- Molecule 1: major capsid protein





- Molecule 1: major capsid protein





TYR
LEU
GLN
CYS
LEU
LEU
ILE
LEU
ARG
CYS
ILE



• Molecule 1: major capsid protein



• Molecule 1: major capsid protein



• Molecule 1: major capsid protein



• Molecule 2: unknown

Chain M:  100%

There are no outlier residues recorded for this chain.

- Molecule 2: unknown

Chain N:  95% 5%



- Molecule 2: unknown

Chain O:  100%


There are no outlier residues recorded for this chain.


- Molecule 2: unknown

Chain P:  95% 5%




- Molecule 2: unknown

Chain Q:  76% 24%





- Molecule 2: unknown

Chain R:  71% 19% 10%



- Molecule 2: unknown

Chain S:  86% 14%




- Molecule 2: unknown

Chain T:  100%


There are no outlier residues recorded for this chain.

- Molecule 2: unknown

Chain U:  95% 5%





- Molecule 2: unknown

Chain V:  86% 14%




- Molecule 2: unknown

Chain W:  90% 10%



- Molecule 2: unknown

Chain X:  67% 24% 10%



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	86.38Å 390.83Å 130.14Å 90.00° 91.66° 90.00°	Depositor
Resolution (Å)	72.92 – 2.44 72.92 – 2.44	Depositor EDS
% Data completeness (in resolution range)	99.6 (72.92-2.44) 99.8 (72.92-2.44)	Depositor EDS
R_{merge}	0.23	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.97 (at 2.45Å)	Xtrriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.205 , 0.252 0.205 , 0.252	Depositor DCC
R_{free} test set	15720 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	39.2	Xtrriage
Anisotropy	0.133	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	0.039 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	61035	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 46.29 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.1691e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/4901	0.48	0/6697
1	B	0.26	0/4953	0.48	0/6767
1	C	0.27	0/4959	0.49	0/6775
1	D	0.26	0/4901	0.47	0/6697
1	E	0.26	0/4958	0.48	0/6774
1	F	0.26	0/4909	0.48	0/6708
1	G	0.27	0/4915	0.48	0/6716
1	H	0.26	0/4947	0.47	0/6759
1	I	0.26	0/4959	0.47	0/6775
1	J	0.26	0/4909	0.47	0/6708
1	K	0.27	0/4939	0.48	0/6748
1	L	0.26	0/4909	0.48	0/6708
All	All	0.26	0/59159	0.48	0/80832

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4785	0	4711	72	0
1	B	4837	0	4766	61	0
1	C	4842	0	4770	63	0
1	D	4785	0	4711	65	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	4841	0	4768	72	0
1	F	4793	0	4715	57	0
1	G	4799	0	4720	56	0
1	H	4831	0	4761	47	0
1	I	4842	0	4770	59	0
1	J	4793	0	4715	55	0
1	K	4823	0	4750	61	0
1	L	4793	0	4715	57	0
2	M	105	0	24	0	0
2	N	105	0	24	1	0
2	O	105	0	24	0	0
2	P	105	0	24	1	0
2	Q	105	0	24	4	0
2	R	95	0	21	4	0
2	S	105	0	24	3	0
2	T	105	0	24	0	0
2	U	105	0	25	1	0
2	V	105	0	25	3	0
2	W	105	0	24	1	0
2	X	95	0	22	5	0
3	A	173	0	0	3	0
3	B	170	0	0	5	0
3	C	164	0	0	2	0
3	D	191	0	0	4	0
3	E	168	0	0	3	0
3	F	172	0	0	3	0
3	G	176	0	0	5	0
3	H	165	0	0	7	0
3	I	133	0	0	3	0
3	J	163	0	0	4	0
3	K	156	0	0	4	0
3	L	171	0	0	5	0
3	N	2	0	0	1	0
3	O	2	0	0	0	0
3	P	5	0	0	0	0
3	Q	2	0	0	1	0
3	R	2	0	0	2	0
3	S	1	0	0	1	0
3	U	7	0	0	0	0
3	V	3	0	0	0	0
3	W	2	0	0	1	0
3	X	3	0	0	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	61035	0	57157	628	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 628 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:527:ARG:HH12	1:A:529:LYS:CE	1.25	1.44
1:D:62:TYR:CE1	1:E:416:LEU:CD1	2.02	1.43
1:D:62:TYR:CE1	1:E:416:LEU:HD13	1.53	1.40
1:A:527:ARG:NH1	1:A:529:LYS:HE3	1.36	1.36
1:A:351:THR:OG1	1:A:362:LEU:HD22	1.14	1.30

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	607/645 (94%)	593 (98%)	14 (2%)	0	100	100
1	B	614/645 (95%)	597 (97%)	17 (3%)	0	100	100
1	C	614/645 (95%)	598 (97%)	16 (3%)	0	100	100
1	D	607/645 (94%)	591 (97%)	16 (3%)	0	100	100
1	E	614/645 (95%)	597 (97%)	17 (3%)	0	100	100
1	F	608/645 (94%)	583 (96%)	25 (4%)	0	100	100
1	G	609/645 (94%)	592 (97%)	17 (3%)	0	100	100
1	H	613/645 (95%)	593 (97%)	20 (3%)	0	100	100
1	I	614/645 (95%)	596 (97%)	18 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	J	608/645 (94%)	592 (97%)	16 (3%)	0	100	100
1	K	612/645 (95%)	590 (96%)	21 (3%)	1 (0%)	47	57
1	L	608/645 (94%)	588 (97%)	20 (3%)	0	100	100
All	All	7328/7740 (95%)	7110 (97%)	217 (3%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	K	590	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	524/554 (95%)	522 (100%)	2 (0%)	91	94
1	B	530/554 (96%)	530 (100%)	0	100	100
1	C	530/554 (96%)	529 (100%)	1 (0%)	93	96
1	D	524/554 (95%)	523 (100%)	1 (0%)	93	96
1	E	530/554 (96%)	528 (100%)	2 (0%)	91	94
1	F	525/554 (95%)	522 (99%)	3 (1%)	86	91
1	G	526/554 (95%)	524 (100%)	2 (0%)	91	94
1	H	529/554 (96%)	528 (100%)	1 (0%)	93	96
1	I	530/554 (96%)	530 (100%)	0	100	100
1	J	525/554 (95%)	524 (100%)	1 (0%)	93	96
1	K	528/554 (95%)	525 (99%)	3 (1%)	86	91
1	L	525/554 (95%)	522 (99%)	3 (1%)	86	91
All	All	6326/6648 (95%)	6307 (100%)	19 (0%)	92	95

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	K	215	ASP
1	L	492	ASP
1	L	502	PHE
1	L	68	MET
1	F	463	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	L	193	ASN
1	L	318	GLN
1	D	342	GLN
1	F	81	ASN
1	F	318	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	609/645 (94%)	-0.18	1 (0%) 95 95	24, 39, 59, 72	0
1	B	616/645 (95%)	-0.18	3 (0%) 91 91	24, 39, 62, 93	0
1	C	616/645 (95%)	-0.16	9 (1%) 73 71	26, 40, 67, 112	0
1	D	609/645 (94%)	-0.14	4 (0%) 87 87	27, 40, 61, 84	0
1	E	616/645 (95%)	-0.12	11 (1%) 68 64	29, 42, 72, 120	0
1	F	610/645 (94%)	-0.07	7 (1%) 80 79	24, 39, 66, 127	0
1	G	611/645 (94%)	-0.10	3 (0%) 91 91	26, 41, 63, 93	0
1	H	615/645 (95%)	-0.16	5 (0%) 86 85	23, 40, 65, 98	0
1	I	616/645 (95%)	-0.07	13 (2%) 63 60	29, 46, 70, 123	0
1	J	610/645 (94%)	-0.17	5 (0%) 86 85	25, 39, 65, 85	0
1	K	614/645 (95%)	-0.17	6 (0%) 82 81	29, 43, 72, 106	0
1	L	610/645 (94%)	-0.10	3 (0%) 91 91	26, 41, 65, 116	0
2	M	0/21	-	-	-	-
2	N	0/21	-	-	-	-
2	O	0/21	-	-	-	-
2	P	0/21	-	-	-	-
2	Q	0/21	-	-	-	-
2	R	0/21	-	-	-	-
2	S	0/21	-	-	-	-
2	T	0/21	-	-	-	-
2	U	0/21	-	-	-	-
2	V	0/21	-	-	-	-
2	W	0/21	-	-	-	-
2	X	0/21	-	-	-	-
All	All	7352/7992 (91%)	-0.13	70 (0%) 82 81	23, 41, 65, 127	0

The worst 5 of 70 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	I	588	ALA	8.0
1	I	587	ARG	6.1
1	C	587	ARG	6.1
1	H	362	LEU	4.5
1	C	588	ALA	4.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.