



Full wwPDB NMR Structure Validation Report ⓘ

May 29, 2020 – 07:32 am BST

PDB ID : 5JPW
Title : Molecular basis for protein recognition specificity of the DYNLT1/Tctex1 canonical binding groove. Characterization of the interaction with activin receptor IIB
Authors : Rodriguez-Crespo, I.; Merino-Gracia, J.; Bruix, M.; Zamora-Carreras, H.
Deposited on : 2016-05-04

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 72%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	143	
1	B	143	

2 Ensemble composition and analysis i

This entry contains 20 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:13-A:114, A:130-A:137, B:156-B:257, B:273-B:280 (220)	0.92	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 7, 8, 10, 11, 12, 13, 14, 15
2	5, 6, 9, 17, 18, 20
3	4, 16, 19

3 Entry composition i

There is only 1 type of molecule in this entry. The entry contains 4284 atoms, of which 2124 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Dynein light chain Tctex-type 1, Cytoplasmic dynein 1 intermediate chain 2.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	143	2142	678	1062	179	215	8	0
1	B	143	2142	678	1062	179	215	8	0

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP P63172
A	2	SER	-	expression tag	UNP P63172
A	116	GLY	-	linker	UNP P63172
A	117	GLY	-	linker	UNP P63172
A	118	GLY	-	linker	UNP P63172
A	119	SER	-	linker	UNP P63172
A	120	GLY	-	linker	UNP P63172
A	121	GLN	-	linker	UNP P63172
A	122	SER	-	linker	UNP P63172
B	144	GLY	-	expression tag	UNP P63172
B	145	SER	-	expression tag	UNP P63172
B	259	GLY	-	linker	UNP P63172
B	260	GLY	-	linker	UNP P63172
B	261	GLY	-	linker	UNP P63172
B	262	SER	-	linker	UNP P63172
B	263	GLY	-	linker	UNP P63172
B	264	GLN	-	linker	UNP P63172
B	265	SER	-	linker	UNP P63172

4 Residue-property plots [i](#)


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2



- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2




4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1 (medoid)

- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2




- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2




4.2.2 Score per residue for model 2

- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2




- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2




4.2.3 Score per residue for model 3

- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2




- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2




4.2.4 Score per residue for model 4

- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2



- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2






4.2.5 Score per residue for model 5


- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2

Chain A: 71% 6% 23%



- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2

Chain B: 66% 10% 23%



4.2.6 Score per residue for model 6

- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2

Chain A: 69% 8% 23%



- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2


Chain B: 69% 8% 23%




4.2.7 Score per residue for model 7

- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2

Chain A: 65% 12% 23%





- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2


Chain B: 69% 8% 23%



4.2.8 Score per residue for model 8


- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2

Chain A: 61% 15% 23%



- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2


Chain B: 62% 15% 23%



4.2.9 Score per residue for model 9

- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2

Chain A: 68% 9% 23%




- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2

Chain B: 67% 10% 23%




4.2.10 Score per residue for model 10

- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2




- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2




4.2.11 Score per residue for model 11

- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2




- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2



4.2.12 Score per residue for model 12


- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2



V143

- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2

Chain B:

P281
P282
E283
E284
I285
V286

4.2.13 Score per residue for model 13


- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2

Chain A:



- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2

Chain B:

E283
E284
I285
V286

4.2.14 Score per residue for model 14

- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2

Chain A:



- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2

Chain B:



T285
V286

4.2.15 Score per residue for model 15

- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2

Chain A:  65% 11% 23%



P138
P139
R140
E141
I142
V143

- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2

Chain B:  66% 10% 23%



R283
E284
I285
V286

4.2.16 Score per residue for model 16

- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2


Chain A:  62% 15% 23%



M129
V135
P138
P139
R140
E141
I142
V143

- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2


Chain B:  65% 12% 23%




P282
R283
E284
I285
V286

4.2.17 Score per residue for model 17

- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2




- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2




4.2.18 Score per residue for model 18

- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2




- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2




4.2.19 Score per residue for model 19

- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2




- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2




4.2.20 Score per residue for model 20

- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2



- Molecule 1: Dynein light chain Tctex-type 1,Cytoplasmic dynein 1 intermediate chain 2



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *molecular dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure calculation	
Amber	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 6 of this report.

Chemical shift file(s)	input_cs.cif
Number of chemical shift lists	1
Total number of shifts	2748
Number of shifts mapped to atoms	2748
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	72%

No validations of the models with respect to experimental NMR restraints is performed at this time.

COVALENT-GEOMETRY INFOmissingINFO

5.1 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	B	846	836	834	1±2
1	A	846	836	834	1±2
All	All	33840	33440	33360	31

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:85:CYS:SG	1:B:209:ILE:HD11	0.56	2.41	14	4
1:A:66:ILE:HD11	1:B:228:CYS:SG	0.55	2.42	4	2
1:A:79:LEU:HD13	1:B:222:LEU:HD13	0.55	1.77	14	1
1:A:79:LEU:HD11	1:B:220:ALA:HB1	0.52	1.82	14	1
1:A:79:LEU:HD13	1:B:222:LEU:HD11	0.51	1.80	6	1
1:B:224:THR:HB	1:B:278:VAL:HG13	0.51	1.83	12	3
1:B:159:VAL:HA	1:B:162:VAL:HG22	0.51	1.82	12	1
1:A:19:VAL:HG13	1:A:56:LEU:HD22	0.51	1.82	7	1
1:A:77:ALA:HB1	1:B:222:LEU:HD11	0.50	1.82	14	1
1:A:71:ILE:HD12	1:A:106:CYS:CB	0.47	2.39	18	3
1:B:214:ILE:HD12	1:B:249:CYS:HB2	0.46	1.87	8	1
1:A:64:LYS:HE3	1:B:254:PHE:CE2	0.46	2.46	8	1
1:B:162:VAL:HG13	1:B:199:LEU:HD22	0.46	1.87	7	2
1:A:71:ILE:HD12	1:A:106:CYS:HB2	0.45	1.87	8	1
1:A:77:ALA:HB1	1:B:222:LEU:CD1	0.43	2.43	14	1
1:A:79:LEU:CD1	1:B:220:ALA:HB1	0.43	2.42	14	1
1:B:214:ILE:HD12	1:B:249:CYS:CB	0.43	2.44	8	2
1:B:210:VAL:HG12	1:B:253:ALA:HB2	0.42	1.90	5	1
1:A:79:LEU:HD22	1:B:222:LEU:HD22	0.41	1.91	2	1
1:B:157:PHE:CZ	1:B:165:ILE:HD11	0.40	2.51	10	1
1:B:196:LEU:HD13	1:B:199:LEU:HD11	0.40	1.93	11	1

5.2 Torsion angles [i](#)

5.2.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	110/143 (77%)	98±3 (89±2%)	11±3 (10±2%)	1±1 (1±1%)	26	73
1	B	110/143 (77%)	97±3 (89±3%)	12±3 (10±3%)	1±1 (1±1%)	21	69
All	All	4400/5720 (77%)	3910 (89%)	454 (10%)	36 (1%)	24	71

All 19 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	112	GLY	4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	B	255	GLY	4
1	A	76	GLY	3
1	B	221	GLY	3
1	B	219	GLY	3
1	B	207	LYS	3
1	A	38	HIS	2
1	B	203	GLY	2
1	B	181	HIS	2
1	B	182	SER	1
1	A	35	ALA	1
1	A	77	ALA	1
1	A	39	SER	1
1	B	178	ALA	1
1	B	175	GLY	1
1	A	63	PHE	1
1	A	32	GLY	1
1	A	78	GLY	1
1	A	64	LYS	1

5.2.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	95/119 (80%)	82±3 (87±3%)	13±3 (13±3%)	7 47
1	B	95/119 (80%)	83±3 (88±3%)	12±3 (12±3%)	8 50
All	All	3800/4760 (80%)	3310 (87%)	490 (13%)	7 49

All 98 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	B	168	GLU	20
1	A	25	GLU	20
1	B	167	LYS	20
1	A	24	LYS	20
1	A	53	LEU	18
1	A	135	VAL	18

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	B	196	LEU	18
1	B	222	LEU	16
1	A	91	THR	16
1	B	234	THR	16
1	A	79	LEU	12
1	B	245	LYS	11
1	A	102	LYS	11
1	A	72	MET	10
1	B	235	ASP	9
1	A	66	ILE	8
1	B	241	ARG	8
1	A	87	TRP	8
1	A	98	ARG	8
1	A	68	THR	8
1	A	92	ASP	7
1	B	206	PHE	7
1	B	211	THR	7
1	B	256	LEU	7
1	A	137	PHE	6
1	A	80	HIS	6
1	B	157	PHE	6
1	B	215	MET	6
1	B	230	TRP	6
1	B	223	HIS	6
1	A	14	PHE	5
1	B	280	PHE	5
1	A	74	LYS	5
1	B	217	LYS	5
1	A	63	PHE	5
1	B	243	GLU	5
1	A	73	GLN	5
1	B	183	LYS	5
1	A	113	LEU	5
1	B	216	GLN	4
1	A	100	GLU	4
1	A	64	LYS	4
1	A	40	LYS	4
1	B	218	ASN	3
1	A	71	ILE	3
1	A	75	ASN	3
1	B	193	GLU	3
1	A	101	ASN	3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	B	244	ASN	3
1	A	50	GLU	3
1	A	94	SER	3
1	B	160	ASP	3
1	B	207	LYS	2
1	A	65	TYR	2
1	A	21	ASN	2
1	B	257	SER	2
1	B	209	ILE	2
1	A	34	ASN	2
1	B	177	ASN	2
1	A	37	GLN	2
1	B	237	SER	2
1	A	111	PHE	2
1	B	180	GLN	2
1	B	229	PHE	2
1	A	61	LYS	2
1	B	164	ASN	2
1	A	17	ASP	2
1	A	134	GLN	2
1	B	204	LYS	2
1	A	29	SER	1
1	B	274	LYS	1
1	B	161	GLU	1
1	A	107	ILE	1
1	A	88	ASP	1
1	A	89	SER	1
1	A	90	SER	1
1	B	277	GLN	1
1	A	38	HIS	1
1	B	250	ILE	1
1	B	186	GLN	1
1	B	199	LEU	1
1	B	184	VAL	1
1	B	202	LEU	1
1	B	172	SER	1
1	B	247	MET	1
1	A	109	SER	1
1	B	248	TYR	1
1	A	86	PHE	1
1	A	106	CYS	1
1	A	105	TYR	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	28	GLU	1
1	B	171	GLU	1
1	B	232	SER	1
1	A	43	GLN	1
1	B	181	HIS	1
1	B	238	CYS	1
1	B	233	SER	1
1	B	224	THR	1

5.2.3 RNA [i](#)

There are no RNA molecules in this entry.

5.3 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.4 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.5 Ligand geometry [i](#)

There are no ligands in this entry.

5.6 Other polymers [i](#)

There are no such molecules in this entry.

5.7 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 72% for the well-defined parts and 72% for the entire structure.

6.1 Chemical shift list 1

File name: input_cs.cif

Chemical shift list name: *Tctex-IC_dimer.bmrB*

6.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	2748
Number of shifts mapped to atoms	2748
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	8

6.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	280	-0.10 ± 0.06	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	246	-0.08 ± 0.10	None needed (< 0.5 ppm)
$^{13}\text{C}'$	0	—	None (insufficient data)
^{15}N	262	-0.15 ± 0.24	None needed (< 0.5 ppm)

6.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 72%, i.e. 1846 atoms were assigned a chemical shift out of a possible 2578. 32 out of 34 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	854/1096 (78%)	428/438 (98%)	218/440 (50%)	208/218 (95%)
Sidechain	922/1240 (74%)	546/718 (76%)	376/476 (79%)	0/46 (0%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	70/242 (29%)	42/126 (33%)	22/102 (22%)	6/14 (43%)
Overall	1846/2578 (72%)	1016/1282 (79%)	616/1018 (61%)	214/278 (77%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 72%, i.e. 2364 atoms were assigned a chemical shift out of a possible 3288. 36 out of 38 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	1084/1414 (77%)	542/564 (96%)	280/572 (49%)	262/278 (94%)
Sidechain	1198/1616 (74%)	718/942 (76%)	480/616 (78%)	0/58 (0%)
Aromatic	82/258 (32%)	50/134 (37%)	26/110 (24%)	6/14 (43%)
Overall	2364/3288 (72%)	1310/1640 (80%)	786/1298 (61%)	268/350 (77%)

6.1.4 Statistically unusual chemical shifts [i](#)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	B	170	ILE	HG23	-0.80	2.13 – -0.57	-5.9
1	B	170	ILE	HG22	-0.80	2.13 – -0.57	-5.9
1	A	27	ILE	HG21	-0.80	2.13 – -0.57	-5.9
1	A	27	ILE	HG22	-0.80	2.13 – -0.57	-5.9
1	B	170	ILE	HG21	-0.80	2.13 – -0.57	-5.9
1	A	27	ILE	HG23	-0.80	2.13 – -0.57	-5.9
1	B	199	LEU	HB2	-0.23	3.32 – -0.08	-5.4
1	A	56	LEU	HB2	-0.23	3.32 – -0.08	-5.4

6.1.5 Random Coil Index (RCI) plots [i](#)

The images below report *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



Random coil index (RCI) for chain B:

