



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 20, 2023 – 08:02 PM EDT

PDB ID : 5K73  
Title : as-isolated Dbr1 with Fe(II) and Zn(II)  
Authors : Clark, N.E.; Taylor, A.B.; Hart, P.J.  
Deposited on : 2016-05-25  
Resolution : 2.08 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1

# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:  
*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6189 (2.10-2.06)
Clashscore	141614	6738 (2.10-2.06)
Ramachandran outliers	138981	6663 (2.10-2.06)
Sidechain outliers	138945	6664 (2.10-2.06)
RSRZ outliers	127900	6057 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	356	 91% 7%
1	B	356	 91% 7%
1	C	356	 90% 7%
1	D	356	 84% 14%
1	E	356	 90% 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

<b>Mol</b>	<b>Type</b>	<b>Chain</b>	<b>Res</b>	<b>Chirality</b>	<b>Geometry</b>	<b>Clashes</b>	<b>Electron density</b>
4	SO4	A	404	-	-	X	-

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 15014 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA lariat debranching enzyme, putative.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	349	2884	1872	473	523	16	0	3	0
1	B	349	2876	1868	470	523	15	0	2	0
1	C	349	2857	1857	465	520	15	0	0	0
1	D	350	2866	1862	466	523	15	0	0	0
1	E	349	2867	1863	468	521	15	0	1	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP C4M1P9
A	0	ALA	-	expression tag	UNP C4M1P9
B	-1	GLY	-	expression tag	UNP C4M1P9
B	0	ALA	-	expression tag	UNP C4M1P9
C	-1	GLY	-	expression tag	UNP C4M1P9
C	0	ALA	-	expression tag	UNP C4M1P9
D	-1	GLY	-	expression tag	UNP C4M1P9
D	0	ALA	-	expression tag	UNP C4M1P9
E	-1	GLY	-	expression tag	UNP C4M1P9
E	0	ALA	-	expression tag	UNP C4M1P9

- Molecule 2 is FE (II) ION (three-letter code: FE2) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Fe	0	0
			1	1		
2	B	1	Total	Fe	0	0
			1	1		

*Continued on next page...*


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	C	1	Total Fe 1 1	0	0
2	D	1	Total Fe 1 1	0	0
2	E	1	Total Fe 1 1	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Zn 1 1	0	0
3	B	1	Total Zn 1 1	0	0
3	C	1	Total Zn 1 1	0	0
3	D	1	Total Zn 1 1	0	0
3	E	1	Total Zn 1 1	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total 5	O 4	S 1	0	0
4	A	1	Total 5	O 4	S 1	0	0
4	B	1	Total 5	O 4	S 1	0	0
4	B	1	Total 5	O 4	S 1	0	0
4	B	1	Total 5	O 4	S 1	0	0
4	C	1	Total 5	O 4	S 1	0	0
4	C	1	Total 5	O 4	S 1	0	0
4	C	1	Total 5	O 4	S 1	0	0
4	D	1	Total 5	O 4	S 1	0	0
4	D	1	Total 5	O 4	S 1	0	0
4	D	1	Total 5	O 4	S 1	0	0
4	D	1	Total 5	O 4	S 1	0	0
4	E	1	Total 5	O 4	S 1	0	0

- Molecule 5 is HYDROXIDE ION (three-letter code: OH) (formula: HO).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O 1 1	0	0
5	B	1	Total O 1 1	0	0
5	C	1	Total O 1 1	0	0
5	D	1	Total O 1 1	0	0
5	E	1	Total O 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	115	Total O 115 115	0	0
6	B	136	Total O 136 136	0	0
6	C	106	Total O 106 106	0	0
6	D	122	Total O 122 122	0	0
6	E	100	Total O 100 100	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: RNA lariat debranching enzyme, putative




- Molecule 1: RNA lariat debranching enzyme, putative




- Molecule 1: RNA lariat debranching enzyme, putative





- Molecule 1: RNA lariat debranching enzyme, putative







● Molecule 1: RNA lariat debranching enzyme, putative



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.16Å 141.71Å 213.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	69.22 – 2.08 69.22 – 2.08	Depositor EDS
% Data completeness (in resolution range)	86.3 (69.22-2.08) 86.1 (69.22-2.08)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.23 (at 2.08Å)	Xtrriage
Refinement program	PHENIX 1.10_2155	Depositor
R, $R_{free}$	0.178 , 0.213 0.178 , 0.213	Depositor DCC
$R_{free}$ test set	5771 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	41.3	Xtrriage
Anisotropy	0.106	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 46.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	15014	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.82% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FE2, SO4, OH, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.24	0/2963	0.42	0/4004
1	B	0.24	0/2955	0.42	0/3994
1	C	0.24	0/2935	0.42	0/3967
1	D	0.24	0/2944	0.41	0/3979
1	E	0.24	0/2946	0.41	0/3982
All	All	0.24	0/14743	0.42	0/19926

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2884	0	2853	17	0
1	B	2876	0	2844	12	0
1	C	2857	0	2831	16	0
1	D	2866	0	2837	31	0
1	E	2867	0	2837	14	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1	0	0	0	0
2	E	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
4	A	15	0	0	2	0
4	B	15	0	0	0	0
4	C	15	0	0	0	0
4	D	20	0	0	0	0
4	E	5	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	1	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
5	E	1	0	0	0	0
6	A	115	0	0	1	0
6	B	136	0	0	2	0
6	C	106	0	0	1	0
6	D	122	0	0	0	0
6	E	100	0	0	0	0
All	All	15014	0	14202	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 90 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:307:THR:HG22	1:D:310:LEU:H	1.53	0.73
1:E:205:ASP:N	1:E:205:ASP:OD1	2.21	0.73
1:C:193:LYS:NZ	6:C:502:HOH:O	2.30	0.64
1:B:218:LEU:HD21	1:B:242:THR:HG21	1.82	0.62
1:A:201:GLY:O	1:A:204:LYS:NZ	2.33	0.61

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	350/356 (98%)	342 (98%)	8 (2%)	0	100	100
1	B	349/356 (98%)	340 (97%)	8 (2%)	1 (0%)	41	39
1	C	347/356 (98%)	338 (97%)	9 (3%)	0	100	100
1	D	348/356 (98%)	336 (97%)	11 (3%)	1 (0%)	41	39
1	E	348/356 (98%)	341 (98%)	7 (2%)	0	100	100
All	All	1742/1780 (98%)	1697 (97%)	43 (2%)	2 (0%)	51	53

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	13	GLY
1	B	13	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	319/320 (100%)	318 (100%)	1 (0%)	92	95
1	B	318/320 (99%)	314 (99%)	4 (1%)	69	74
1	C	316/320 (99%)	311 (98%)	5 (2%)	62	67
1	D	317/320 (99%)	314 (99%)	3 (1%)	78	83
1	E	317/320 (99%)	314 (99%)	3 (1%)	78	83
All	All	1587/1600 (99%)	1571 (99%)	16 (1%)	76	81

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	205	ASP
1	E	51	TYR
1	C	193	LYS
1	D	307	THR
1	C	63	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	5	GLN
1	E	173	GLN
1	E	352	ASN
1	B	149	ASN
1	A	194	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 29 ligands modelled in this entry, 10 are monoatomic and 5 are modelled with single atom - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	SO4	E	403	-	4,4,4	0.14	0	6,6,6	0.06	0
4	SO4	D	404	-	4,4,4	0.15	0	6,6,6	0.04	0
4	SO4	A	405	-	4,4,4	0.15	0	6,6,6	0.06	0
4	SO4	C	404	-	4,4,4	0.14	0	6,6,6	0.07	0
4	SO4	B	404	-	4,4,4	0.14	0	6,6,6	0.04	0
4	SO4	D	403	-	4,4,4	0.14	0	6,6,6	0.04	0
4	SO4	D	406	-	4,4,4	0.14	0	6,6,6	0.05	0
4	SO4	D	405	-	4,4,4	0.14	0	6,6,6	0.05	0
4	SO4	A	403	-	4,4,4	0.14	0	6,6,6	0.07	0
4	SO4	B	405	-	4,4,4	0.14	0	6,6,6	0.05	0
4	SO4	C	405	-	4,4,4	0.14	0	6,6,6	0.05	0
4	SO4	C	401	-	4,4,4	0.14	0	6,6,6	0.05	0
4	SO4	B	403	-	4,4,4	0.15	0	6,6,6	0.05	0
4	SO4	A	404	-	4,4,4	0.14	0	6,6,6	0.04	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	404	SO4	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	349/356 (98%)	0.35	4 (1%) 80 83	32, 45, 68, 97	0
1	B	349/356 (98%)	0.44	5 (1%) 75 78	32, 43, 68, 98	0
1	C	349/356 (98%)	0.43	8 (2%) 60 64	35, 49, 75, 98	0
1	D	350/356 (98%)	0.51	20 (5%) 23 28	32, 48, 88, 128	0
1	E	349/356 (98%)	0.47	12 (3%) 45 50	31, 49, 82, 113	0
All	All	1746/1780 (98%)	0.44	49 (2%) 53 58	31, 47, 77, 128	0

The worst 5 of 49 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	252	TYR	5.1
1	E	352	ASN	4.6
1	D	200	PRO	4.3
1	E	353	ILE	4.3
1	D	198	PHE	4.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,



median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	OH	B	406	1/1	0.85	0.57	57,57,57,57	0
4	SO4	D	406	5/5	0.86	0.21	101,103,105,106	0
4	SO4	D	404	5/5	0.86	0.15	108,110,111,114	0
4	SO4	A	404	5/5	0.87	0.20	112,115,116,117	0
4	SO4	B	404	5/5	0.89	0.17	105,112,114,115	0
5	OH	C	406	1/1	0.90	0.21	74,74,74,74	0
4	SO4	D	405	5/5	0.91	0.20	113,116,117,120	0
4	SO4	C	401	5/5	0.92	0.23	84,89,92,98	0
4	SO4	A	405	5/5	0.94	0.11	70,71,78,82	0
4	SO4	C	405	5/5	0.96	0.21	70,77,82,90	0
4	SO4	B	405	5/5	0.96	0.18	72,75,80,81	0
5	OH	E	404	1/1	0.96	0.29	50,50,50,50	0
2	FE2	D	401	1/1	0.97	0.10	59,59,59,59	0
3	ZN	C	403	1/1	0.97	0.09	53,53,53,53	0
4	SO4	A	403	5/5	0.97	0.16	62,64,72,75	0
5	OH	A	406	1/1	0.98	0.37	53,53,53,53	0
3	ZN	D	402	1/1	0.98	0.09	59,59,59,59	0
4	SO4	C	404	5/5	0.98	0.19	53,56,65,67	0
4	SO4	E	403	5/5	0.98	0.16	49,51,58,63	0
4	SO4	D	403	5/5	0.99	0.19	44,48,53,55	0
2	FE2	E	401	1/1	0.99	0.12	43,43,43,43	0
3	ZN	A	402	1/1	0.99	0.10	49,49,49,49	0
4	SO4	B	403	5/5	0.99	0.21	48,53,61,64	0
3	ZN	B	402	1/1	0.99	0.12	49,49,49,49	0
2	FE2	B	401	1/1	0.99	0.11	47,47,47,47	0
2	FE2	C	402	1/1	0.99	0.12	49,49,49,49	0
3	ZN	E	402	1/1	0.99	0.11	45,45,45,45	0
5	OH	D	407	1/1	0.99	0.11	47,47,47,47	0
2	FE2	A	401	1/1	0.99	0.10	47,47,47,47	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.