



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 7, 2024 – 03:28 am GMT

PDB ID : 5NXQ  
Title : Crystal structure of the carboxy-terminal domain of yeast Ctf4 bound to a stapled Sld5 CIP  
Authors : Wu, Y.; Pellegrini, L.  
Deposited on : 2017-05-10  
Resolution : 2.41 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4647 (2.44-2.40)
Clashscore	141614	5161 (2.44-2.40)
Ramachandran outliers	138981	5073 (2.44-2.40)
Sidechain outliers	138945	5074 (2.44-2.40)
RSRZ outliers	127900	4543 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	479	 2% 83% 6% 11%
1	B	479	 3% 85% 5% 10%
1	C	479	 6% 58% 38%
2	D	19	 16% 79% 21%
2	E	19	 11% 74% 5% 21%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9931 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase alpha-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	424	Total 3434	C 2205	N 570	O 644	S 15	0	3	0
1	B	431	Total 3482	C 2233	N 579	O 654	S 16	0	2	0
1	C	296	Total 2405	C 1562	N 392	O 440	S 11	0	1	0

There are 66 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	449	MET	-	initiating methionine	UNP Q01454
A	450	GLY	-	expression tag	UNP Q01454
A	451	SER	-	expression tag	UNP Q01454
A	452	SER	-	expression tag	UNP Q01454
A	453	HIS	-	expression tag	UNP Q01454
A	454	HIS	-	expression tag	UNP Q01454
A	455	HIS	-	expression tag	UNP Q01454
A	456	HIS	-	expression tag	UNP Q01454
A	457	HIS	-	expression tag	UNP Q01454
A	458	HIS	-	expression tag	UNP Q01454
A	459	SER	-	expression tag	UNP Q01454
A	460	GLN	-	expression tag	UNP Q01454
A	461	ASP	-	expression tag	UNP Q01454
A	462	PRO	-	expression tag	UNP Q01454
A	463	GLU	-	expression tag	UNP Q01454
A	464	ASN	-	expression tag	UNP Q01454
A	465	LEU	-	expression tag	UNP Q01454
A	466	TYR	-	expression tag	UNP Q01454
A	467	PHE	-	expression tag	UNP Q01454
A	468	GLN	-	expression tag	UNP Q01454
A	469	GLY	-	expression tag	UNP Q01454
A	470	SER	-	expression tag	UNP Q01454
B	449	MET	-	initiating methionine	UNP Q01454

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	450	GLY	-	expression tag	UNP Q01454
B	451	SER	-	expression tag	UNP Q01454
B	452	SER	-	expression tag	UNP Q01454
B	453	HIS	-	expression tag	UNP Q01454
B	454	HIS	-	expression tag	UNP Q01454
B	455	HIS	-	expression tag	UNP Q01454
B	456	HIS	-	expression tag	UNP Q01454
B	457	HIS	-	expression tag	UNP Q01454
B	458	HIS	-	expression tag	UNP Q01454
B	459	SER	-	expression tag	UNP Q01454
B	460	GLN	-	expression tag	UNP Q01454
B	461	ASP	-	expression tag	UNP Q01454
B	462	PRO	-	expression tag	UNP Q01454
B	463	GLU	-	expression tag	UNP Q01454
B	464	ASN	-	expression tag	UNP Q01454
B	465	LEU	-	expression tag	UNP Q01454
B	466	TYR	-	expression tag	UNP Q01454
B	467	PHE	-	expression tag	UNP Q01454
B	468	GLN	-	expression tag	UNP Q01454
B	469	GLY	-	expression tag	UNP Q01454
B	470	SER	-	expression tag	UNP Q01454
C	449	MET	-	initiating methionine	UNP Q01454
C	450	GLY	-	expression tag	UNP Q01454
C	451	SER	-	expression tag	UNP Q01454
C	452	SER	-	expression tag	UNP Q01454
C	453	HIS	-	expression tag	UNP Q01454
C	454	HIS	-	expression tag	UNP Q01454
C	455	HIS	-	expression tag	UNP Q01454
C	456	HIS	-	expression tag	UNP Q01454
C	457	HIS	-	expression tag	UNP Q01454
C	458	HIS	-	expression tag	UNP Q01454
C	459	SER	-	expression tag	UNP Q01454
C	460	GLN	-	expression tag	UNP Q01454
C	461	ASP	-	expression tag	UNP Q01454
C	462	PRO	-	expression tag	UNP Q01454
C	463	GLU	-	expression tag	UNP Q01454
C	464	ASN	-	expression tag	UNP Q01454
C	465	LEU	-	expression tag	UNP Q01454
C	466	TYR	-	expression tag	UNP Q01454
C	467	PHE	-	expression tag	UNP Q01454
C	468	GLN	-	expression tag	UNP Q01454
C	469	GLY	-	expression tag	UNP Q01454

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	470	SER	-	expression tag	UNP Q01454

- Molecule 2 is a protein called MET-ASP-ILE-UA1-ILE-ASP-ASP-ILE-LEU-UA2-GLU-LEU-ASP-LYS-GLU.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
2	D	15	Total	C	N	O	S	0	0	0
			134	84	22	27	1			
2	E	15	Total	C	N	O	S	0	0	0
			134	84	22	27	1			

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	130	Total	O	0	0
			130	130		
4	B	121	Total	O	0	0
			121	121		
4	C	82	Total	O	0	0
			82	82		

Continued on next page...


*Continued from previous page...*


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
4	D	1	Total O 1 1	0	0
4	E	2	Total O 2 2	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase alpha-binding protein

Chain A: 




- Molecule 1: DNA polymerase alpha-binding protein

Chain B: 



- Molecule 1: DNA polymerase alpha-binding protein



Chain C: 





ALA ALA LEU ASN GLY TYR ASP LYS LEU ARG LEU PHE SER CYS SER ASP GLN VAL GLU LYS ALA SER LEU ALA HIS GLU LYS GLN ASP ARG ALA THR ALA VAL LYS ILE SER GLU ARG ALA GLU LEU PRO SER LEU VAL LYS LYS ILE ASN

ASN ILE ARG GLU ALA TYR GLU GLN LEU LYS

- Molecule 2: MET-ASP-ILE-UA1-ILE-ASP-ASP-ILE-LEU-UA2-GLU-LEU-ASP-LYS-GLU



- Molecule 2: MET-ASP-ILE-UA1-ILE-ASP-ASP-ILE-LEU-UA2-GLU-LEU-ASP-LYS-GLU





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	88.68Å 100.29Å 219.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.21 – 2.41 49.21 – 2.41	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.21-2.41) 100.0 (49.21-2.41)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.44 (at 2.42Å)	Xtrriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, $R_{free}$	0.184 , 0.211 0.185 , 0.212	Depositor DCC
$R_{free}$ test set	3806 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	59.7	Xtrriage
Anisotropy	0.310	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 47.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	9931	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	68.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.73% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 9G2, 9FZ, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.24	0/3520	0.43	0/4767
1	B	0.25	0/3569	0.43	0/4832
1	C	0.25	0/2481	0.45	0/3370
2	D	0.19	0/104	0.31	0/135
2	E	0.20	0/104	0.34	0/135
All	All	0.24	0/9778	0.43	0/13239

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3434	0	3372	17	0
1	B	3482	0	3424	15	0
1	C	2405	0	2326	13	0
2	D	134	0	107	0	0
2	E	134	0	107	1	0
3	B	6	0	8	0	0
4	A	130	0	0	1	0
4	B	121	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	82	0	0	2	0
4	D	1	0	0	0	0
4	E	2	0	0	0	0
All	All	9931	0	9344	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 38 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:846:GLU:OE2	1:B:852:ASN:ND2	2.18	0.76
1:A:846:GLU:OE2	1:A:852:ASN:ND2	2.27	0.66
1:A:785:LYS:HE3	1:C:569:ILE:HD12	1.83	0.61
1:C:577:ARG:NH1	4:C:1001:HOH:O	2.34	0.59
1:C:722:LEU:HD21	1:C:774:MET:HE2	1.87	0.57

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	421/479 (88%)	406 (96%)	14 (3%)	1 (0%)	47 61
1	B	429/479 (90%)	416 (97%)	11 (3%)	2 (0%)	29 40
1	C	293/479 (61%)	279 (95%)	13 (4%)	1 (0%)	41 54
2	D	11/19 (58%)	11 (100%)	0	0	100 100
2	E	11/19 (58%)	11 (100%)	0	0	100 100
All	All	1165/1475 (79%)	1123 (96%)	38 (3%)	4 (0%)	41 54

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	508	SER
1	A	749	TYR
1	B	749	TYR
1	B	508	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	379/423 (90%)	378 (100%)	1 (0%)	92	97
1	B	385/423 (91%)	384 (100%)	1 (0%)	92	97
1	C	267/423 (63%)	267 (100%)	0	100	100
2	D	13/16 (81%)	13 (100%)	0	100	100
2	E	13/16 (81%)	13 (100%)	0	100	100
All	All	1057/1301 (81%)	1055 (100%)	2 (0%)	93	97

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	827	TYR
1	B	827	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	559	HIS
1	C	634	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GOL	B	1001	-	5,5,5	0.37	0	5,5,5	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	1001	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	1001	GOL	O1-C1-C2-C3
3	B	1001	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	424/479 (88%)	0.15	10 (2%) 59 56	44, 60, 101, 127	0
1	B	431/479 (89%)	0.17	15 (3%) 44 42	45, 61, 103, 160	0
1	C	296/479 (61%)	0.55	28 (9%) 8 7	49, 66, 124, 161	0
2	D	13/19 (68%)	1.64	3 (23%) 0 0	86, 97, 118, 123	0
2	E	13/19 (68%)	1.55	2 (15%) 2 1	80, 92, 120, 123	0
All	All	1177/1475 (79%)	0.29	58 (4%) 29 27	44, 62, 109, 161	0

The worst 5 of 58 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	668	MET	8.3
1	C	729	TRP	7.7
1	B	666	SER	6.9
1	C	738	THR	6.6
1	C	734	GLY	6.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GOL	B	1001	6/6	0.76	0.24	80,87,94,96	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.