



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 22, 2020 – 01:47 am BST

PDB ID : 5SV0  
Title : Structure of the ExbB/ExbD complex from E. coli at pH 7.0  
Authors : Celia, H.; Botos, I.; Lloubes, R.; Buchanan, S.K.; Noinaj, N.  
Deposited on : 2016-08-04  
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	244	<p>2% 69% 23% • 5%</p>
1	B	244	<p>70% 18% • 8%</p>
1	C	244	<p>70% 17% • 9%</p>
1	D	244	<p>1% 65% 18% 5% 11%</p>
1	E	244	<p>2% 65% 20% • • 11%</p>
1	F	244	<p>3% 71% 18% • 9%</p>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	G	244	 <p>% 70% 20% • 6%</p>
1	H	244	 <p>% 70% 20% • 8%</p>
1	I	244	 <p>64% 24% • 9%</p>
1	J	244	 <p>% 66% 19% • 12%</p>

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 16861 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Biopolymer transport protein ExbB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	225	Total 1692	C 1080	N 295	O 311	S 6	0	0	0
1	C	222	Total 1651	C 1050	N 289	O 306	S 6	0	0	0
1	D	215	Total 1615	C 1028	N 282	O 300	S 5	0	0	0
1	E	216	Total 1609	C 1025	N 279	O 300	S 5	0	0	0
1	F	223	Total 1651	C 1050	N 290	O 306	S 5	0	0	0
1	G	229	Total 1731	C 1104	N 301	O 320	S 6	0	2	0
1	H	225	Total 1700	C 1085	N 296	O 313	S 6	0	0	0
1	I	223	Total 1678	C 1068	N 294	O 310	S 6	0	0	0
1	J	214	Total 1596	C 1017	N 279	O 295	S 5	0	0	0
1	A	233	Total 1749	C 1111	N 308	O 324	S 6	0	0	0

- Molecule 2 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: C<sub>8</sub>H<sub>18</sub>N<sub>2</sub>O<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	D	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
2	F	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
2	I	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		
3	I	1	Total	Ca	0	0
			1	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	14	Total	O	0	0
			14	14		
4	C	11	Total	O	0	0
			11	11		
4	D	12	Total	O	0	0
			12	12		
4	E	16	Total	O	0	0
			16	16		

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	F	16	Total 16	O 16	0	0
4	G	15	Total 15	O 15	0	0
4	H	13	Total 13	O 13	0	0
4	I	15	Total 15	O 15	0	0
4	J	20	Total 20	O 20	0	0
4	A	10	Total 10	O 10	0	0







• Molecule 1: Biopolymer transport protein ExbB




• Molecule 1: Biopolymer transport protein ExbB




• Molecule 1: Biopolymer transport protein ExbB




• Molecule 1: Biopolymer transport protein ExbB








- Molecule 1: Biopolymer transport protein ExbB



- Molecule 1: Biopolymer transport protein ExbB



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	137.17Å 104.78Å 149.36Å 90.00° 91.77° 90.00°	Depositor
Resolution (Å)	46.00 – 2.60 49.76 – 2.58	Depositor EDS
% Data completeness (in resolution range)	94.4 (46.00-2.60) 89.3 (49.76-2.58)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.12	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.08 (at 2.58Å)	Xtrriage
Refinement program	PHENIX (1.10_2142: ???)	Depositor
R, $R_{free}$	0.211 , 0.258 0.210 , 0.258	Depositor DCC
$R_{free}$ test set	2008 reflections (1.59%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	55.7	Xtrriage
Anisotropy	0.185	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 59.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.000 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	16861	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	79.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.96% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, EPE, MLY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.47	0/1764	0.63	0/2392
1	B	0.44	0/1707	0.68	0/2315
1	C	0.44	0/1661	0.61	0/2249
1	D	0.50	0/1625	0.62	0/2200
1	E	0.44	0/1619	0.64	2/2195 (0.1%)
1	F	0.48	0/1662	0.62	1/2253 (0.0%)
1	G	0.46	0/1747	0.66	1/2370 (0.0%)
1	H	0.46	0/1715	0.63	0/2324
1	I	0.48	0/1690	0.64	0/2289
1	J	0.52	0/1605	0.68	0/2174
All	All	0.47	0/16795	0.64	4/22761 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	2
1	D	0	1
1	E	0	1
1	G	0	3
All	All	0	8

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	167	LEU	C-N-CA	6.74	138.55	121.70

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	77	ASP	C-N-CA	6.27	137.37	121.70
1	F	110	ARG	NE-CZ-NH2	-5.74	117.43	120.30
1	G	110	ARG	NE-CZ-NH2	-5.45	117.58	120.30

There are no chirality outliers.

5 of 8 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	17	GLN	Peptide
1	C	18	HIS	Peptide
1	D	162	THR	Peptide
1	E	167	LEU	Peptide
1	G	161	GLN	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1749	0	1759	45	0
1	B	1692	0	1714	40	0
1	C	1651	0	1679	28	0
1	D	1615	0	1651	36	0
1	E	1609	0	1636	40	0
1	F	1651	0	1663	41	0
1	G	1731	0	1734	41	0
1	H	1700	0	1729	30	0
1	I	1678	0	1708	53	0
1	J	1596	0	1630	42	0
2	D	15	0	17	0	0
2	F	15	0	17	1	0
2	I	15	0	17	1	0
3	A	1	0	0	0	0
3	I	1	0	0	0	0
4	A	10	0	0	0	0
4	B	14	0	0	4	0
4	C	11	0	0	0	0
4	D	12	0	0	1	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	E	16	0	0	0	0
4	F	16	0	0	1	0
4	G	15	0	0	1	0
4	H	13	0	0	1	0
4	I	15	0	0	1	0
4	J	20	0	0	0	0
All	All	16861	0	16954	348	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 348 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:117:ARG:NH1	1:C:211:ASP:OD1	1.88	1.06
1:H:136:ILE:O	1:H:140:SER:HB3	1.68	0.93
1:B:90:GLU:HG3	1:B:118:ARG:HH22	1.33	0.92
1:C:196:ASN:HB3	1:C:200:ARG:HH11	1.39	0.87
1:A:141:PRO:HD3	1:A:188:ALA:HB2	1.54	0.87

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	230/244 (94%)	216 (94%)	12 (5%)	2 (1%)	17	35
1	B	222/244 (91%)	215 (97%)	7 (3%)	0	100	100
1	C	219/244 (90%)	217 (99%)	2 (1%)	0	100	100
1	D	212/244 (87%)	206 (97%)	6 (3%)	0	100	100
1	E	213/244 (87%)	206 (97%)	7 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	F	220/244 (90%)	214 (97%)	6 (3%)	0	100	100
1	G	228/244 (93%)	222 (97%)	6 (3%)	0	100	100
1	H	222/244 (91%)	214 (96%)	7 (3%)	1 (0%)	29	52
1	I	220/244 (90%)	215 (98%)	3 (1%)	2 (1%)	17	35
1	J	211/244 (86%)	207 (98%)	4 (2%)	0	100	100
All	All	2197/2440 (90%)	2132 (97%)	60 (3%)	5 (0%)	47	71

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	I	12	VAL
1	A	168	ALA
1	H	68	LEU
1	I	21	ILE
1	A	166	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	176/189 (93%)	161 (92%)	15 (8%)	10	21
1	B	171/189 (90%)	156 (91%)	15 (9%)	10	19
1	C	166/189 (88%)	149 (90%)	17 (10%)	7	14
1	D	164/189 (87%)	147 (90%)	17 (10%)	7	13
1	E	162/189 (86%)	140 (86%)	22 (14%)	3	6
1	F	163/189 (86%)	155 (95%)	8 (5%)	25	48
1	G	173/189 (92%)	159 (92%)	14 (8%)	11	23
1	H	173/189 (92%)	156 (90%)	17 (10%)	8	15
1	I	170/189 (90%)	156 (92%)	14 (8%)	11	22
1	J	160/189 (85%)	142 (89%)	18 (11%)	6	10

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1678/1890 (89%)	1521 (91%)	157 (9%)	8 17

5 of 157 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	22	VAL
1	G	102	ASP
1	A	37	THR
1	F	37	THR
1	G	8	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

10 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	MLY	F	108	1	9,10,11	0.66	0	6,11,13	0.82	0
1	MLY	D	108	1	9,10,11	0.81	0	6,11,13	0.77	0
1	MLY	I	108	1	9,10,11	0.77	0	6,11,13	0.66	0
1	MLY	G	108	1	9,10,11	0.81	0	6,11,13	0.79	0
1	MLY	A	108	1	9,10,11	0.60	0	6,11,13	0.82	0
1	MLY	E	108	1	9,10,11	0.82	0	6,11,13	0.81	0
1	MLY	C	108	1	9,10,11	0.88	0	6,11,13	0.80	0
1	MLY	J	108	1	9,10,11	0.68	0	6,11,13	0.74	0
1	MLY	H	108	1	9,10,11	0.94	0	6,11,13	0.97	0
1	MLY	B	108	1	9,10,11	0.79	0	6,11,13	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MLY	F	108	1	-	1/8/9/11	-
1	MLY	D	108	1	-	2/8/9/11	-
1	MLY	I	108	1	-	0/8/9/11	-
1	MLY	G	108	1	-	0/8/9/11	-
1	MLY	A	108	1	-	0/8/9/11	-
1	MLY	E	108	1	-	1/8/9/11	-
1	MLY	C	108	1	-	1/8/9/11	-
1	MLY	J	108	1	-	0/8/9/11	-
1	MLY	H	108	1	-	1/8/9/11	-
1	MLY	B	108	1	-	1/8/9/11	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	D	108	MLY	CD-CE-NZ-CH2
1	E	108	MLY	CD-CE-NZ-CH2
1	F	108	MLY	CD-CE-NZ-CH2
1	C	108	MLY	CD-CE-NZ-CH2
1	D	108	MLY	CD-CE-NZ-CH1

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and



the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	EPE	D	301	-	15,15,15	0.95	1 (6%)	18,20,20	1.86	7 (38%)
2	EPE	F	301	-	15,15,15	0.93	1 (6%)	18,20,20	2.17	5 (27%)
2	EPE	I	302	-	15,15,15	0.91	1 (6%)	18,20,20	2.03	7 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EPE	D	301	-	-	3/9/19/19	0/1/1/1
2	EPE	F	301	-	-	4/9/19/19	0/1/1/1
2	EPE	I	302	-	-	4/9/19/19	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	301	EPE	C10-S	3.27	1.82	1.77
2	I	302	EPE	C10-S	3.11	1.81	1.77
2	D	301	EPE	C10-S	2.89	1.81	1.77

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	301	EPE	C5-N4-C3	5.67	121.58	108.83
2	I	302	EPE	C7-N4-C5	4.09	121.69	111.23
2	I	302	EPE	C5-N4-C3	4.08	118.01	108.83
2	F	301	EPE	O1S-S-C10	3.93	111.64	106.92
2	D	301	EPE	C5-N4-C3	3.69	117.13	108.83

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	301	EPE	C9-C10-S-O2S
2	D	301	EPE	C9-C10-S-O3S
2	F	301	EPE	C8-C7-N4-C5
2	F	301	EPE	C9-C10-S-O2S
2	F	301	EPE	C9-C10-S-O3S

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	301	EPE	1	0
2	I	302	EPE	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	232/244 (95%)	-0.11	6 (2%) 56 50	42, 74, 134, 191	0
1	B	224/244 (91%)	-0.31	1 (0%) 92 91	46, 76, 121, 164	0
1	C	221/244 (90%)	-0.26	1 (0%) 91 89	42, 77, 115, 160	0
1	D	214/244 (87%)	-0.21	2 (0%) 84 82	41, 68, 128, 200	0
1	E	215/244 (88%)	-0.12	5 (2%) 60 54	42, 76, 134, 164	0
1	F	222/244 (90%)	-0.13	7 (3%) 47 40	43, 73, 147, 218	0
1	G	228/244 (93%)	-0.27	3 (1%) 77 73	44, 76, 132, 214	0
1	H	224/244 (91%)	-0.36	3 (1%) 77 73	46, 75, 116, 197	0
1	I	222/244 (90%)	-0.32	1 (0%) 91 89	45, 77, 123, 155	0
1	J	213/244 (87%)	-0.32	2 (0%) 84 82	40, 70, 142, 181	0
All	All	2215/2440 (90%)	-0.24	31 (1%) 75 71	40, 75, 132, 218	0

The worst 5 of 31 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	13	TRP	5.8
1	J	232	ALA	5.8
1	A	3	ASN	4.8
1	F	12	VAL	4.2
1	E	234	PRO	4.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	MLY	C	108	11/12	0.97	0.14	38,44,60,61	0
1	MLY	B	108	11/12	0.97	0.17	38,43,50,53	0
1	MLY	I	108	11/12	0.98	0.11	33,39,65,67	0
1	MLY	G	108	11/12	0.98	0.12	32,43,52,53	0
1	MLY	E	108	11/12	0.98	0.19	52,56,68,73	0
1	MLY	F	108	11/12	0.98	0.16	44,49,62,66	0
1	MLY	J	108	11/12	0.98	0.13	31,40,59,65	0
1	MLY	D	108	11/12	0.98	0.14	36,40,49,56	0
1	MLY	H	108	11/12	0.99	0.11	35,45,57,60	0
1	MLY	A	108	11/12	0.99	0.13	33,37,55,65	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	EPE	D	301	15/15	0.76	0.15	95,119,159,163	0
3	CA	I	301	1/1	0.81	0.16	104,104,104,104	0
2	EPE	I	302	15/15	0.82	0.14	85,127,167,169	0
2	EPE	F	301	15/15	0.86	0.29	88,114,157,161	0
3	CA	A	301	1/1	0.97	0.16	97,97,97,97	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.