

Full wwPDB EM Validation Report (i)

Nov 27, 2022 – 01:14 AM EST

PDB ID : 5T0G

EMDB ID : EMD-8334

Title: Structural basis for dynamic regulation of the human 26S proteasome

Authors: Chen, S.; Wu, J.; Lu, Y.; Ma, Y.B.; Lee, B.H.; Yu, Z.; Ouyang, Q.; Finley, D.;

Kirschner, M.W.; Mao, Y.

Deposited on : 2016-08-16

Resolution : 4.40 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43

Mogul : 1.8.5 (274361), CSD as541be (2020)

MolProbity : 4.02b-467 buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

MapQ : FAILED

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.31.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 4.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain	
1	G	245	97%	
2	Н	233	100%	•
3	I	260	96%	-
4	J	247	96%	-
5	K	240	93% • 5%	.
6	L	268	88% 11%	ı
7	M	254	94% • 6%	6
8	N	238	80% 20%	-
9	О	276	80% 20%	-
10	Р	204	100%	_



Continued from previous page...

Mol	Chain	Length	Quality of chain	
11	Q	201	99%	
12	R	262	77%	23%
13	S	240	89%	11%
14	Т	263	82%	18%
15	A	433	82%	17%
16	В	440	76%	22%
17	D	418	88%	• 9%
18	E	403	86%	• 12%
19	F	439	81%	17%
20	С	398	94%	
21	U	953	84%	• 15%
22	V	533	88%	• 10%
23	W	456	98%	
24	X	422	90%	10%
25	Y	389	96%	
26	Z	324	87%	• 12%
27	a	376	98%	
28	b	377	50% 49%	
29	С	309	90%	• 7%
30	d	349	72%	26%
31	е	70	56% • 43%	
32	f	749	91%	• 7%



2 Entry composition (i)

There are 34 unique types of molecules in this entry. The entry contains 77800 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Proteasome subunit alpha type-6.

Mol	Chain	Residues		At	oms			AltConf	Trace
1	G	240	Total 1826	C 1160	N 305	O 348	S 13	0	0

• Molecule 2 is a protein called Proteasome subunit alpha type-2.

Mol	Chain	Residues		Ato	oms			AltConf	Trace
2	Н	233	Total 1713	C 1084	N 290	O 334	S 5	0	0

• Molecule 3 is a protein called Proteasome subunit alpha type-4.

Mol	Chain	Residues		Ato	oms	\mathbf{Atoms}				
3	I	250	Total 1912	C 1204	N 329	O 371	S 8	0	0	

• Molecule 4 is a protein called Proteasome subunit alpha type-7.

Mol	Chain	Residues		Ato	oms			AltConf	Trace
4	J	239	Total 1704	C 1056	N 308	O 335	S 5	0	0

• Molecule 5 is a protein called Proteasome subunit alpha type-5.

Mol	Chain	Residues		At	AltConf	Trace			
5	K	228	Total 1722	C 1080	N 284	O 348	S 10	0	0

• Molecule 6 is a protein called Proteasome subunit alpha type-1.

Mol	Chain	Residues		At	oms			AltConf	Trace
6	L	238	Total 1850	C 1159	N 334	O 346	S 11	0	0



• Molecule 7 is a protein called Proteasome subunit alpha type-3.

Mol	Chain	Residues		At	oms			AltConf	Trace
7	M	240	Total 1856	C 1178	N 314	O 353	S 11	0	0

• Molecule 8 is a protein called Proteasome subunit beta type-6.

Mol	Chain	Residues		\mathbf{A}	toms	AltConf	Trace		
8	N	191	Total 1430	C 893	N 245	O 280	S 12	0	0

• Molecule 9 is a protein called Proteasome subunit beta type-7.

Mol	Chain	Residues		At	oms			AltConf	Trace
9	О	220	Total 1643	C 1033	N 280	O 318	S 12	0	0

• Molecule 10 is a protein called Proteasome subunit beta type-3.

Mol	Chain	Residues		At	oms			AltConf	Trace
10	Р	204	Total 1585	C 1010	N 262	O 294	S 19	0	0

• Molecule 11 is a protein called Proteasome subunit beta type-2.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	Q	199	Total 1570	C 1006	N 265	O 290	S 9	0	0

• Molecule 12 is a protein called Proteasome subunit beta type-5.

Mol	Chain	Residues		At	oms			AltConf	Trace
19	R	201	Total	С	N	О	S	0	0
12	10	201	1548	974	273	292	9		U

• Molecule 13 is a protein called Proteasome subunit beta type-1.

Mol	Chain	Residues		At	oms	${f Atoms}$				
13	S	213	Total 1641	C 1036	N 282	O 313	S 10	0	0	

• Molecule 14 is a protein called Proteasome subunit beta type-4.



Mol	Chain	Residues		At	oms			AltConf	Trace
14	Т	215	Total 1667	C 1052	N 285	O 318	S 12	0	0

• Molecule 15 is a protein called 26S protease regulatory subunit 7.

Mo	Chain	Residues		At	oms			AltConf	Trace
15	A	361	Total 2835	C 1788	N 501	O 528	S 18	0	0

• Molecule 16 is a protein called 26S protease regulatory subunit 4.

\mathbf{Mol}	Chain	Residues		At	oms			AltConf	Trace
16	В	341	Total 2662	C 1671	N 453	O 526	S 12	0	0

• Molecule 17 is a protein called 26S protease regulatory subunit 6B.

Mol	Chain	Residues		At	oms			AltConf	Trace
17	D	380	Total 3040	C 1923	N 524	O 580	S 13	0	0

• Molecule 18 is a protein called 26S protease regulatory subunit 10B.

Mol	Chain	Residues		At	oms			AltConf	Trace
18	Е	353	Total 2790	C 1755	N 494	O 525	S 16	0	0

• Molecule 19 is a protein called 26S protease regulatory subunit 6A.

Mol	Chain	Residues		At	oms			AltConf	Trace
19	F	366	Total 2863	C 1802	N 496	O 549	S 16	0	0

• Molecule 20 is a protein called 26S protease regulatory subunit 8.

Mol	Chain	Residues	Atoms					AltConf	Trace
20	С	384	Total 3015	C 1894	N 540	O 564	S 17	0	0

• Molecule 21 is a protein called 26S proteasome non-ATPase regulatory subunit 1.



Mol	Chain	Residues		${f Atoms}$					Trace
21	U	806	Total 6287	C 3990	N 1075	O 1178	S 44	0	0

• Molecule 22 is a protein called 26S proteasome non-ATPase regulatory subunit 3.

Mo	Chain	Residues		At	oms			AltConf	Trace
22	V	480	Total 3852	C 2444	N 684	O 710	S 14	0	0

• Molecule 23 is a protein called 26S proteasome non-ATPase regulatory subunit 12.

Mol	Chain	Residues		At	oms			AltConf	Trace
23	W	456	Total 3703	C 2339	N 635	O 704	S 25	0	0

• Molecule 24 is a protein called 26S proteasome non-ATPase regulatory subunit 11.

Mol	Chain	Residues		At	oms			AltConf	Trace
24	X	380	Total 3009	C 1918	N 509	O 570	S 12	0	0

• Molecule 25 is a protein called 26S proteasome non-ATPase regulatory subunit 6.

Mol	Chain	Residues		At	AltConf	Trace			
25	Y	378	Total 3115	C 1987	N 533	O 578	S 17	0	0

• Molecule 26 is a protein called 26S proteasome non-ATPase regulatory subunit 7.

Mol	Chain	Residues		Ato	AltConf	Trace			
26	Z	286	Total 2281	C 1457	N 392	O 427	S 5	0	0

• Molecule 27 is a protein called 26S proteasome non-ATPase regulatory subunit 13.

Mol	Chain	Residues		At	AltConf	Trace			
27	a	373	Total 2995	C 1911	N 510	O 559	S 15	0	0

• Molecule 28 is a protein called 26S proteasome non-ATPase regulatory subunit 4.



Mol	Chain	Residues		At	oms		AltConf	Trace	
28	h	191	Total	С	N	О	S	0	0
20	D	191	1458	910	261	279	8	0	U

• Molecule 29 is a protein called 26S proteasome non-ATPase regulatory subunit 14.

Mol	Chain	Residues		At	AltConf	Trace			
29	c	287	Total 2260	C 1430	N 389	O 422	S 19	0	0

• Molecule 30 is a protein called 26S proteasome non-ATPase regulatory subunit 8.

Mol	Chain	Residues		At	AltConf	Trace			
30	d	257	Total 2116	C 1371	N 346	O 390	S 9	0	0

• Molecule 31 is a protein called 26S proteasome complex subunit DSS1.


Mol	Chain	Residues		Ato	ms			AltConf	Trace
91		40	Total	С	N	О	S	0	0
31	е	40	334	200	55	77	2	0	U

• Molecule 32 is a protein called 26S proteasome non-ATPase regulatory subunit 2.

Mol	Chain	Residues		\mathbf{A}^{\dagger}	AltConf	Trace			
32	f	694	Total 5331	C 3364	N 899	O 1027	S 41	0	0

• Molecule 33 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).





Mol	Chain	Residues		Ato	oms			AltConf
33	A	1	Total	С	N	О	Р	0
33	A	1	31	10	5	13	3	U
33	В	1	Total	С	N	О	Р	0
00	Ъ	1	31	10	5	13	3	U
33	D	1	Total	С	N	О	Р	0
00	D	1	31	10	5	13	3	U
33	E	1	Total	С	N	Ο	Р	0
00	Ľ	1	31	10	5	13	3	U
33	F	1	Total	С	N	О	Р	0
	I.	1	31	10	5	13	3	O
33	C	1	Total	С	N	Ο	Р	0
55		1	31	10	5	13	3	

 \bullet Molecule 34 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	AltConf
34	c	1	Total Zn 1 1	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Proteasome subunit alpha type-6













• Molecule 19: 26S protease regulatory subunit 6A
















4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	139236	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TECNAI ARCTICA	Depositor
Voltage (kV)	200	Depositor
Electron dose $(e^-/\text{Å}^2)$	30	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

N / L - 1	Cl :-	Во	nd lengths	В	ond angles
Mol	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	G	0.30	0/1859	0.50	0/2523
2	Н	0.30	0/1747	0.49	0/2376
3	I	0.29	0/1942	0.51	0/2628
4	J	0.28	0/1728	0.48	0/2358
5	K	0.28	0/1747	0.48	0/2364
6	L	0.28	0/1885	0.48	0/2552
7	M	0.28	0/1891	0.46	0/2552
8	N	0.28	0/1454	0.48	0/1967
9	О	0.27	0/1670	0.49	0/2265
10	Р	0.28	0/1614	0.47	0/2177
11	Q	0.29	0/1603	0.49	0/2174
12	R	0.28	0/1579	0.45	0/2134
13	S	0.28	0/1671	0.47	0/2253
14	Т	0.28	0/1700	0.46	0/2305
15	A	0.30	0/2886	0.53	0/3899
16	В	0.29	0/2700	0.51	0/3645
17	D	0.31	0/3090	0.59	2/4168 (0.0%)
18	Е	0.30	0/2835	0.52	$1/3821 \; (0.0\%)$
19	F	0.29	0/2903	0.51	1/3912~(0.0%)
20	С	0.29	0/3054	0.56	3/4107 (0.1%)
21	U	0.29	0/6396	0.49	0/8646
22	V	0.31	0/3929	0.57	0/5309
23	W	0.29	0/3751	0.54	3/5042~(0.1%)
24	X	0.27	0/3053	0.44	0/4115
25	Y	0.30	0/3173	0.53	2/4273~(0.0%)
26	Z	0.27	0/2324	0.55	0/3150
27	a	0.36	1/3053~(0.0%)	0.52	0/4133
28	b	0.27	0/1478	0.48	0/2001
29	c	0.33	0/2302	0.60	1/3110 (0.0%)
30	d	0.30	0/2162	0.57	0/2919
31	е	0.28	0/338	0.56	0/450
32	f	0.33	2/5413~(0.0%)	0.63	3/7317 (0.0%)



Mol	Chain	Bo	nd lengths	Bond angles		
MIOI	Mol Chain		# Z > 5	RMSZ	# Z >5	
All	All	0.30	3/78930 (0.0%)	0.52	16/106645 (0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
17	D	0	1
20	С	0	1
29	c	0	1
All	All	0	3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	Ideal(Å)
27	a	145	LEU	C-N	11.92	1.56	1.34
32	f	681	LEU	C-N	6.42	1.46	1.34
32	f	181	MET	C-N	-5.33	1.21	1.34

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
20	С	167	LEU	CB-CG-CD2	6.98	122.86	111.00
32	f	459	GLU	N-CA-C	6.02	127.25	111.00
20	С	217	SER	C-N-CA	5.95	136.58	121.70
17	D	151	ILE	C-N-CA	5.85	136.32	121.70
32	f	107	LEU	CA-CB-CG	5.62	128.22	115.30
25	Y	63	TRP	C-N-CA	5.52	135.51	121.70
23	W	40	LEU	CA-CB-CG	5.50	127.95	115.30
23	W	92	LYS	C-N-CA	5.44	135.31	121.70
32	f	636	GLY	N-CA-C	5.43	126.68	113.10
25	Y	287	LEU	CA-CB-CG	5.42	127.78	115.30
19	F	204	LEU	CA-CB-CG	5.37	127.64	115.30
23	W	135	LYS	C-N-CA	5.33	135.04	121.70
29	c	243	SER	C-N-CA	5.33	135.03	121.70
20	С	217	SER	CA-C-N	5.27	128.79	117.20
17	D	373	ALA	C-N-CA	5.04	134.31	121.70
18	Е	62	LYS	CD-CE-NZ	5.04	123.28	111.70

There are no chirality outliers.



All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
20	С	171	HIS	Peptide
17	D	148	ASP	Peptide
29	c	243	SER	Peptide

5.2 Too-close contacts (i)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	G	238/245~(97%)	223 (94%)	13 (6%)	2 (1%)	19	60
2	Н	230/233 (99%)	216 (94%)	14 (6%)	0	100	100
3	I	$248/260 \ (95\%)$	230 (93%)	17 (7%)	1 (0%)	34	72
4	J	237/247 (96%)	223 (94%)	12 (5%)	2 (1%)	19	60
5	K	224/240 (93%)	204 (91%)	16 (7%)	4 (2%)	8	42
6	L	236/268 (88%)	218 (92%)	18 (8%)	0	100	100
7	M	238/254 (94%)	219 (92%)	18 (8%)	1 (0%)	34	72
8	N	189/238 (79%)	182 (96%)	7 (4%)	0	100	100
9	О	218/276 (79%)	212 (97%)	6 (3%)	0	100	100
10	Р	202/204 (99%)	192 (95%)	10 (5%)	0	100	100
11	Q	197/201 (98%)	185 (94%)	12 (6%)	0	100	100
12	R	199/262 (76%)	191 (96%)	8 (4%)	0	100	100
13	S	211/240 (88%)	203 (96%)	8 (4%)	0	100	100
14	Т	213/263 (81%)	207 (97%)	6 (3%)	0	100	100
15	A	359/433 (83%)	317 (88%)	37 (10%)	5 (1%)	11	47



 $Continued\ from\ previous\ page...$

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
16	В	339/440~(77%)	308 (91%)	27 (8%)	4 (1%)	13	50
17	D	$378/418 \; (90\%)$	339 (90%)	32 (8%)	7 (2%)	8	41
18	E	351/403~(87%)	321 (92%)	27 (8%)	3 (1%)	17	56
19	F	362/439~(82%)	326 (90%)	33 (9%)	3 (1%)	19	60
20	С	382/398~(96%)	337 (88%)	42 (11%)	3 (1%)	19	60
21	U	798/953 (84%)	762 (96%)	34 (4%)	2 (0%)	41	76
22	V	478/533~(90%)	431 (90%)	39 (8%)	8 (2%)	9	43
23	W	$454/456 \; (100\%)$	412 (91%)	38 (8%)	4 (1%)	17	56
24	X	378/422~(90%)	363 (96%)	15 (4%)	0	100	100
25	Y	$376/389 \; (97\%)$	341 (91%)	31 (8%)	4 (1%)	14	52
26	Z	284/324~(88%)	257 (90%)	23 (8%)	4 (1%)	11	47
27	a	371/376 (99%)	343 (92%)	23 (6%)	5 (1%)	12	48
28	b	189/377~(50%)	180 (95%)	8 (4%)	1 (0%)	29	68
29	c	285/309~(92%)	253 (89%)	26 (9%)	6 (2%)	7	39
30	d	255/349 (73%)	227 (89%)	25 (10%)	3 (1%)	13	50
31	e	36/70 (51%)	32 (89%)	3 (8%)	1 (3%)	5	33
32	f	686/749 (92%)	571 (83%)	110 (16%)	5 (1%)	22	62
All	All	9841/11269 (87%)	9025 (92%)	738 (8%)	78 (1%)	24	60

All (78) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	111	VAL
5	K	12	VAL
21	U	364	VAL
23	W	136	ILE
25	Y	350	VAL
29	С	157	ILE
29	С	244	VAL
32	f	62	ILE
32	f	447	VAL
4	J	199	VAL
15	A	159	PRO
15	A	206	ILE
16	В	218	PRO
16	В	263	GLY



Continued from previous page...

Continued from previous page							
Mol	Chain	Res	Type				
17	D	149	SER				
17	D	151	ILE				
18	Е	247	THR				
22	V	82	LEU				
27	a	340	VAL				
32	f	131	VAL				
1	G	185	LYS				
3	I	108	GLU				
17	D	305	VAL				
20	C C V	219	LEU				
20	С	298	ILE				
22	V	59	ALA				
22	V	96	ARG				
25	Y	287	LEU				
26	Z	144	VAL				
27	a	336	VAL				
32	f	281	ILE				
16	В	230	THR				
17	D	170	MET				
17	D	374	ASP				
19	F	168	TYR				
23	W	40	LEU				
23	W	138	VAL				
25	Y	64	GLN				
25	Y	67	VAL				
26	Z	33	LYS				
28	b	77	THR				
29	С	156	VAL				
30	d	213	ARG				
32	f	435	LEU				
5	K	179	SER				
7	M	235	ALA				
15	A	268	LYS				
17	D	155	THR				
18	Е	208	ILE				
19	F	426	GLU				
23	W	427	ASP				
27	a	69	HIS				
27	a	260	ASP				
30	d	214	GLY				
31	е	4	LYS				
18	Е	292	PRO				



Continued from previous page...

Mol	Chain	Res	Type
20	С	68	GLU
22	V	54	LYS
22	V	319	HIS
4	J	98	VAL
15	A	345	LEU
29	c	105	PRO
16	В	227	PRO
17	D	125	LYS
19	F	326	VAL
22	V	464	ILE
26	Z	240	VAL
15	A	172	VAL
21	U	134	VAL
22	V	322	VAL
27	a	166	ILE
29	c	151	VAL
5	K	11	GLY
22	V	317	PRO
5	K	89	ILE
29	С	189	ILE
26	Z	221	PRO
30	d	37	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	G	$193/209\ (92\%)$	192 (100%)	1 (0%)	88 93
2	Н	$164/190\ (86\%)$	164 (100%)	0	100 100
3	I	193/220 (88%)	193 (100%)	0	100 100
4	J	152/210~(72%)	151 (99%)	1 (1%)	84 90
5	K	$186/202\ (92\%)$	186 (100%)	0	100 100
6	L	$198/229\ (86\%)$	197 (100%)	1 (0%)	88 93
7	M	192/211 (91%)	191 (100%)	1 (0%)	88 93



 $Continued\ from\ previous\ page...$

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
8	N	148/180 (82%)	148 (100%)	0	100	100
9	О	177/227 (78%)	177 (100%)	0	100	100
10	Р	172/173 (99%)	172 (100%)	0	100	100
11	Q	164/171 (96%)	164 (100%)	0	100	100
12	R	153/201 (76%)	153 (100%)	0	100	100
13	S	174/198 (88%)	174 (100%)	0	100	100
14	T	175/214 (82%)	175 (100%)	0	100	100
15	A	308/372 (83%)	306 (99%)	2 (1%)	86	92
16	В	298/385 (77%)	296 (99%)	2 (1%)	84	90
17	D	333/366 (91%)	331 (99%)	2 (1%)	86	92
18	E	308/353 (87%)	306 (99%)	2 (1%)	86	92
19	F	312/379 (82%)	307 (98%)	5 (2%)	62	79
20	С	332/346 (96%)	329 (99%)	3 (1%)	78	88
21	U	685/816 (84%)	681 (99%)	4 (1%)	86	92
22	V	414/459 (90%)	409 (99%)	5 (1%)	71	84
23	W	416/416 (100%)	413 (99%)	3 (1%)	84	90
24	X	327/362~(90%)	325 (99%)	2 (1%)	86	92
25	Y	334/344 (97%)	334 (100%)	0	100	100
26	Z	257/295 (87%)	256 (100%)	1 (0%)	91	94
27	a	333/336 (99%)	333 (100%)	0	100	100
28	b	167/312 (54%)	167 (100%)	0	100	100
29	c	252/267 (94%)	250 (99%)	2 (1%)	81	89
30	d	231/293 (79%)	230 (100%)	1 (0%)	91	94
31	е	38/63 (60%)	38 (100%)	0	100	100
32	f	582/628 (93%)	578 (99%)	4 (1%)	84	90
All	All	8368/9627 (87%)	8326 (100%)	42 (0%)	89	93

All (42) residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	G	161	CYS
4	J	21	TYR
6	L	38	LEU



Continued from previous page...

			ous page
Mol	Chain	Res	Type
7	M	238	TYR
15	A	153	LEU
15	A	347	ASP
16	В	178	LYS
16	В	190	LEU
17	D	121	ARG
17	D	143	LEU
18	Е	262	ASN
18	Е	358	ASP
19	F	76	ASN
19	F	134	LEU
19	F	170	SER
19	F	299	GLU
19	F	323	ASN
20	С	50	ASN
20	С	53	ASN
20	C C	175	PHE
21	U	147	TYR
21	U	172	ASP
21	U	345	ASN
21	U	384	GLN
22	V	36	GLU
22	V	240	LEU
22	V	258	TYR
22	V	281	ASN
22	V	324	PHE
23	W	214	PHE
23	W	273	TYR
23	W	361	HIS
24	X	62	GLN
24	X	157	LEU
26	Z	196	HIS
29	c	38	LEU
29	c	234	TYR
30	d	3	GLU
32	f	89	LEU
32	f	107	LEU
32	f	526	THR
32	f	600	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (37) such sidechains are listed below:



Mol	Chain	Res	Type
3	I	119	GLN
4	J	68	ASN
5	K	23	GLN
11	Q	24	ASN
12	R	29	GLN
15	A	94	GLN
15	A	197	HIS
16	В	368	HIS
17	D	83	GLN
17	D	286	GLN
17	D	414	HIS
18	Е	75	ASN
19	F	194	GLN
20	С	296	ASN
21	U	18	GLN
21	U	338	HIS
22	V	33	GLN
23	W	257	GLN
23	W	414	ASN
27	a	244	ASN
29	c	172	HIS
29	С	298	GLN
31	е	6	GLN
32	f	39	HIS
32	f	40	ASN
32	f	212	ASN
32	f	228	GLN
32	f	237	ASN
32	f	269	GLN
32	f	293	ASN
32	f	316	ASN
32	f	372	ASN
32	f	406	ASN
32	f	491	GLN
32	f	565	ASN
32	f	578	ASN
32	f	588	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Chain	Res	Link	Вс	ond leng	ths	В	ond ang	les
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
33	ATP	A	501	-	26,33,33	0.90	1 (3%)	31,52,52	1.84	7 (22%)
33	ATP	D	501	-	26,33,33	0.92	1 (3%)	31,52,52	1.53	5 (16%)
33	ATP	В	501	-	26,33,33	0.91	1 (3%)	31,52,52	1.58	5 (16%)
33	ATP	F	501	-	26,33,33	0.92	1 (3%)	31,52,52	1.60	5 (16%)
33	ATP	С	501	-	26,33,33	0.92	1 (3%)	31,52,52	1.65	6 (19%)
33	ATP	Е	401	-	26,33,33	0.94	1 (3%)	31,52,52	1.85	7 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
33	ATP	A	501	-	-	2/18/38/38	0/3/3/3
33	ATP	D	501	-	-	1/18/38/38	0/3/3/3
33	ATP	В	501	-	-	1/18/38/38	0/3/3/3
33	ATP	F	501	-	-	3/18/38/38	0/3/3/3
33	ATP	С	501	-	-	1/18/38/38	0/3/3/3
33	ATP	E	401	-	-	3/18/38/38	0/3/3/3



All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
33	Ε	401	ATP	C5-C4	2.51	1.47	1.40
33	С	501	ATP	C5-C4	2.43	1.47	1.40
33	D	501	ATP	C5-C4	2.40	1.47	1.40
33	F	501	ATP	C5-C4	2.39	1.47	1.40
33	В	501	ATP	C5-C4	2.38	1.47	1.40
33	A	501	ATP	C5-C4	2.37	1.47	1.40

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
33	A	501	ATP	PA-O3A-PB	-5.34	114.50	132.83
33	Ε	401	ATP	PA-O3A-PB	-4.75	116.52	132.83
33	Ε	401	ATP	PB-O3B-PG	-4.56	117.17	132.83
33	F	501	ATP	PB-O3B-PG	-4.53	117.29	132.83
33	A	501	ATP	PB-O3B-PG	-4.10	118.75	132.83
33	С	501	ATP	PB-O3B-PG	-4.09	118.79	132.83
33	D	501	ATP	PA-O3A-PB	-3.72	120.05	132.83
33	С	501	ATP	PA-O3A-PB	-3.70	120.12	132.83
33	Ε	401	ATP	C3'-C2'-C1'	3.70	106.55	100.98
33	В	501	ATP	PB-O3B-PG	-3.69	120.15	132.83
33	A	501	ATP	N3-C2-N1	-3.49	123.22	128.68
33	Ε	401	ATP	N3-C2-N1	-3.36	123.42	128.68
33	В	501	ATP	N3-C2-N1	-3.33	123.48	128.68
33	A	501	ATP	C3'-C2'-C1'	3.30	105.95	100.98
33	С	501	ATP	N3-C2-N1	-3.25	123.61	128.68
33	D	501	ATP	N3-C2-N1	-3.25	123.61	128.68
33	F	501	ATP	N3-C2-N1	-3.22	123.65	128.68
33	С	501	ATP	C3'-C2'-C1'	3.19	105.78	100.98
33	В	501	ATP	C3'-C2'-C1'	3.13	105.69	100.98
33	В	501	ATP	PA-O3A-PB	-3.00	122.52	132.83
33	F	501	ATP	PA-O3A-PB	-2.97	122.62	132.83
33	D	501	ATP	PB-O3B-PG	-2.75	123.40	132.83
33	D	501	ATP	C3'-C2'-C1'	2.70	105.04	100.98
33	D	501	ATP	C4-C5-N7	-2.66	106.63	109.40
33	F	501	ATP	C3'-C2'-C1'	2.65	104.97	100.98
33	A	501	ATP	C4-C5-N7	-2.52	106.77	109.40
33	В	501	ATP	C4-C5-N7	-2.49	106.81	109.40
33	С	501	ATP	C4-C5-N7	-2.43	106.86	109.40
33	Ε	401	ATP	C4-C5-N7	-2.42	106.88	109.40
33	F	501	ATP	C4-C5-N7	-2.36	106.94	109.40
33	A	501	ATP	C2-N1-C6	2.14	122.42	118.75
33	С	501	ATP	C2-N1-C6	2.09	122.34	118.75



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
33	Ε	401	ATP	C2-N1-C6	2.07	122.30	118.75
33	A	501	ATP	O3G-PG-O2G	2.06	115.50	107.64
33	Е	401	ATP	O3G-PG-O2G	2.03	115.39	107.64

There are no chirality outliers.

All (11) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
33	A	501	ATP	C5'-O5'-PA-O1A
33	Е	401	ATP	C5'-O5'-PA-O3A
33	F	501	ATP	C5'-O5'-PA-O3A
33	F	501	ATP	PA-O3A-PB-O2B
33	Е	401	ATP	C5'-O5'-PA-O1A
33	Е	401	ATP	C5'-O5'-PA-O2A
33	F	501	ATP	C5'-O5'-PA-O1A
33	D	501	ATP	PB-O3A-PA-O2A
33	С	501	ATP	PB-O3A-PA-O1A
33	A	501	ATP	C5'-O5'-PA-O3A
33	В	501	ATP	O4'-C4'-C5'-O5'

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





























5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
32	f	3
2	Н	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	\mid Distance (Å) \mid
1	f	110:ALA	С	111:LEU	N	9.72
1	f	79:ASN	С	80:TYR	N	7.23
1	f	348:ASP	С	349:SER	N	5.88
1	Н	2:ALA	С	3:GLU	N	5.52



6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-8334. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)

This section was not generated.

6.2 Central slices (i)

This section was not generated.

6.3 Largest variance slices (i)

This section was not generated.

6.4 Orthogonal surface views (i)

This section was not generated.

6.5 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)

This section was not generated.

7.2 Volume estimate versus contour level (i)

This section was not generated.

7.3 Rotationally averaged power spectrum (i)

This section was not generated. The rotationally averaged power spectrum had issues being displayed.



8 Fourier-Shell correlation (i)

This section was not generated. No FSC curve or half-maps provided.



9 Map-model fit (i)

This section was not generated.

