

wwPDB EM Validation Summary Report (i)

Nov 27, 2022 – 02:01 AM EST

PDB ID : 5T0H

EMDB ID : EMD-8335

Title: Structural basis for dynamic regulation of the human 26S proteasome

Authors: Chen, S.; Wu, J.; Lu, Y.; Ma, Y.B.; Lee, B.H.; Yu, Z.; Ouyang, Q.; Finley, D.;

Kirschner, M.W.; Mao, Y.

Deposited on : 2016-08-16

Resolution : 6.80 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43

Mogul : 1.8.5 (274361), CSD as541be (2020)

MolProbity : 4.02b-467 buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

MapQ : FAILED

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.31.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 6.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826


The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain	
1	A	433	82%	• 17%
2	В	440	75%	22%
3	С	398	93%	
4	D	418	90%	• 9%
5	Е	403	87%	• 12%
6	F	439	81%	• 17%
7	G	245	98%	
8	Н	233	98%	
9	I	260	96%	
10	J	247	96%	

Continued on next page...



 $Continued\ from\ previous\ page...$





2 Entry composition (i)

There are 34 unique types of molecules in this entry. The entry contains 73509 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called 26S protease regulatory subunit 7.

Mol	Chain	Residues		At	AltConf	Trace			
1	A	361	Total 2835	C 1788	N 501	O 528	S 18	0	0

• Molecule 2 is a protein called 26S protease regulatory subunit 4.

Mol	Chain	Residues		At	AltConf	Trace			
2	В	341	Total 2662	C 1671	N 453	O 526	S 12	0	0

• Molecule 3 is a protein called 26S protease regulatory subunit 8.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	С	384	Total 3015	C 1894	N 540	O 564	S 17	0	0

• Molecule 4 is a protein called 26S protease regulatory subunit 6B.

Mol	Chain	Residues		At	AltConf	Trace			
4	D	380	Total 3040	C 1923	N 524	O 580	S 13	0	0

• Molecule 5 is a protein called 26S protease regulatory subunit 10B.

Mol	Chain	Residues		At	AltConf	Trace			
5	Е	353	Total 2790	C 1755	N 494	O 525	S 16	0	0

• Molecule 6 is a protein called 26S protease regulatory subunit 6A.

Mol	Chain	Residues		At	AltConf	Trace			
6	F	366	Total 2863	C 1802	N 496	O 549	S 16	0	0



• Molecule 7 is a protein called Proteasome subunit alpha type-6.

Mol	Chain	Residues		At	AltConf	Trace			
7	G	240	Total 1826	C 1160	N 305	O 348	S 13	0	0

• Molecule 8 is a protein called Proteasome subunit alpha type-2.

Mol	Chain	Residues	Atoms					AltConf	Trace
0	П	232	Total	С	N	О	S	0	0
0	11	232	1708	1081	289	333	5	0	0

• Molecule 9 is a protein called Proteasome subunit alpha type-4.

\mathbf{Mol}	Chain	Residues		Ato	AltConf	Trace			
9	I	250	Total 1912	C 1204	N 329	O 371	S 8	0	0

• Molecule 10 is a protein called Proteasome subunit alpha type-7.

Mol	Chain	Residues	${f Atoms}$					AltConf	Trace
10	J	239	Total 1704	C 1056	N 308	O 335	S 5	0	0

• Molecule 11 is a protein called Proteasome subunit alpha type-5.

Mol	Chain	Residues		At	AltConf	Trace			
11	К	228	Total 1722	C 1080	N 284	O 348	S 10	0	0

• Molecule 12 is a protein called Proteasome subunit alpha type-1.

Mol	Chain	Residues		At	AltConf	Trace			
12	L	238	Total 1850	C 1159	N 334	O 346	S 11	0	0

• Molecule 13 is a protein called Proteasome subunit alpha type-3.

Mol	Chain	Residues		At	AltConf	Trace			
13	M	240	Total 1856	C 1178	N 314	O 353	S 11	0	0

• Molecule 14 is a protein called Proteasome subunit beta type-6.



Mol	Chain	Residues		\mathbf{A}^{1}	toms	AltConf	Trace		
14	N	191	Total 1430	C 893	N 245	O 280	S 12	0	0

• Molecule 15 is a protein called Proteasome subunit beta type-7.

Mol	Chain	Residues		At	AltConf	Trace			
15	О	220	Total 1643	C 1033	N 280	O 318	S 12	0	0

• Molecule 16 is a protein called Proteasome subunit beta type-3.

Mol	Chain	Residues		At	AltConf	Trace			
16	Р	204	Total 1585	C 1010	N 262	O 294	S 19	0	0

• Molecule 17 is a protein called Proteasome subunit beta type-2.

Mol	Chain	Residues		Ato	oms			AltConf	Trace
17	Q	199	Total 1570	C 1006	N 265	O 290	S 9	0	0

• Molecule 18 is a protein called Proteasome subunit beta type-5.

Mol	Chain	Residues	Atoms					AltConf	Trace
1 Q	B	201	Total	С	N	О	S	0	0
10	10	201	1548	974	273	292	9		

• Molecule 19 is a protein called Proteasome subunit beta type-1.

Mol	Chain	Residues		At	AltConf	Trace			
19	S	213	Total 1641	C 1036	N 282	O 313	S 10	0	0

• Molecule 20 is a protein called Proteasome subunit beta type-4.

Mol	Chain	Residues		At	AltConf	Trace			
20	Т	215	Total 1667	C 1052	N 285	O 318	S 12	0	0

 \bullet Molecule 21 is a protein called 26S proteasome non-ATP ase regulatory subunit 1.



Mol	Chain	Residues		A	AltConf	Trace			
21	U	806	Total 6287	C 3990	N 1075	O 1178	S 44	0	0

• Molecule 22 is a protein called 26S proteasome non-ATPase regulatory subunit 3.

Mol	Chain	Residues		At	AltConf	Trace			
22	V	480	Total 3852	C 2444	N 684	O 710	S 14	0	0

• Molecule 23 is a protein called 26S proteasome non-ATPase regulatory subunit 12.

ľ	Mol	Chain	Residues	Atoms					AltConf	Trace
	23	W	236	Total 1940	C 1237	N 331	O 361	S 11	0	0

• Molecule 24 is a protein called 26S proteasome non-ATPase regulatory subunit 11.

Mol	Chain	Residues	Atoms					AltConf	Trace
24	X	81	Total 647	C 414	N 107	O 124	S 2	0	0

• Molecule 25 is a protein called 26S proteasome non-ATPase regulatory subunit 6.

Mol	Chain	Residues	Atoms					AltConf	Trace
25	Y	378	Total 3115	C 1987	N 533	O 578	S 17	0	0

• Molecule 26 is a protein called 26S proteasome non-ATPase regulatory subunit 7.

Mol	Chain	Residues	Atoms					AltConf	Trace
26	Z	286	Total 2281	C 1457	N 392	O 427	S 5	0	0

• Molecule 27 is a protein called 26S proteasome non-ATPase regulatory subunit 13.

Mol	Chain	Residues	Atoms					AltConf	Trace
27	a	373	Total 2995	C 1911	N 510	O 559	S 15	0	0

• Molecule 28 is a protein called 26S proteasome non-ATPase regulatory subunit 4.



Mol	Chain	Residues	${f Atoms}$					AltConf	Trace
20	h	191	Total	С	N	О	S	0	0
20	D	191	1458	910	261	279	8		0

• Molecule 29 is a protein called 26S proteasome non-ATPase regulatory subunit 14.

Mol	Chain	Residues	Atoms					AltConf	Trace
29	c	287	Total 2260	C 1430	N 389	O 422	S 19	0	0

• Molecule 30 is a protein called 26S proteasome non-ATPase regulatory subunit 8.

Mol	Chain	Residues	Atoms					AltConf	Trace
30	d	257	Total 2116	C 1371	N 346	O 390	S 9	0	0

• Molecule 31 is a protein called 26S proteasome complex subunit DSS1.


Mol	Chain	Residues	Atoms					AltConf	Trace
91		24	Total	С	N	О	S	0	0
31	е	24	197	121	34	40	2	0	U

• Molecule 32 is a protein called 26S proteasome non-ATPase regulatory subunit 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
32	f	694	Total 5331	C 3364	N 899	O 1027	S 41	0	0

• Molecule 33 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).





Mol	Chain	Residues		Ato	oms			AltConf
33	A	1	Total	С	N	О	Р	n
33	A	1	27	10	5	10	2	U
33	В	1	Total	С	N	О	Р	0
00	Ъ	1	27	10	5	10	2	0
33	С	1	Total	С	N	О	Р	0
00		1	27	10	5	10	2	U
33	D	1	Total	С	N	Ο	Р	0
00	D	1	27	10	5	10	2	U
33	E	1	Total	С	N	Ο	Р	0
55	Ľ	1	27	10	5	10	2	U
33	F	1	Total	С	N	Ο	Р	0
55	I.	1	27	10	5	10	2	

 \bullet Molecule 34 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	AltConf
34	c	1	Total Zn 1 1	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 26S protease regulatory subunit 7









• Molecule 12: Proteasome subunit alpha type-1






• Molecule 19: Proteasome subunit beta type-1





• Molecule 24: 26S proteasome non-ATPase regulatory subunit 11









Chain e:

GLY LYS LYS ASP LYS LYS GLU GLU GLU ASP LYS LYS

• Molecule 29: 26S proteasome non-ATPase regulatory subunit 14




 \bullet Molecule 30: 26S proteasome non-ATPase regulatory subunit 8



• Molecule 31: 26S proteasome complex subunit DSS1



• Molecule 32: 26S proteasome non-ATPase regulatory subunit 2





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	18443	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TECNAI ARCTICA	Depositor
Voltage (kV)	200	Depositor
Electron dose $(e^-/\text{Å}^2)$	30	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Clasia.	Во	nd lengths	В	ond angles
Mol	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.27	0/2886	0.54	0/3899
2	В	0.27	0/2700	0.55	0/3645
3	С	0.25	0/3054	0.52	1/4107 (0.0%)
4	D	0.26	0/3090	0.53	0/4168
5	Е	0.25	0/2835	0.48	0/3821
6	F	0.26	0/2903	0.54	0/3912
7	G	0.23	0/1859	0.45	0/2523
8	Н	0.25	0/1743	0.46	0/2372
9	I	0.23	0/1942	0.45	0/2628
10	J	0.23	0/1728	0.44	0/2358
11	K	0.24	0/1747	0.44	0/2364
12	L	0.24	0/1885	0.44	0/2552
13	M	0.23	0/1891	0.42	0/2552
14	N	0.24	0/1454	0.42	0/1967
15	О	0.23	0/1670	0.43	0/2265
16	Р	0.24	0/1614	0.41	0/2177
17	Q	0.23	0/1603	0.41	0/2174
18	R	0.23	0/1579	0.39	0/2134
19	S	0.24	0/1671	0.41	0/2253
20	Т	0.25	0/1700	0.41	0/2305
21	U	0.23	0/6396	0.42	0/8646
22	V	0.25	0/3929	0.52	$2/5309 \ (0.0\%)$
23	W	0.24	0/1975	0.46	0/2659
24	X	0.22	0/655	0.40	0/877
25	Y	0.24	0/3173	0.47	2/4273 (0.0%)
26	Z	1.86	6/2324 (0.3%)	0.53	0/3150
27	a	1.52	2/3052~(0.1%)	0.55	4/4130 (0.1%)
28	b	0.25	0/1478	0.43	0/2001
29	С	0.25	0/2302	0.53	2/3110 (0.1%)
30	d	0.25	0/2162	0.51	0/2919
31	е	0.24	0/198	0.53	0/258
32	f	0.27	1/5413~(0.0%)	0.53	3/7317 (0.0%)



Mol	Chain	Bo	nd lengths	Е	Bond angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z >5
All	All	0.51	9/74611 (0.0%)	0.48	14/100825 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	В	0	2
3	С	0	1
5	Е	0	1
6	F	0	2
21	U	0	3
22	V	0	2
25	Y	0	1
30	d	0	1
32	f	0	2
All	All	0	15

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(\text{\AA})$
27	a	215	GLU	CG-CD	82.57	2.75	1.51
26	Z	228	TYR	CD2-CE2	46.09	2.08	1.39
26	Z	228	TYR	CD1-CE1	45.24	2.07	1.39
26	Z	228	TYR	CE1-CZ	34.72	1.83	1.38
26	Z	228	TYR	CE2-CZ	33.47	1.82	1.38

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
27	a	215	GLU	OE1-CD-OE2	-19.13	100.35	123.30
27	a	215	GLU	CG-CD-OE1	6.35	131.00	118.30
29	c	243	SER	C-N-CA	6.19	137.17	121.70
27	a	215	GLU	CB-CG-CD	6.17	130.86	114.20
32	f	459	GLU	N-CA-C	6.13	127.56	111.00

There are no chirality outliers.

5 of 15 planarity outliers are listed below:



Mol	Chain	Res	Type	Group
2	В	133	VAL	Peptide
2	В	264	PRO	Peptide
3	С	171	HIS	Peptide
5	Е	292	PRO	Peptide
6	F	87	PRO	Peptide

5.2 Too-close contacts (i)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	359/433~(83%)	304 (85%)	53 (15%)	2 (1%)	25 66
2	В	339/440 (77%)	298 (88%)	38 (11%)	3 (1%)	17 57
3	С	382/398 (96%)	317 (83%)	62 (16%)	3 (1%)	19 60
4	D	378/418 (90%)	328 (87%)	49 (13%)	1 (0%)	41 77
5	Е	351/403 (87%)	299 (85%)	51 (14%)	1 (0%)	41 77
6	F	362/439 (82%)	316 (87%)	45 (12%)	1 (0%)	41 77
7	G	238/245 (97%)	221 (93%)	16 (7%)	1 (0%)	34 72
8	Н	230/233 (99%)	210 (91%)	19 (8%)	1 (0%)	34 72
9	I	248/260 (95%)	230 (93%)	18 (7%)	0	100 100
10	J	237/247 (96%)	221 (93%)	15 (6%)	1 (0%)	34 72
11	K	224/240 (93%)	203 (91%)	20 (9%)	1 (0%)	34 72
12	L	236/268 (88%)	222 (94%)	14 (6%)	0	100 100
13	M	238/254 (94%)	218 (92%)	20 (8%)	0	100 100
14	N	189/238 (79%)	175 (93%)	14 (7%)	0	100 100
15	О	218/276 (79%)	209 (96%)	9 (4%)	0	100 100

Continued on next page...



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
16	Р	$202/204\ (99\%)$	188 (93%)	14 (7%)	0	100	100
17	Q	197/201 (98%)	183 (93%)	14 (7%)	0	100	100
18	R	$199/262\ (76\%)$	190 (96%)	9 (4%)	0	100	100
19	S	211/240 (88%)	199 (94%)	12 (6%)	0	100	100
20	Т	213/263 (81%)	204 (96%)	9 (4%)	0	100	100
21	U	798/953 (84%)	729 (91%)	67 (8%)	2 (0%)	41	77
22	V	478/533 (90%)	416 (87%)	60 (13%)	2 (0%)	34	72
23	W	234/456~(51%)	213 (91%)	21 (9%)	0	100	100
24	X	79/422~(19%)	75 (95%)	4 (5%)	0	100	100
25	Y	$376/389 \; (97\%)$	336 (89%)	38 (10%)	2 (0%)	29	69
26	Z	284/324 (88%)	251 (88%)	32 (11%)	1 (0%)	34	72
27	a	369/376~(98%)	340 (92%)	27 (7%)	2 (0%)	29	69
28	b	189/377 (50%)	178 (94%)	11 (6%)	0	100	100
29	С	285/309 (92%)	242 (85%)	39 (14%)	4 (1%)	11	46
30	d	255/349 (73%)	219 (86%)	34 (13%)	2 (1%)	19	60
31	e	20/70 (29%)	16 (80%)	4 (20%)	0	100	100
32	f	686/749 (92%)	574 (84%)	108 (16%)	4 (1%)	25	66
All	All	9304/11269 (83%)	8324 (90%)	946 (10%)	34 (0%)	38	72

5 of 34 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
7	G	111	VAL
11	K	12	VAL
21	U	364	VAL
25	Y	350	VAL
29	c	244	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	308/372~(83%)	305 (99%)	3 (1%)	76	86
2	В	298/385 (77%)	294 (99%)	4 (1%)	69	82
3	С	332/346 (96%)	324 (98%)	8 (2%)	49	69
4	D	333/366 (91%)	329 (99%)	4 (1%)	71	83
5	Е	308/353 (87%)	306 (99%)	2 (1%)	86	92
6	F	312/379 (82%)	303 (97%)	9 (3%)	42	64
7	G	193/209 (92%)	192 (100%)	1 (0%)	88	93
8	Н	164/190 (86%)	162 (99%)	2 (1%)	71	83
9	I	193/220 (88%)	193 (100%)	0	100	100
10	J	152/210 (72%)	151 (99%)	1 (1%)	84	90
11	K	186/202 (92%)	185 (100%)	1 (0%)	88	93
12	L	198/229 (86%)	197 (100%)	1 (0%)	88	93
13	M	192/211 (91%)	191 (100%)	1 (0%)	88	93
14	N	148/180 (82%)	148 (100%)	0	100	100
15	О	177/227 (78%)	177 (100%)	0	100	100
16	Р	172/173 (99%)	172 (100%)	0	100	100
17	Q	164/171 (96%)	164 (100%)	0	100	100
18	R	153/201 (76%)	153 (100%)	0	100	100
19	S	174/198 (88%)	174 (100%)	0	100	100
20	Т	175/214 (82%)	174 (99%)	1 (1%)	86	92
21	U	685/816 (84%)	679 (99%)	6 (1%)	78	87
22	V	414/459 (90%)	408 (99%)	6 (1%)	67	80
23	W	218/416 (52%)	215 (99%)	3 (1%)	67	80
24	X	74/362~(20%)	74 (100%)	0	100	100
25	Y	334/344 (97%)	330 (99%)	4 (1%)	71	83
26	Z	257/295 (87%)	254 (99%)	3 (1%)	71	83
27	a	333/336 (99%)	333 (100%)	0	100	100
28	b	167/312~(54%)	166 (99%)	1 (1%)	86	92
29	c	252/267~(94%)	249 (99%)	3 (1%)	71	83
30	d	231/293 (79%)	229 (99%)	2 (1%)	78	87
31	е	$22/63\ (35\%)$	22 (100%)	0	100	100
32	f	582/628 (93%)	574 (99%)	8 (1%)	67	80

Continued on next page...



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	7901/9627~(82%)	7827 (99%)	74 (1%)	79 87

5 of 74 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
26	Z	133	LEU
32	f	600	LEU
26	Z	191	ILE
30	d	116	HIS
6	F	161	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 61 such sidechains are listed below:

Mol	Chain	Res	Type
7	G	123	GLN
29	С	287	HIS
21	U	685	GLN
29	С	241	ASN
32	f	269	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The



Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	oe Chain Res L		Link	Вс	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
33	ADP	Е	401	-	24,29,29	1.00	2 (8%)	29,45,45	1.47	4 (13%)	
33	ADP	A	501	-	24,29,29	0.99	1 (4%)	29,45,45	1.42	4 (13%)	
33	ADP	С	501	-	24,29,29	0.98	1 (4%)	29,45,45	1.44	4 (13%)	
33	ADP	В	501	-	24,29,29	0.96	1 (4%)	29,45,45	1.35	4 (13%)	
33	ADP	D	501	-	24,29,29	0.93	1 (4%)	29,45,45	1.41	4 (13%)	
33	ADP	F	501	-	24,29,29	1.01	2 (8%)	29,45,45	1.34	4 (13%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
33	ADP	Е	401	-	-	6/12/32/32	0/3/3/3
33	ADP	A	501	-	-	3/12/32/32	0/3/3/3
33	ADP	С	501	-	-	4/12/32/32	0/3/3/3
33	ADP	В	501	-	-	5/12/32/32	0/3/3/3
33	ADP	D	501	-	-	2/12/32/32	0/3/3/3
33	ADP	F	501	-	-	7/12/32/32	0/3/3/3

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	Ideal(A)
33	A	501	ADP	C5-C4	2.63	1.47	1.40
33	С	501	ADP	C5-C4	2.56	1.47	1.40
33	F	501	ADP	C5-C4	2.55	1.47	1.40
33	Е	401	ADP	C5-C4	2.48	1.47	1.40
33	В	501	ADP	C5-C4	2.44	1.47	1.40

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
33	С	501	ADP	PA-O3A-PB	-3.48	120.87	132.83
33	A	501	ADP	PA-O3A-PB	-3.28	121.56	132.83
33	С	501	ADP	C3'-C2'-C1'	3.28	105.91	100.98

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(^{o})$	$Ideal(^{o})$
33	Е	401	ADP	PA-O3A-PB	-3.26	121.63	132.83
33	Е	401	ADP	N3-C2-N1	-3.26	123.59	128.68

There are no chirality outliers.

5 of 27 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
33	A	501	ADP	O4'-C4'-C5'-O5'
33	В	501	ADP	PA-O3A-PB-O3B
33	В	501	ADP	C5'-O5'-PA-O3A
33	С	501	ADP	C5'-O5'-PA-O1A
33	Е	401	ADP	C5'-O5'-PA-O1A


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



















5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks				
32	f	3				
27	a	1				

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	f	110:ALA	С	111:LEU	N	8.67
1	f	79:ASN	С	80:TYR	N	7.26
1	f	348:ASP	С	349:SER	N	6.44
1	a	341:LEU	С	342:ASP	N	5.77



6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-8335. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)

This section was not generated.

6.2 Central slices (i)

This section was not generated.

6.3 Largest variance slices (i)

This section was not generated.

6.4 Orthogonal surface views (i)

This section was not generated.

6.5 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)

This section was not generated.

7.2 Volume estimate versus contour level (i)

This section was not generated.

7.3 Rotationally averaged power spectrum (i)

This section was not generated. The rotationally averaged power spectrum had issues being displayed.



8 Fourier-Shell correlation (i)

This section was not generated. No FSC curve or half-maps provided.



9 Map-model fit (i)

This section was not generated.

