

Full wwPDB X-ray Structure Validation Report (i)

Sep 24, 2023 – 07:47 AM EDT

PDB ID : 5UAI

Title : Crystal structure of Methionyl-tRNA formyltransferase from Pseudomonas

aeruginosa

Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)

Deposited on : 2016-12-19

Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

 $Mol Probity \quad : \quad 4.02b\text{--}467$

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.35.1

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\AA)}) \end{array}$
R_{free}	130704	1235 (2.78-2.74)
Clashscore	141614	1277 (2.78-2.74)
Ramachandran outliers	138981	1257 (2.78-2.74)
Sidechain outliers	138945	1257 (2.78-2.74)
RSRZ outliers	127900	1207 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	322	89%	7%	-
1	В	322	89%	8%	-
1	С	322	89%	7%	
1	D	322	86%	10%	



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 9181 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Methionyl-tRNA formyltransferase.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	310	Total	С	N	О	S	0	0	0
1	A	310	2230	1408	402	411	9	0	U	
1	В	311	Total	С	N	О	S	0	0	0
1	Ъ	311	2252	1420	403	420	9	0	0	
1	С	311	Total	С	N	О	S	0	0	0
1		311	2236	1413	400	414	9	0	U	
1	D	311	Total	С	N	О	S	0	1	0
1	ע	911	2244	1416	403	416	9		1	

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	initiating methionine	UNP O85732
A	-6	ALA	-	expression tag	UNP O85732
A	-5	HIS	-	expression tag	UNP O85732
A	-4	HIS	-	expression tag	UNP O85732
A	-3	HIS	-	expression tag	UNP O85732
A	-2	HIS	-	expression tag	UNP O85732
A	-1	HIS	-	expression tag	UNP O85732
A	0	HIS	-	expression tag	UNP O85732
В	-7	MET	-	initiating methionine	UNP O85732
В	-6	ALA	-	expression tag	UNP O85732
В	-5	HIS	-	expression tag	UNP O85732
В	-4	HIS	-	expression tag	UNP O85732
В	-3	HIS	-	expression tag	UNP O85732
В	-2	HIS	-	expression tag	UNP O85732
В	-1	HIS	-	expression tag	UNP O85732
В	0	HIS	-	expression tag	UNP O85732
С	-7	MET	-	initiating methionine	UNP O85732
С	-6	ALA	-	expression tag	UNP O85732
С	-5	HIS	-	expression tag	UNP O85732
С	-4	HIS	-	expression tag	UNP O85732
С	-3	HIS	-	expression tag	UNP O85732



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
С	-2	HIS	-	expression tag	UNP O85732
С	-1	HIS	-	expression tag	UNP O85732
С	0	HIS	-	expression tag	UNP O85732
D	-7	MET	-	initiating methionine	UNP O85732
D	-6	ALA	-	expression tag	UNP O85732
D	-5	HIS	-	expression tag	UNP O85732
D	-4	HIS	-	expression tag	UNP O85732
D	-3	HIS	-	expression tag	UNP O85732
D	-2	HIS	-	expression tag	UNP O85732
D	-1	HIS	-	expression tag	UNP O85732
D	0	HIS	-	expression tag	UNP O85732

 \bullet Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $\mathrm{C_2H_6O_2}).$



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
2	В	1	Total 4	C 2	O 2	0	0

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	61	Total O 61 61	0	0
3	В	62	Total O 62 62	0	0
3	С	55	Total O 55 55	0	0



Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	37	Total O 37 37	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Methionyl-tRNA formyltransferase





4 Data and refinement statistics (i)

Property	Value	Source	
Space group	P 21 21 2	Depositor	
Cell constants	113.92Å 109.06Å 109.93Å	Donogiton	
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor	
Resolution (Å)	49.08 - 2.75	Depositor	
Resolution (A)	49.08 - 2.75	EDS	
% Data completeness	95.8 (49.08-2.75)	Depositor	
(in resolution range)	95.9 (49.08-2.75)	EDS	
R_{merge}	0.10	Depositor	
R_{sum}	(Not available)	Depositor	
$< I/\sigma(I) > 1$	4.67 (at 2.77Å)	Xtriage	
Refinement program	PHENIX 1.11.1_2575	Depositor	
R, R_{free}	0.167 , 0.234	Depositor	
it, it free	0.167 , 0.234	DCC	
R_{free} test set	2195 reflections (6.31%)	wwPDB-VP	
Wilson B-factor (Å ²)	36.8	Xtriage	
Anisotropy	0.167	Xtriage	
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	$0.33 \; , 51.7$	EDS	
L-test for twinning ²	$< L > = 0.48, < L^2> = 0.31$	Xtriage	
	0.026 for l,-k,h		
	0.020 for -k,-h,-l		
Estimated twinning fraction	0.016 for -h,l,k	Xtriage	
	0.003 for k,l,h		
	0.003 for l,h,k		
F_o, F_c correlation	0.94	EDS	
Total number of atoms	9181	wwPDB-VP	
Average B, all atoms (\mathring{A}^2)	36.0	wwPDB-VP	

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 12.61% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z >5	RMSZ	# Z > 5	
1	A	0.43	0/2273	0.64	0/3104	
1	В	0.43	0/2295	0.63	0/3133	
1	С	0.40	0/2279	0.61	1/3112 (0.0%)	
1	D	0.40	0/2291	0.62	0/3129	
All	All	0.41	0/9138	0.62	1/12478 (0.0%)	

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	С	253	LEU	CB-CG-CD2	-5.13	102.27	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2230	0	2233	12	2
1	В	2252	0	2256	16	2
1	С	2236	0	2238	15	0
1	D	2244	0	2233	21	0
2	В	4	0	6	1	0
3	A	61	0	0	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	В	62	0	0	2	0
3	С	55	0	0	1	0
3	D	37	0	0	1	0
All	All	9181	0	8966	59	2

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 3.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
		${ m distance}({ m \AA})$	overlap (Å)
1:A:36:GLN:HE21	1:A:92:GLY:HA3	1.47	0.80
1:C:26:THR:HG21	1:C:181:ILE:HD12	1.73	0.70
1:A:26:THR:HG21	1:A:181:ILE:HD12	1.74	0.69
1:B:26:THR:HG21	1:B:181:ILE:HD12	1.78	0.65
1:D:36:GLN:HE21	1:D:92:GLY:HA3	1.64	0.61
1:A:163:ASP:OD1	1:A:167:SER:OG	2.15	0.60
1:B:104:ARG:NH2	3:B:502:HOH:O	2.38	0.57
1:C:20:LEU:HD23	1:C:55:LEU:HD23	1.87	0.57
1:C:117:ARG:HG2	1:C:118:TRP:CD1	2.40	0.57
1:B:263:ILE:HD13	1:B:271:LEU:HD11	1.88	0.55
1:C:234:TRP:CD1	1:C:235:PRO:HA	2.43	0.53
1:D:34:TYR:CE1	1:D:64:MET:HG3	2.44	0.52
1:D:137:VAL:HG11	1:D:176:GLY:HA2	1.91	0.52
1:C:24:LEU:HD21	1:C:61:LEU:HD11	1.92	0.52
1:C:5:LEU:HD11	1:C:188:ALA:HB2	1.92	0.51
1:C:39:ARG:HB2	1:D:39:ARG:HD3	1.92	0.50
1:D:36:GLN:O	1:D:65:GLN:NE2	2.42	0.50
1:D:70:ARG:HD2	3:D:423:HOH:O	2.13	0.49
1:A:246:LEU:HD21	1:A:285:LEU:HD22	1.94	0.49
1:B:284:ARG:HD3	3:B:559:HOH:O	2.11	0.49
1:A:88:VAL:O	1:A:109:ASN:ND2	2.46	0.49
1:D:24:LEU:HD23	1:D:30:ILE:HD12	1.95	0.48
1:D:88:VAL:O	1:D:109:ASN:HA	2.13	0.48
1:B:12:THR:HB	1:B:13:PRO:HD3	1.96	0.48
1:D:141:GLN:O	1:D:149:GLY:HA3	2.13	0.48
1:B:206:HIS:HA	2:B:401:EDO:H12	1.97	0.47
1:A:305:GLN:O	1:A:310:GLN:NE2	2.43	0.47
1:A:88:VAL:O	1:A:109:ASN:HA	2.14	0.47
1:C:39:ARG:CB	1:D:39:ARG:HD3	2.45	0.46
1:C:116:PRO:HG2	1:C:201:LEU:HB2	1.98	0.46
		Continu	ed on nert nage



 $Continued\ from\ previous\ page...$

A Land		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)
1:B:88:VAL:O	1:B:109:ASN:HA	2.16	0.46
1:A:39:ARG:HB3	1:B:39:ARG:HD3	1.97	0.46
1:C:39:ARG:HG3	1:D:36:GLN:OE1	2.15	0.46
1:A:36:GLN:OE1	1:B:39:ARG:HG2	2.17	0.45
1:D:10:ALA:HA	1:D:34:TYR:O	2.17	0.45
1:B:24:LEU:HD21	1:B:61:LEU:HD11	1.98	0.45
1:B:34:TYR:CE1	1:B:64:MET:HG3	2.52	0.44
1:C:116:PRO:O	1:C:202:ALA:HA	2.17	0.43
1:C:177:PRO:O	1:C:181:ILE:HG12	2.18	0.43
1:D:12:THR:HB	1:D:13:PRO:HD3	2.01	0.43
1:C:12:THR:HB	1:C:13:PRO:HD3	2.01	0.43
1:D:234:TRP:CD1	1:D:235:PRO:HA	2.55	0.42
1:C:171:ARG:HD3	3:C:421:HOH:O	2.18	0.42
1:D:247:LYS:HB2	1:D:286:GLN:HB3	2.01	0.42
1:B:187:LEU:HD23	1:B:187:LEU:HA	1.92	0.42
1:B:34:TYR:HA	1:B:64:MET:O	2.20	0.41
1:D:253:LEU:HD12	1:D:253:LEU:HA	1.86	0.41
1:D:117:ARG:HG2	1:D:118:TRP:CD1	2.54	0.41
1:A:234:TRP:CD1	1:A:235:PRO:HA	2.54	0.41
1:A:215:LEU:HD23	1:A:215:LEU:HA	1.92	0.41
1:D:158:PRO:O	1:D:171:ARG:NH1	2.54	0.41
1:B:163:ASP:OD1	1:B:167:SER:OG	2.31	0.41
1:C:102:ILE:HB	1:C:103:PRO:HD3	2.02	0.41
1:A:215:LEU:HD22	1:A:224:LEU:HD22	2.02	0.41
1:D:23:LEU:HD23	1:D:23:LEU:HA	1.80	0.41
1:D:113:SER:HB3	1:D:123:PRO:HB2	2.02	0.41
1:B:177:PRO:O	1:B:181:ILE:HG12	2.20	0.41
1:B:140:MET:CE	1:B:147:ASP:HA	2.52	0.40
1:D:241:LEU:HB2	1:D:312:LEU:HD22	2.03	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:A:202:ALA:O	1:B:226:ARG:NH2[3_444]	2.01	0.19
1:A:226:ARG:NH2	1:B:202:ALA:O[3_444]	2.05	0.15



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Chain Analysed Favoured Allowed		Outliers	Perce	$_{ m ntiles}$	
1	A	308/322 (96%)	298 (97%)	8 (3%)	2 (1%)	25	42
1	В	309/322~(96%)	299 (97%)	8 (3%)	2 (1%)	25	42
1	С	309/322~(96%)	303 (98%)	4 (1%)	2 (1%)	25	42
1	D	310/322 (96%)	303 (98%)	4 (1%)	3 (1%)	15	27
All	All	$1236/1288 \; (96\%)$	1203 (97%)	24 (2%)	9 (1%)	22	39

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	303	ARG
1	A	236	VAL
1	В	12	THR
1	С	12	THR
1	С	236	VAL
1	A	12	THR
1	В	236	VAL
1	D	12	THR
1	D	236	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	in Analysed Rotameric		Outliers	Percentiles
1	A	217/242 (90%)	215 (99%)	2 (1%)	78 87
1	В	222/242 (92%)	221 (100%)	1 (0%)	88 92



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Rotameric Outliers	
1	C	218/242 (90%)	215 (99%)	3 (1%)	67 79
1	D	218/242 (90%)	217 (100%)	1 (0%)	88 92
All	All	875/968 (90%)	868 (99%)	7 (1%)	81 88

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	140	MET
1	A	281	ARG
1	В	287	LEU
1	С	39	ARG
1	С	64	MET
1	С	140	MET
1	D	64	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	206	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and



the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	B	ond leng	$_{ m gths}$	В	ond ang	gles
WIOI	туре	Chain	nes	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	В	401	-	3,3,3	0.50	0	2,2,2	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	В	401	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	В	401	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	401	EDO	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q<0.9
1	A	310/322 (96%)	-0.31	5 (1%) 72 79	15, 31, 62, 92	0
1	В	311/322 (96%)	-0.30	0 100 100	15, 31, 55, 74	0
1	С	311/322 (96%)	-0.17	2 (0%) 89 92	18, 38, 61, 77	0
1	D	311/322 (96%)	-0.25	3 (0%) 82 87	18, 35, 69, 94	0
All	All	1243/1288 (96%)	-0.26	10 (0%) 86 90	15, 33, 63, 94	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	299	TYR	3.3
1	A	296	ALA	3.0
1	D	299	TYR	3.0
1	D	306	PHE	2.6
1	A	302	ARG	2.5
1	A	4	ALA	2.3
1	D	302	ARG	2.3
1	A	295	PHE	2.3
1	С	27	PRO	2.0
1	С	24	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	EDO	В	401	4/4	0.89	0.22	31,32,35,45	0

6.5 Other polymers (i)

There are no such residues in this entry.

