

wwPDB NMR Structure Validation Summary Report (i)

Jun 3, 2023 – 05:21 AM EDT

PDB ID : 5URN BMRB ID : 30243

Title: NMR structure of the complex between the PH domain of the Tfb1 subunit

from TFIIH and the transactivation domain 1 of p65

Authors: Lecoq, L.; Omichinski, J.G.; Raiola, L.; Cyr, N.; Chabot, P.; Arseneault, G.;

Legault, P.

Deposited on : 2017-02-11

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

wwPDB-RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

 $\begin{array}{ccc} wwPDB\text{-}ShiftChecker &: & v1.2 \\ BMRB \ Restraints \ Analysis &: & v1.2 \\ \end{array}$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.33

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 94%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	NMR archive	
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries})$	
Clashscore	158937	12864	
Ramachandran outliers	154571	11451	
Sidechain outliers	154315	11428	

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain				
1	A	115	76%	15%	10%		
2	В	33	100%				



2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 5 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues					
Well-defined core Residue range (total) Backbone RMSD (Å) Medoid model					
1	A:4-A:65, A:87-A:112 (88)	0.18	5		
2	A:68-A:83 (16)	0.34	4		

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters. No single-model clusters were found.

Cluster number	Models
1	3, 5, 6, 8, 9, 11, 12, 14, 15, 17, 18
2	2, 10, 13, 16, 19, 20
3	1, 4, 7



3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 2274 atoms, of which 1135 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called RNA polymerase II transcription factor B subunit 1.

Mol	Chain	Residues		Atoms			Trace		
1	Λ	115	Total	С	Н	N	О	S	0
1	A	110	1821	559	919	161	177	5	U

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	PRO	-	expression tag	UNP P32776

• Molecule 2 is a protein called Transcription factor p65.

Mol	Chain	Residues	Atoms			Trace			
2	D	33	Total	С	Н	N	О	S	0
	Б	ეე	453	146	216	35	55	1	U

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	519	GLY	-	expression tag	UNP Q04206
В	520	SER	-	expression tag	UNP Q04206
В	523	TYR	LEU	engineered mutation	UNP Q04206




4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


• Molecule 1: RNA polymerase II transcription factor B subunit 1



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 5. Colouring as in section 4.1 above.

• Molecule 1: RNA polymerase II transcription factor B subunit 1





Refinement protocol and experimental data overview (i) 5



The models were refined using the following method: *simulated annealing*.

Of the 260 calculated structures, 20 were deposited, based on the following criterion: structures with the lowest energy.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	refinement	
YASARA	refinement	
ARIA	structure solution	
CCPNMR	structure solution	
VNMR	structure solution	
NMRPipe	structure solution	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1812
Number of shifts mapped to atoms	1812
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	94%



6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0 ± 0.0	1.0 ± 0.0
All	All	0	20

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

All unique planar outliers are listed below.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	23	SER	Peptide	20

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	815	843	841	12±4
2	В	0	0	0	0±0
All	All	16300	16860	16820	249

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

5 of 99 unique clashes are listed below, sorted by their clash magnitude.

Atom 1	Atom-1 Atom-2		$\operatorname{Distance}(\operatorname{\AA})$	Models	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:74:ASP:CB	1:A:76:GLU:OE2	0.98	2.10	11	1



Continued from previous page...

Atom-1	Atom 2	Clash(Å)	$Distance(\mathring{A})$	${f Models}$	
Atom-1	Atom-2 Clash(Distance(A)	Worst	Total
1:A:74:ASP:HB2	1:A:76:GLU:OE2	0.84	1.73	11	1
1:A:74:ASP:HB3	1:A:76:GLU:OE2	0.83	1.73	11	1
1:A:53:ALA:HB2	1:A:94:ARG:HD2	0.81	1.52	8	3
1:A:51:THR:HG23	1:A:59:MET:HG2	0.74	1.57	11	2

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	104/115 (90%)	99±1 (95±1%)	5±1 (5±1%)	0±0 (0±0%)	100 100
2	В	0	-	-	-	-
All	All	2080/2960 (70%)	1983 (95%)	97 (5%)	0 (0%)	100 100

There are no Ramachandran outliers.

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Analysed Rotameric Out		Percentiles
1	A	93/103 (90%)	90±1 (97±1%)	3±1 (3±1%)	41 87
2	В	0	-	-	-
All	All	1860/2600 (72%)	1798 (97%)	62 (3%)	41 87

5 of 18 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	48	LEU	16
1	A	16	ILE	15



Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	26	GLU	6
1	A	90	SER	3
1	A	32	THR	3

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 94% for the well-defined parts and 93% for the entire structure.

7.1 Chemical shift list 1

File name: working cs.cif

Chemical shift list name: CS_Tfb1_p65_all.str

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1812
Number of shifts mapped to atoms	1812
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	4

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\mathrm{C}_{\alpha}$	148	-0.03 ± 0.07	None needed ($< 0.5 \text{ ppm}$)
$^{13}C_{\beta}$	139	0.06 ± 0.15	None needed (< 0.5 ppm)
¹³ C′	136	0.03 ± 0.08	None needed ($< 0.5 \text{ ppm}$)
^{15}N	137	-0.28 ± 0.42	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 94%, i.e. 1347 atoms were assigned a chemical shift out of a possible 1433. 0 out of 14 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathrm{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Backbone	511/519 (98%)	209/210 (100%)	202/208 (97%)	100/101 (99%)
Sidechain	778/847 (92%)	523/549 (95%)	244/263 (93%)	11/35 (31%)



Continued from previous page...

	Total	$^{1}{ m H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Aromatic	58/67 (87%)	29/33~(88%)	28/29 (97%)	1/5 (20%)
Overall	1347/1433 (94%)	761/792 (96%)	474/500 (95%)	112/141 (79%)

7.1.4 Statistically unusual chemical shifts (i)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	44	THR	HG1	4.96	0.08 - 2.19	18.1
1	A	10	GLU	HG2	0.69	1.24 - 3.30	-7.6
1	A	9	PHE	HB2	0.31	1.20 - 4.80	-7.5
1	A	10	GLU	HG3	0.91	1.20 - 3.30	-6.4

7.1.5 Random Coil Index (RCI) plots (i)


The image below reports random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



Random coil index (RCI) for chain B:







8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	3460
Intra-residue ($ i-j =0$)	1470
Sequential (i-j =1)	630
Medium range ($ i-j >1$ and $ i-j <5$)	402
Long range (i-j ≥5)	912
Inter-chain	46
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	211
Number of unmapped restraints	0
Number of restraints per residue	24.8
Number of long range restraints per residue ¹	6.2

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	104.5	0.2
0.2-0.5 (Medium)	180.1	0.5
>0.5 (Large)	316.9	6.62



8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation.

Bins $(^{\circ})$	Average number of violations per model	\mathbf{Max} (°)
1.0-10.0 (Small)	15.6	10.0
10.0-20.0 (Medium)	2.6	12.9
>20.0 (Large)	None	None



9 Distance violation analysis (i)

9.1 Summary of distance violations (i)


The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Destroints tune	Count	% ¹	Vi	${f Violated^3}$			tently	$\mathbf{Violated}^4$
Restraints type	Count	701	Count	$\%^2$	$\%^{1}$	Count	$\frac{1}{2}$	$\%^1$
Intra-residue (i-j =0)	1470	42.5	281	19.1	8.1	54	3.7	1.6
Backbone-Backbone	131	3.8	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	770	22.3	184	23.9	5.3	33	4.3	1.0
Sidechain-Sidechain	569	16.4	97	17.0	2.8	21	3.7	0.6
Sequential (i-j =1)	630	18.2	148	23.5	4.3	39	6.2	1.1
Backbone-Backbone	243	7.0	14	5.8	0.4	1	0.4	0.0
Backbone-Sidechain	344	9.9	105	30.5	3.0	28	8.1	0.8
Sidechain-Sidechain	43	1.2	29	67.4	0.8	10	23.3	0.3
Medium range ($ i-j >1 \& i-j <5$)	402	11.6	177	44.0	5.1	34	8.5	1.0
Backbone-Backbone	77	2.2	8	10.4	0.2	1	1.3	0.0
Backbone-Sidechain	194	5.6	78	40.2	2.3	18	9.3	0.5
Sidechain-Sidechain	131	3.8	91	69.5	2.6	15	11.5	0.4
Long range ($ i-j \ge 5$)	912	26.4	454	49.8	13.1	87	9.5	2.5
Backbone-Backbone	82	2.4	6	7.3	0.2	1	1.2	0.0
Backbone-Sidechain	350	10.1	159	45.4	4.6	32	9.1	0.9
Sidechain-Sidechain	480	13.9	289	60.2	8.4	54	11.2	1.6
Inter-chain	46	1.3	46	100.0	1.3	2	4.3	0.1
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	6	0.2	6	100.0	0.2	0	0.0	0.0
Sidechain-Sidechain	40	1.2	40	100.0	1.2	2	5.0	0.1
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	3460	100.0	1106	32.0	32.0	216	6.2	6.2
Backbone-Backbone	533	15.4	28	5.3	0.8	3	0.6	0.1
Backbone-Sidechain	1664	48.1	532	32.0	15.4	111	6.7	3.2
Sidechain-Sidechain	1263	36.5	546	43.2	15.8	102	8.1	2.9

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models



9.1.1 Bar chart: Distribution of distance restraints and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Madal ID		Nun	nber o	f viola	ations	5	M (8)	M (Å)	CD6 (Å)	Madian (8)
Model ID	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (Å)	Max (Å)	\mathbf{SD}^6 (Å)	Median (Å)
1	138	85	94	252	19	588	0.83	5.35	0.83	0.52
2	149	82	98	249	29	607	0.88	4.64	0.86	0.56
3	149	83	93	263	18	606	0.88	6.62	0.89	0.53
4	150	78	99	242	16	585	0.81	5.22	0.75	0.56
5	157	84	102	250	26	619	0.89	4.47	0.86	0.57
6	152	88	108	233	25	606	0.83	4.7	0.81	0.53
7	146	82	98	241	23	590	0.84	4.72	0.79	0.53
8	141	87	102	236	16	582	0.82	4.49	0.79	0.52
9	149	82	102	244	26	603	0.84	4.93	0.84	0.53
10	139	84	90	244	19	576	0.76	4.67	0.72	0.48
11	158	86	93	258	34	629	0.87	5.37	0.86	0.55




Continued from previous page...

Model ID		Nun	nber o	f viola	ations	3	Mean (Å)	Max (Å)	SD^6 (Å)	Median (Å)
Model 1D	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (A)	Max (A)	$SD^*(A)$	Median (A)
12	138	77	92	251	27	585	0.9	5.32	0.86	0.57
13	148	79	100	253	20	600	0.82	4.67	0.78	0.55
14	154	81	97	251	26	609	0.82	4.43	0.77	0.54
15	160	81	96	243	22	602	0.85	5.2	0.82	0.55
16	148	78	102	249	23	600	0.85	4.62	0.82	0.54
17	142	83	93	231	19	568	0.85	5.12	0.87	0.54
18	152	81	102	253	23	611	0.84	5.46	0.84	0.54
19	139	82	97	281	21	620	0.9	5.9	0.89	0.58
20	144	84	105	281	28	642	0.89	4.94	0.82	0.6

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Standard deviation

9.2.1 Bar graph: Distance Violation statistics for each model (i)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right



9.3 Distance violation statistics for the ensemble (i)


Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 2354(IR:1189, SQ:482, MR:225, LR:458, IC:0) restraints are not violated in the ensemble.

Nu	$\overline{\mathbf{mber}}$	of vio	lated	restra	aints	Fraction	n of the ensemble
IR^1	SQ^2	MR^3	LR^4	IC^5	Total	$Count^6$	%
45	23	16	69	2	155	1	5.0
24	11	9	19	3	66	2	10.0
19	9	15	16	3	62	3	15.0
12	6	14	29	1	62	4	20.0
3	5	11	16	3	38	5	25.0
8	4	4	13	4	33	6	30.0
10	1	5	17	1	34	7	35.0
10	4	1	15	3	33	8	40.0
7	3	4	8	1	23	9	45.0
7	3	8	9	1	28	10	50.0
4	7	2	11	3	27	11	55.0
9	2	4	15	4	34	12	60.0
4	1	1	10	4	20	13	65.0
6	2	6	17	3	34	14	70.0
6	6	8	22	2	44	15	75.0
9	4	5	16	0	34	16	80.0
10	5	9	23	4	51	17	85.0
9	2	9	9	1	30	18	90.0
25	11	12	33	1	82	19	95.0
54	39	34	87	2	216	20	100.0

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Number of models with violations



9.3.1 Bar graph: Distance violation statistics for the ensemble (i)




9.4 Most violated distance restraints in the ensemble (i)

9.4.1 Histogram: Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	\mathbf{SD}^1 (Å)	Median (Å)
(1,1669)	1:A:27:LEU:HD12	1:A:38:HIS:HD2	20	4.52	0.19	4.47
(1,1669)	1:A:27:LEU:HD11	1:A:38:HIS:HD2	20	4.52	0.19	4.47
(1,1669)	1:A:27:LEU:HD13	1:A:38:HIS:HD2	20	4.52	0.19	4.47
(1,936)	1:A:70:LYS:HG2	1:A:82:PRO:HB3	20	4.16	0.35	4.18
(1,1759)	1:A:94:ARG:HG2	1:A:58:MET:HE2	20	3.93	1.47	4.36
(1,1759)	1:A:94:ARG:HG2	1:A:58:MET:HE1	20	3.93	1.47	4.36
(1,1759)	1:A:94:ARG:HG2	1:A:58:MET:HE3	20	3.93	1.47	4.36
(1,541)	1:A:40:VAL:HG22	1:A:38:HIS:HD2	20	3.45	0.14	3.42
(1,541)	1:A:40:VAL:HG21	1:A:38:HIS:HD2	20	3.45	0.14	3.42
(1,541)	1:A:40:VAL:HG23	1:A:38:HIS:HD2	20	3.45	0.14	3.42
(1,1752)	1:A:57:LYS:HD2	1:A:56:GLU:HB3	20	3.1	0.82	3.33
(1,1224)	1:A:88:MET:HG3	1:A:59:MET:HG2	20	3.03	1.6	2.96
(1,955)	1:A:70:LYS:HD3	1:A:82:PRO:HD3	20	3.0	0.27	3.07
(1,241)	1:A:19:ASN:HA	1:A:3:HIS:HD2	20	2.85	0.85	3.2
(1,874)	1:A:65:LYS:HE2	1:A:111:TYR:HB2	20	2.82	1.33	3.06
(1,1067)	1:A:82:PRO:HD2	1:A:80:VAL:HG22	20	2.81	0.42	3.0




¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints (i)

9.5.1 Histogram: Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table: All distance violations (i)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,1029)	1:A:79:GLU:HG2	1:A:73:LYS:HE2	3	6.62
(1,1030)	1:A:79:GLU:HG3	1:A:73:LYS:HE2	3	6.35
(1,1208)	1:A:94:ARG:HD3	1:A:58:MET:HE2	19	5.9
(1,1208)	1:A:94:ARG:HD3	1:A:58:MET:HE1	19	5.9
(1,1208)	1:A:94:ARG:HD3	1:A:58:MET:HE3	19	5.9
(1,1759)	1:A:94:ARG:HG2	1:A:58:MET:HE2	18	5.46
(1,1759)	1:A:94:ARG:HG2	1:A:58:MET:HE1	18	5.46
(1,1759)	1:A:94:ARG:HG2	1:A:58:MET:HE3	18	5.46



Continued from previous page...

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,1759)	1:A:94:ARG:HG2	1:A:58:MET:HE2	19	5.44
(1,1759)	1:A:94:ARG:HG2	1:A:58:MET:HE1	19	5.44
(1,1759)	1:A:94:ARG:HG2	1:A:58:MET:HE3	19	5.44
(1,1224)	1:A:88:MET:HG3	1:A:59:MET:HG2	11	5.37
(1,1224)	1:A:88:MET:HG3	1:A:59:MET:HG2	1	5.35
(1,1207)	1:A:94:ARG:HD2	1:A:58:MET:HE2	3	5.35
(1,1207)	1:A:94:ARG:HD2	1:A:58:MET:HE1	3	5.35
(1,1207)	1:A:94:ARG:HD2	1:A:58:MET:HE3	3	5.35
(1,1759)	1:A:94:ARG:HG2	1:A:58:MET:HE2	12	5.32
(1,1759)	1:A:94:ARG:HG2	1:A:58:MET:HE1	12	5.32
(1,1759)	1:A:94:ARG:HG2	1:A:58:MET:HE3	12	5.32
(1,1759)	1:A:94:ARG:HG2	1:A:58:MET:HE2	11	5.3



10 Dihedral-angle violation analysis (i)


10.1 Summary of dihedral-angle violations (i)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle true	Count	$\%^{1}$	Vie	olated	3	Consis	tent	${ m ly~Violated^4}$
Angle type	Count	70	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^1$
PSI	105	49.8	19	18.1	9.0	3	2.9	1.4
PHI	106	50.2	19	17.9	9.0	5	4.7	2.4
Total	211	100.0	38	18.0	18.0	8	3.8	3.8

 $^{^1}$ percentage calculated with respect to total number of dihedral-angle restraints, 2 percentage calculated with respect to number of restraints in a particular dihedral-angle type, 3 violated in at least one model, 4 violated in all the models

10.1.1 Bar chart: Distribution of dihedral-angles and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories




10.2 Dihedral-angle violation statistics for each model (i)

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

Model ID	Nun	nber o	f violations	Mean (°)	Mov (°)	SD (°)	Modian (°)
Wiodei 1D	PSI	PHI	Total	Mean ()	$\mathbf{Max} \ (^{\circ})$	\mathbf{SD} (°)	\mid Median (°) \mid
1	8	12	20	3.72	11.2	3.24	1.9
2	8	8	16	4.05	11.7	3.51	2.25
3	7	9	16	4.47	11.7	3.54	2.6
4	8	10	18	3.92	12.3	3.54	2.35
5	7	9	16	4.39	12.2	3.56	3.1
6	9	10	19	3.87	12.9	3.73	1.8
7	8	8	16	4.36	12.2	3.64	3.4
8	7	10	17	4.31	11.1	3.23	2.6
9	9	12	21	3.58	12.5	3.4	1.9
10	10	11	21	3.6	11.7	3.27	1.9
11	8	12	20	3.89	12.0	3.41	2.45
12	15	8	23	3.22	12.1	3.28	1.8
13	9	12	21	4.0	12.5	3.45	2.3
14	5	12	17	4.2	11.2	3.45	2.5
15	8	11	19	3.94	12.1	3.39	2.3
16	7	8	15	4.45	11.9	3.57	2.7
17	6	10	16	4.41	11.0	3.22	2.85
18	8	8	16	4.33	12.4	3.64	2.95
19	9	10	19	3.83	12.1	3.53	2.3
20	9	9	18	3.82	11.9	3.49	2.05



10.2.1 Bar graph: Dihedral violation statistics for each model (i)



The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right

10.3 Dihedral-angle violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of ensemble.

Nun	nber o	f violated restraints	Fraction of the ensemble		
PSI	PHI	Total	Count ¹	%	
3	1	4	1	5.0	
1	2	3	2	10.0	
1	1	2	3	15.0	
4	2	6	4	20.0	
2	1	3	5	25.0	
0	1	1	6	30.0	
0	0	0	7	35.0	
0	1	1	8	40.0	
1	0	1	9	45.0	
0	1	1	10	50.0	
0	2	2	11	55.0	




Continued from previous page...

Nun	nber o	of violated restraints	Fraction of the ensemble			
PSI	PHI	Total	Count ¹	%		
0	0	0	12	60.0		
0	0	0	13	65.0		
0	0	0	14	70.0		
3	0	3	15	75.0		
0	2	2	16	80.0		
1	0	1	17	85.0		
0	0	0	18	90.0		
0	0	0	19	95.0		
3	5	8	20	100.0		

¹ Number of models with violations

10.3.1 Bar graph: Dihedral-angle Violation statistics for the ensemble (i)




10.4 Most violated dihedral-angle restraints in the ensemble (i)

10.4.1 Histogram: Distribution of mean dihedral-angle violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models



in the ensemble



10.4.2 Table: Most violated dihedral-angle restraints (i)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	\mathbf{Models}^1	Mean	\mathbf{SD}^2	Median
(1,29)	1:A:22:VAL:C	1:A:23:SER:N	1:A:23:SER:CA	1:A:23:SER:C	20	11.93	0.5	12.05
(1,31)	1:A:23:SER:C	1:A:24:PRO:N	1:A:24:PRO:CA	1:A:24:PRO:C	20	10.71	0.66	10.7
(1,28)	1:A:22:VAL:N	1:A:22:VAL:CA	1:A:22:VAL:C	1:A:23:SER:N	20	10.4	0.52	10.45
(1,32)	1:A:24:PRO:N	1:A:24:PRO:CA	1:A:24:PRO:C	1:A:25:ALA:N	20	5.6	0.51	5.4
(1,92)	1:A:58:MET:N	1:A:58:MET:CA	1:A:58:MET:C	1:A:59:MET:N	20	4.47	1.08	4.35
(1,115)	1:A:70:LYS:C	1:A:71:LYS:N	1:A:71:LYS:CA	1:A:71:LYS:C	20	4.01	1.22	3.9
(1,27)	1:A:21:ASP:C	1:A:22:VAL:N	1:A:22:VAL:CA	1:A:22:VAL:C	20	2.51	0.79	2.1
(1,170)	1:A:103:THR:C	1:A:104:LEU:N	1:A:104:LEU:CA	1:A:104:LEU:C	20	1.49	0.26	1.4
(1,125)	1:A:80:VAL:N	1:A:80:VAL:CA	1:A:80:VAL:C	1:A:81:VAL:N	17	1.85	0.66	1.5
(1,134)	1:A:84:PRO:C	1:A:85:GLN:N	1:A:85:GLN:CA	1:A:85:GLN:C	16	2.01	0.61	1.9


¹ Number of violated models, ²Standard deviation, All angle values are in degree (°)



10.5 All violated dihedral-angle restraints (i)

10.5.1 Histogram: Distribution of violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



10.5.2 Table: All violated dihedral-angle restraints (i)

The following table provides the list of violations for the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation (°)
(1,29)	1:A:22:VAL:C	1:A:23:SER:N	1:A:23:SER:CA	1:A:23:SER:C	6	12.9
(1,29)	1:A:22:VAL:C	1:A:23:SER:N	1:A:23:SER:CA	1:A:23:SER:C	9	12.5
(1,29)	1:A:22:VAL:C	1:A:23:SER:N	1:A:23:SER:CA	1:A:23:SER:C	13	12.5
(1,29)	1:A:22:VAL:C	1:A:23:SER:N	1:A:23:SER:CA	1:A:23:SER:C	18	12.4
(1,29)	1:A:22:VAL:C	1:A:23:SER:N	1:A:23:SER:CA	1:A:23:SER:C	4	12.3
(1,29)	1:A:22:VAL:C	1:A:23:SER:N	1:A:23:SER:CA	1:A:23:SER:C	5	12.2
(1,29)	1:A:22:VAL:C	1:A:23:SER:N	1:A:23:SER:CA	1:A:23:SER:C	7	12.2
(1,29)	1:A:22:VAL:C	1:A:23:SER:N	1:A:23:SER:CA	1:A:23:SER:C	12	12.1
(1,29)	1:A:22:VAL:C	1:A:23:SER:N	1:A:23:SER:CA	1:A:23:SER:C	15	12.1
(1,29)	1:A:22:VAL:C	1:A:23:SER:N	1:A:23:SER:CA	1:A:23:SER:C	19	12.1

