



Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 02:14 pm BST

PDB ID : 5VU7
Title : TNA polymerase, open ternary complex
Authors : Chim, N.; Chaput, J.C.
Deposited on : 2017-05-18
Resolution : 2.72 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3359 (2.74-2.70)
Clashscore	141614	3686 (2.74-2.70)
Ramachandran outliers	138981	3622 (2.74-2.70)
Sidechain outliers	138945	3623 (2.74-2.70)
RSRZ outliers	127900	3276 (2.74-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	774	 2% 74% 23%
2	T	16	 44% 38% 19%
3	P	12	 83% 17%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	FA2	A	803	-	-	-	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6746 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	756	6214	4003	1053	1142	16	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	141	ALA	ASP	engineered mutation	UNP D0VWU9
A	143	ALA	GLU	engineered mutation	UNP D0VWU9
A	147	HIS	GLU	engineered mutation	UNP D0VWU9
A	485	ARG	ALA	engineered mutation	UNP D0VWU9
A	584	LYS	GLU	engineered mutation	UNP D0VWU9
A	664	ILE	GLU	engineered mutation	UNP D0VWU9

- Molecule 2 is a DNA chain called DNA template.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	T	13	264	126	45	80	13	0	0	0

- Molecule 3 is a DNA chain called DNA/TNA hybrid primer.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	P	12	242	115	49	67	11	0	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is 5-(6-AMINO-9H-PURIN-9-YL)-4-HYDROXYTETRAHYDROFURAN-3-YL DIHYDROGEN PHOSPHATE (three-letter code: FA2) (formula: C₉H₁₂N₅O₆P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	0	0
			20	9	5	5	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase



- Molecule 2: DNA template



- Molecule 3: DNA/TNA hybrid primer



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, α , β , γ	66.55Å 112.61Å 145.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.10 – 2.72 66.55 – 2.72	Depositor EDS
% Data completeness (in resolution range)	97.4 (49.10-2.72) 91.6 (66.55-2.72)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.38 (at 2.73Å)	Xtrriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, R_{free}	0.214 , 0.268 0.214 , 0.268	Depositor DCC
R_{free} test set	2000 reflections (6.81%)	wwPDB-VP
Wilson B-factor (Å ²)	40.8	Xtrriage
Anisotropy	0.661	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 30.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	6746	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FA2, MG, 9O4, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/6353	0.45	0/8570
2	T	0.54	0/294	0.93	0/451
3	P	0.61	0/250	0.89	1/384 (0.3%)
All	All	0.30	0/6897	0.51	1/9405 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	P	7	DC	C1'-O4'-C4'	-5.61	104.49	110.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6214	0	6264	103	0
2	T	264	0	148	5	0
3	P	242	0	123	1	0
4	A	1	0	0	0	0
5	A	5	0	0	0	0
6	A	20	0	9	2	0
All	All	6746	0	6544	107	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (107) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:58:ARG:NH1	1:A:92:ASP:OD1	2.15	0.79
1:A:689:ARG:NH2	1:A:719:GLU:OE2	2.16	0.79
1:A:265:ARG:NH2	1:A:665:GLN:OE1	2.16	0.77
1:A:657:PRO:HA	1:A:660:LEU:HD12	1.66	0.76
1:A:685:ARG:NH2	1:A:711:GLY:O	2.19	0.75
1:A:398:GLU:HG2	1:A:585:ARG:HG2	1.68	0.73
1:A:683:ALA:HB2	1:A:699:ILE:HD11	1.74	0.69
1:A:743:ARG:NH2	2:T:11:DT:OP1	2.26	0.68
1:A:525:ILE:HD12	1:A:536:VAL:HG11	1.80	0.64
1:A:597:ILE:HB	1:A:603:ILE:HD13	1.78	0.64
1:A:713:ARG:NH2	1:A:728:ASP:OD1	2.30	0.64
1:A:329:MET:SD	1:A:485:ARG:NH1	2.70	0.63
1:A:45:ASP:OD2	1:A:70:LYS:NZ	2.30	0.63
1:A:674:LYS:NZ	3:P:6:DA:OP1	2.28	0.63
1:A:392:PRO:HB3	1:A:538:TYR:HD2	1.64	0.62
1:A:525:ILE:HG23	1:A:536:VAL:HG21	1.82	0.61
1:A:285:GLN:HG2	1:A:286:PRO:HD2	1.81	0.61
1:A:241:ILE:HG22	1:A:250:VAL:HG22	1.82	0.60
1:A:54:ILE:HG23	1:A:96:ILE:HD11	1.82	0.59
1:A:738:LEU:HD11	1:A:756:ARG:HB3	1.84	0.59
1:A:35:GLU:OE2	1:A:67:ARG:NH2	2.35	0.58
1:A:521:ILE:O	1:A:525:ILE:HG12	2.04	0.57
1:A:580:GLU:OE1	1:A:583:TYR:OH	2.19	0.57
1:A:419:SER:OG	1:A:421:ASP:OD1	2.13	0.56
1:A:622:THR:HG21	1:A:646:VAL:HG11	1.86	0.56
1:A:137:MET:HG2	1:A:205:VAL:HB	1.88	0.55
1:A:157:ILE:N	1:A:187:GLU:OE1	2.37	0.55
1:A:392:PRO:HD3	1:A:591:LYS:HG3	1.88	0.55
1:A:196:ARG:NH1	1:A:200:GLU:OE1	2.35	0.54
1:A:432:ASP:CG	1:A:443:LYS:HD2	2.28	0.54
1:A:397:TRP:HB2	1:A:400:ILE:HD11	1.89	0.53
1:A:22:GLU:HB2	1:A:27:LYS:HE3	1.90	0.53
1:A:589:VAL:HG23	1:A:590:THR:HG22	1.90	0.53
2:T:13:DC:H2"	2:T:14:DG:C8	2.44	0.52
1:A:734:GLU:OE1	1:A:756:ARG:NH1	2.43	0.52
1:A:334:SER:OG	1:A:339:GLN:O	2.21	0.52
1:A:403:LEU:HB3	1:A:577:LEU:HD21	1.92	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:803:FA2:H2	2:T:5:DT:C2	2.45	0.52
1:A:683:ALA:HB2	1:A:699:ILE:CD1	2.40	0.52
1:A:428:CYS:HB2	1:A:431:TYR:CZ	2.45	0.51
1:A:710:ILE:HD12	2:T:12:DT:H3'	1.92	0.51
1:A:145:LEU:HG	1:A:158:LEU:HD21	1.92	0.50
1:A:195:LEU:HD21	1:A:230:PHE:CD2	2.46	0.50
1:A:625:ARG:HE	1:A:642:ILE:HD11	1.76	0.50
1:A:207:ILE:HG22	1:A:319:THR:HG22	1.93	0.50
1:A:359:ARG:O	1:A:363:GLU:HG3	2.12	0.49
1:A:58:ARG:HG2	1:A:59:HIS:ND1	2.28	0.48
1:A:745:LEU:C	1:A:747:ALA:H	2.15	0.48
1:A:142:ILE:HG22	1:A:160:ILE:HG12	1.95	0.48
1:A:174:LYS:HE2	1:A:299:TRP:CE2	2.48	0.48
1:A:499:TYR:CZ	1:A:501:ARG:HB2	2.49	0.48
1:A:332:GLN:HE21	1:A:478:LEU:HD22	1.79	0.47
1:A:147:HIS:HB2	1:A:150:GLU:CD	2.34	0.47
1:A:597:ILE:HD11	1:A:601:GLY:HA2	1.96	0.47
1:A:464:LYS:HG2	1:A:483:GLN:HG3	1.97	0.47
1:A:548:ILE:HD11	1:A:559:LYS:HD3	1.96	0.47
1:A:123:ASP:HA	1:A:362:TYR:CE2	2.50	0.46
1:A:636:VAL:HG12	1:A:748:PHE:CE1	2.51	0.46
1:A:473:PRO:HG2	1:A:474:ILE:HD12	1.98	0.46
1:A:617:GLU:HB2	1:A:659:LYS:HA	1.98	0.46
1:A:670:LEU:HA	1:A:673:TYR:HD2	1.80	0.46
1:A:625:ARG:NE	1:A:642:ILE:HD11	2.31	0.46
1:A:293:GLU:O	1:A:297:THR:HG23	2.16	0.46
1:A:496:TYR:CZ	1:A:502:ALA:HB1	2.51	0.46
1:A:626:VAL:O	1:A:630:LEU:HG	2.17	0.45
1:A:644:LYS:HA	1:A:644:LYS:HD2	1.82	0.45
1:A:425:ARG:HE	1:A:445:PHE:HE1	1.64	0.45
1:A:406:ARG:NH2	1:A:576:GLU:OE1	2.48	0.45
2:T:4:DT:H2'	2:T:5:DT:C6	2.52	0.45
1:A:676:THR:HA	1:A:680:VAL:HG21	1.98	0.45
1:A:150:GLU:O	1:A:151:GLU:HB3	2.16	0.45
1:A:609:GLU:HA	1:A:612:ARG:HG3	1.99	0.45
1:A:89:HIS:CG	1:A:90:PRO:HD2	2.51	0.45
1:A:650:LEU:HB3	1:A:733:ILE:HD11	1.99	0.45
1:A:130:GLU:HG3	1:A:335:ARG:HE	1.81	0.45
1:A:505:TYR:OH	1:A:507:LYS:HB2	2.17	0.44
1:A:584:LYS:HD2	1:A:599:GLU:HA	2.00	0.44
1:A:670:LEU:HD23	1:A:684:LYS:HA	2.00	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:15:VAL:HG22	1:A:32:ARG:HG2	1.99	0.44
1:A:402:TYR:CE2	1:A:404:ASP:HB3	2.53	0.44
1:A:392:PRO:HB3	1:A:538:TYR:CD2	2.50	0.44
1:A:389:VAL:HG12	1:A:591:LYS:HG2	1.99	0.44
1:A:23:ASN:N	1:A:133:GLU:OE1	2.50	0.43
1:A:153:ALA:HA	1:A:218:TYR:CZ	2.53	0.43
1:A:91:GLN:O	1:A:94:PRO:HD2	2.18	0.43
1:A:528:ILE:HB	1:A:563:PHE:CZ	2.53	0.43
1:A:58:ARG:HD2	1:A:91:GLN:HB3	2.01	0.42
1:A:612:ARG:O	1:A:620:LYS:NZ	2.38	0.42
1:A:637:GLU:O	1:A:641:ARG:HG3	2.19	0.42
1:A:403:LEU:HB3	1:A:577:LEU:CD2	2.48	0.42
1:A:171:ILE:HG22	1:A:190:MET:HG3	2.01	0.42
1:A:460:ARG:O	1:A:464:LYS:HG3	2.20	0.42
1:A:642:ILE:O	1:A:645:GLU:HG2	2.20	0.42
6:A:803:FA2:H2'	6:A:803:FA2:H8	1.78	0.42
1:A:198:VAL:HG11	1:A:232:LEU:HD22	2.02	0.42
1:A:243:ARG:HA	1:A:248:PHE:HD1	1.84	0.42
1:A:528:ILE:HG12	1:A:534:PHE:HB2	2.02	0.42
1:A:207:ILE:HD13	1:A:323:GLY:HA3	2.02	0.41
1:A:610:ILE:HG13	1:A:623:GLN:HE21	1.86	0.41
1:A:158:LEU:HD22	1:A:299:TRP:CE2	2.55	0.41
1:A:417:ASN:HD21	1:A:446:PRO:HA	1.85	0.41
1:A:54:ILE:HG21	1:A:100:ILE:HD11	2.02	0.41
1:A:728:ASP:OD2	1:A:731:TYR:N	2.40	0.41
1:A:93:VAL:HG22	1:A:94:PRO:HD3	2.03	0.41
1:A:354:GLU:OE1	1:A:496:TYR:OH	2.18	0.41
1:A:656:PRO:HA	1:A:657:PRO:HD3	1.96	0.41
1:A:737:VAL:O	1:A:741:VAL:HG22	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	754/774 (97%)	719 (95%)	34 (4%)	1 (0%)	51 77

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	673	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	658/673 (98%)	639 (97%)	19 (3%)	42 70

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	LYS
1	A	44	ASP
1	A	179	PRO
1	A	215	ASP
1	A	315	ASP
1	A	324	LYS
1	A	327	LEU
1	A	354	GLU
1	A	483	GLN
1	A	497	TYR
1	A	516	TRP
1	A	567	ILE
1	A	575	LEU
1	A	577	LEU
1	A	630	LEU
1	A	668	ARG
1	A	694	ARG
1	A	719	GLU
1	A	751	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	9O4	P	12	3	16,21,22	5.66	11 (68%)	13,30,33	3.34	5 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	9O4	P	12	3	-	0/0/17/18	0/3/3/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	P	12	9O4	C2-N3	13.63	1.54	1.32
3	P	12	9O4	C4-N3	10.51	1.50	1.35
3	P	12	9O4	C8-N7	-7.13	1.22	1.34
3	P	12	9O4	C6-C5	-7.08	1.17	1.43
3	P	12	9O4	C6-N1	-5.26	1.14	1.37
3	P	12	9O4	C5-C4	4.33	1.52	1.40
3	P	12	9O4	C2'-C3'	-4.06	1.43	1.52
3	P	12	9O4	O4'-C4'	3.90	1.52	1.43
3	P	12	9O4	O4'-C1'	3.69	1.48	1.41
3	P	12	9O4	C6-N6	3.44	1.46	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	P	12	9O4	O3'-C3'	-2.06	1.43	1.45

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	P	12	9O4	C4-C5-N7	-8.07	100.99	109.40
3	P	12	9O4	C2-N1-C6	5.62	128.36	118.75
3	P	12	9O4	C2'-C3'-C4'	4.88	109.56	102.72
3	P	12	9O4	N3-C2-N1	-3.38	123.40	128.68
3	P	12	9O4	O4'-C1'-C2'	2.46	110.46	107.10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SO4	A	802	-	4,4,4	0.14	0	6,6,6	0.07	0
6	FA2	A	803	-	16,22,23	6.29	12 (75%)	15,32,35	3.03	6 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	FA2	A	803	-	-	1/1/21/22	0/3/3/3

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	803	FA2	C2-N3	13.68	1.54	1.32
6	A	803	FA2	O2'-C2'	-10.44	1.18	1.43
6	A	803	FA2	C4-N3	10.28	1.49	1.35
6	A	803	FA2	C8-N7	-7.20	1.21	1.34
6	A	803	FA2	C6-C5	-7.16	1.16	1.43
6	A	803	FA2	C6-N1	-5.25	1.14	1.37
6	A	803	FA2	C2'-C1'	-4.61	1.46	1.53
6	A	803	FA2	C5-C4	4.31	1.52	1.40
6	A	803	FA2	O4'-C1'	4.13	1.48	1.41
6	A	803	FA2	O4'-C4'	3.77	1.52	1.43
6	A	803	FA2	C2'-C3'	-3.69	1.43	1.53
6	A	803	FA2	C6-N6	3.35	1.46	1.34

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	803	FA2	C4-C5-N7	-7.65	101.43	109.40
6	A	803	FA2	C2-N1-C6	5.30	127.82	118.75
6	A	803	FA2	C4'-C3'-C2'	4.23	109.67	102.28
6	A	803	FA2	C3'-C2'-C1'	3.28	107.19	99.92
6	A	803	FA2	N3-C2-N1	-3.20	123.68	128.68
6	A	803	FA2	C1'-N9-C4	-2.64	122.00	126.64

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	A	803	FA2	C2'-C3'-O3'-P

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	803	FA2	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	756/774 (97%)	0.19	19 (2%) 57 59	26, 47, 90, 127	0
2	T	13/16 (81%)	-0.14	0 100 100	44, 56, 87, 94	0
3	P	11/12 (91%)	-0.21	0 100 100	49, 62, 94, 101	0
All	All	780/802 (97%)	0.18	19 (2%) 59 60	26, 48, 90, 127	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	471	ILE	3.9
1	A	148	GLU	3.9
1	A	691	VAL	2.9
1	A	589	VAL	2.9
1	A	247	ARG	2.8
1	A	394	ARG	2.7
1	A	746	ARG	2.7
1	A	608	LEU	2.7
1	A	744	ILE	2.7
1	A	549	PRO	2.6
1	A	741	VAL	2.6
1	A	747	ALA	2.4
1	A	742	GLU	2.3
1	A	145	LEU	2.2
1	A	698	VAL	2.2
1	A	749	GLY	2.1
1	A	755	LEU	2.1
1	A	646	VAL	2.1
1	A	717	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	9O4	P	12	19/20	0.96	0.19	22,39,49,51	0

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	FA2	A	803	20/21	0.77	0.72	34,39,75,81	20
4	MG	A	801	1/1	0.86	0.15	44,44,44,44	0
5	SO4	A	802	5/5	0.89	0.20	105,107,108,117	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.