



wwPDB X-ray Structure Validation Summary Report ⓘ

May 16, 2020 – 07:38 pm BST

PDB ID : 5X41
Title : 3.5A resolution structure of a cobalt energy-coupling factor transporter using LCP method-CbiMQO
Authors : Bao, Z.; Qi, X.; Zhao, W.; Li, D.; Zhang, P.
Deposited on : 2017-02-09
Resolution : 3.47 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.47 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1379 (3.56-3.40)
Clashscore	141614	1461 (3.56-3.40)
Ramachandran outliers	138981	1424 (3.56-3.40)
Sidechain outliers	138945	1425 (3.56-3.40)
RSRZ outliers	127900	1289 (3.56-3.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	280	
1	B	280	
1	C	280	
1	D	280	
2	E	222	
2	M	222	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	256	
3	Q	256	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 14359 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cobalt ABC transporter ATP-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	263	Total 1935	C 1221	N 349	O 362	S 3	0	0	0
1	B	258	Total 1899	C 1197	N 345	O 353	S 4	0	0	0
1	C	270	Total 1986	C 1251	N 362	O 369	S 4	0	0	0
1	D	267	Total 1964	C 1237	N 357	O 366	S 4	0	0	0

- Molecule 2 is a protein called Cobalt transport protein CbiM.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	M	207	Total 1470	C 975	N 236	O 252	S 7	0	0	0
2	E	207	Total 1470	C 975	N 236	O 252	S 7	0	0	0


- Molecule 3 is a protein called Uncharacterized protein CbiQ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	Q	243	Total 1815	C 1175	N 333	O 301	S 6	0	0	0
3	F	244	Total 1820	C 1178	N 334	O 302	S 6	0	0	0


LEU
ALA
ALA
GLN
MET
ALA
GLY
TRP
ARG
ARG

• Molecule 1: Cobalt ABC transporter ATP-binding protein







• Molecule 2: Cobalt transport protein CbiM



• Molecule 3: Uncharacterized protein CbiQ



• Molecule 3: Uncharacterized protein CbiQ





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	96.16Å 135.04Å 204.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.96 – 3.47 47.96 – 3.47	Depositor EDS
% Data completeness (in resolution range)	97.5 (47.96-3.47) 97.5 (47.96-3.47)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.53 (at 3.48Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.299 , 0.335 0.306 , 0.336	Depositor DCC
R_{free} test set	1727 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	99.3	Xtrriage
Anisotropy	0.800	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 63.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	14359	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.68	2/1960 (0.1%)	1.04	16/2665 (0.6%)
1	B	0.58	1/1923 (0.1%)	1.07	19/2611 (0.7%)
1	C	0.69	3/2013 (0.1%)	1.16	30/2736 (1.1%)
1	D	0.75	3/1990 (0.2%)	1.12	21/2704 (0.8%)
2	E	0.65	0/1504	0.96	7/2055 (0.3%)
2	M	0.64	2/1504 (0.1%)	0.95	10/2055 (0.5%)
3	F	0.70	2/1855 (0.1%)	1.25	31/2532 (1.2%)
3	Q	0.73	2/1850 (0.1%)	1.15	21/2525 (0.8%)
All	All	0.68	15/14599 (0.1%)	1.10	155/19883 (0.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	4
1	C	0	6
1	D	0	1
2	E	0	4
2	M	0	1
3	F	0	8
3	Q	0	13
All	All	0	39

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	242	PRO	N-CD	13.57	1.66	1.47
3	F	89	PRO	N-CD	5.77	1.55	1.47
1	A	200	VAL	CB-CG1	-5.75	1.40	1.52
3	Q	41	PRO	N-CD	5.61	1.55	1.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	57	PRO	N-CD	5.34	1.55	1.47

The worst 5 of 155 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	2	SER	N-CA-CB	-15.41	87.39	110.50
1	D	205	ALA	CB-CA-C	14.15	131.33	110.10
3	F	58	ALA	CB-CA-C	13.42	130.23	110.10
3	F	139	ALA	CB-CA-C	-13.00	90.61	110.10
3	Q	207	ARG	N-CA-CB	-11.81	89.35	110.60

There are no chirality outliers.

5 of 39 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	165	ASP	Peptide
1	A	216	GLY	Peptide
1	B	223	ALA	Peptide
1	B	65	GLY	Mainchain
1	B	95	PHE	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1935	0	2015	510	4
1	B	1899	0	1989	430	2
1	C	1986	0	2074	323	0
1	D	1964	0	2055	285	0
2	E	1470	0	1540	181	0
2	M	1470	0	1540	144	0
3	F	1820	0	1951	383	4
3	Q	1815	0	1948	357	2
All	All	14359	0	15112	2429	8

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 82.

The worst 5 of 2429 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:50:HIS:CE1	1:B:57:PRO:HG3	1.26	1.70
2:E:61:PRO:HD3	3:F:178:TRP:CZ3	1.25	1.63
1:C:243:LEU:CD2	1:D:243:LEU:HD21	1.16	1.61
1:C:252:ARG:CB	1:C:257:LEU:HD23	1.16	1.61
1:B:10:LEU:CD1	1:B:21:LEU:CD2	1.80	1.59

The worst 5 of 8 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:113:SER:CB	1:A:254:HIS:CE1[4_479]	1.33	0.87
1:B:251:ALA:CB	3:F:100:ALA:CB[2_374]	1.51	0.69
1:A:113:SER:OG	1:A:254:HIS:NE2[4_479]	1.58	0.62
1:A:113:SER:OG	1:A:254:HIS:CE1[4_479]	1.66	0.54
1:A:113:SER:CB	1:A:254:HIS:NE2[4_479]	2.02	0.18

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	259/280 (92%)	206 (80%)	36 (14%)	17 (7%)	1	12
1	B	254/280 (91%)	202 (80%)	38 (15%)	14 (6%)	2	16
1	C	268/280 (96%)	224 (84%)	32 (12%)	12 (4%)	2	20
1	D	265/280 (95%)	221 (83%)	33 (12%)	11 (4%)	3	22
2	E	205/222 (92%)	160 (78%)	39 (19%)	6 (3%)	4	29
2	M	205/222 (92%)	176 (86%)	20 (10%)	9 (4%)	2	20
3	F	242/256 (94%)	186 (77%)	42 (17%)	14 (6%)	1	14
3	Q	241/256 (94%)	182 (76%)	40 (17%)	19 (8%)	1	9

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	1939/2076 (93%)	1557 (80%)	280 (14%)	102 (5%)	2	16

5 of 102 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	63	LEU
1	A	114	GLU
1	A	140	MET
1	A	145	GLN
1	A	243	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	197/209 (94%)	142 (72%)	55 (28%)	0	2
1	B	194/209 (93%)	159 (82%)	35 (18%)	1	8
1	C	202/209 (97%)	168 (83%)	34 (17%)	2	11
1	D	200/209 (96%)	167 (84%)	33 (16%)	2	12
2	E	143/155 (92%)	123 (86%)	20 (14%)	3	17
2	M	143/155 (92%)	134 (94%)	9 (6%)	18	50
3	F	179/190 (94%)	155 (87%)	24 (13%)	4	19
3	Q	179/190 (94%)	145 (81%)	34 (19%)	1	7
All	All	1437/1526 (94%)	1193 (83%)	244 (17%)	2	11

5 of 244 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	Q	65	TRP
1	C	44	LYS
3	F	53	PHE
3	Q	135	TRP
3	Q	181	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
3	Q	208	ASN
3	Q	210	GLN
2	E	69	HIS
2	M	69	HIS
3	Q	86	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	263/280 (93%)	0.84	38 (14%) 2 3	28, 91, 141, 154	0
1	B	258/280 (92%)	0.73	27 (10%) 6 8	56, 80, 111, 132	0
1	C	270/280 (96%)	0.52	17 (6%) 20 20	54, 65, 99, 134	0
1	D	267/280 (95%)	0.45	24 (8%) 9 11	48, 62, 84, 113	0
2	E	207/222 (93%)	0.11	4 (1%) 66 63	53, 62, 81, 116	0
2	M	207/222 (93%)	0.27	9 (4%) 35 33	56, 62, 86, 147	0
3	F	244/256 (95%)	0.44	20 (8%) 11 13	53, 62, 114, 148	0
3	Q	243/256 (94%)	0.26	7 (2%) 51 48	51, 69, 85, 122	0
All	All	1959/2076 (94%)	0.47	146 (7%) 14 16	28, 68, 110, 154	0

The worst 5 of 146 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	142	SER	11.4
1	B	249	LEU	9.6
1	B	250	LEU	8.0
3	F	96	PRO	7.6
1	A	65	GLY	6.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.