

Full wwPDB EM Validation Report (i)

Nov 6, 2022 – 12:48 PM EST

PDB ID : 6C9A

EMDB ID : EMD-7435

Title: Cryo-EM structure of mouse TPC1 channel in the PtdIns(3,5)P2-bound state

Authors: She, J.; Guo, J.; Chen, Q.; Bai, X.; Jiang, Y.

Deposited on : 2018-01-25

Resolution : 3.20 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at

https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43

Mogul : 1.8.5 (274361), CSD as541be (2020)

MolProbity : 4.02b-467 buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

MapQ: 1.9.9

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.31.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# ext{Entries})$	${ m EM~structures} \ (\#{ m Entries})$
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
			20%		
1	A	825	81%	7%	12%
			22%		
1	В	825	81%	7%	12%
			100%		
2	$^{\mathrm{C}}$	2	100%		
			50%		
2	D	2	100%		



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 12182 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Two pore calcium channel protein 1.

Mol	Chain	Residues	\mathbf{Atoms}			AltConf	Trace		
1	В	723	Total 6002	C 3956	N 986	O 1022	S 38	0	0
1	A	723	Total 6002	C 3956	N 986	O 1022	S 38	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	818	VAL	-	expression tag	UNP Q9EQJ0
В	819	GLU	-	expression tag	UNP Q9EQJ0
В	820	ALA	-	expression tag	UNP Q9EQJ0
В	821	GLY	-	expression tag	UNP Q9EQJ0
В	822	LEU	-	expression tag	UNP Q9EQJ0
В	823	VAL	-	expression tag	UNP Q9EQJ0
В	824	PRO	-	expression tag	UNP Q9EQJ0
В	825	ARG	-	expression tag	UNP Q9EQJ0
A	818	VAL	-	expression tag	UNP Q9EQJ0
A	819	GLU	-	expression tag	UNP Q9EQJ0
A	820	ALA	-	expression tag	UNP Q9EQJ0
A	821	GLY	-	expression tag	UNP Q9EQJ0
A	822	LEU	-	expression tag	UNP Q9EQJ0
A	823	VAL	-	expression tag	UNP Q9EQJ0
A	824	PRO	-	expression tag	UNP Q9EQJ0
A	825	ARG	-	expression tag	UNP Q9EQJ0


• Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.





Mol	Chain	Residues	Atoms			AltConf	Trace	
2	С	2	Total 28				0	0
2	D	9	Total				0	0
2	D	2	28	16	2	10	U	U


 $\hbox{$\bullet$ Molecule 3 is (2R)-3-\{[(S)-hydroxy\{[(1S,2R,3R,4S,5S,6R)-2,4,6-trihydroxy-3,5-bis(phosphonoxy)cyclohexyl]oxy\}phosphoryl]oxy\}propane-1,2-diyl dioctanoate (three-letter code: EUJ) (formula: $C_{25}H_{49}O_{19}P_3$). }$



Mol	Chain	Residues	Atoms			AltConf	
2	B	1	Total				0
3	Б	1	47	25	19	3	0
2	Λ	1	Total				0
J	A	1	47	25	19	3	U

 \bullet Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $\rm C_8H_{15}NO_6).$






Mol	Chain	Residues	Atoms		AltConf
1	D	1	Total C N	О	0
4	Ъ	1	14 8 1	5	U
1	Λ	1	Total C N	О	0
4	A	1	14 8 1	5	U



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Two pore calcium channel protein 1 22% Chain B: GLN ASP SER GLU VAL ASP • Molecule 1: Two pore calcium channel protein 1 20% Chain A:



 \bullet Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C: 100%



 \bullet Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D: 100%





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	82819	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	50	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.215	Depositor
Minimum map value	-0.111	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.007	Depositor
Recommended contour level	0.0446	Depositor
Map size (Å)	256.80002, 256.80002, 256.80002	wwPDB
Map dimensions	240, 240, 240	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.07, 1.07, 1.07	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, EUJ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond	angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.45	0/6159	0.59	0/8339
1	В	0.45	0/6159	0.59	0/8339
All	All	0.45	0/12318	0.59	0/16678

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6002	0	6067	34	0
1	В	6002	0	6067	33	0
2	С	28	0	25	0	0
2	D	28	0	25	0	0
3	A	47	0	0	0	0
3	В	47	0	0	0	0
4	A	14	0	13	0	0
4	В	14	0	13	0	0
All	All	12182	0	12210	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.



All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A		Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:435:ILE:HG22	1:A:497:GLY:HA3	1.49	0.94
1:B:435:ILE:HG22	1:B:497:GLY:HA3	1.49	0.91
1:B:466:LEU:HD12	1:B:470:ASN:HB3	1.66	0.78
1:A:466:LEU:HD12	1:A:470:ASN:HB3	1.66	0.76
1:B:467:LYS:HG3	1:B:467:LYS:O	1.90	0.72
1:A:467:LYS:O	1:A:467:LYS:HG3	1.90	0.71
1:B:466:LEU:CD1	1:B:470:ASN:HB3	2.30	0.61
1:A:466:LEU:CD1	1:A:470:ASN:HB3	2.30	0.59
1:A:435:ILE:CG2	1:A:497:GLY:HA3	2.30	0.58
1:B:435:ILE:CG2	1:B:497:GLY:HA3	2.30	0.58
1:A:658:THR:HG21	1:A:666:ARG:HG3	1.87	0.57
1:B:658:THR:HG21	1:B:666:ARG:HG3	1.87	0.57
1:B:219:VAL:HG21	1:B:573:LEU:HD23	1.88	0.55
1:A:219:VAL:HG21	1:A:573:LEU:HD23	1.89	0.55
1:A:232:PRO:HG2	1:A:323:THR:HG21	1.90	0.53
1:B:232:PRO:HG2	1:B:323:THR:HG21	1.90	0.53
1:A:365:GLU:HA	1:A:380:ARG:HD3	1.90	0.53
1:A:219:VAL:HG13	1:A:574:THR:HG22	1.91	0.52
1:B:365:GLU:HA	1:B:380:ARG:HD3	1.90	0.52
1:B:219:VAL:HG13	1:B:574:THR:HG22	1.91	0.52
1:B:694:ARG:HG3	1:B:769:LEU:HD21	1.93	0.51
1:B:568:MET:HG2	1:B:687:ILE:HD13	1.94	0.49
1:A:694:ARG:HG3	1:A:769:LEU:HD21	1.93	0.49
1:B:633:ASN:ND2	1:B:636:ASN:OD1	2.45	0.49
1:A:568:MET:HG2	1:A:687:ILE:HD13	1.94	0.48
1:A:633:ASN:ND2	1:A:636:ASN:OD1	2.45	0.48
1:A:133:VAL:HG13	1:A:134:LEU:HD12	1.97	0.47
1:B:133:VAL:HG13	1:B:134:LEU:HD12	1.97	0.47
1:A:607:ALA:HB1	1:A:626:TYR:HD2	1.79	0.47
1:B:173:ARG:NE	1:B:212:ASP:OD1	2.48	0.47
1:B:607:ALA:HB1	1:B:626:TYR:HD2	1.79	0.47
1:A:173:ARG:NE	1:A:212:ASP:OD1	2.48	0.46
1:B:138:THR:OG1	1:B:195:GLN:OE1	2.33	0.46
1:B:356:PRO:HB2	1:B:394:LEU:HD11	1.98	0.46
1:A:138:THR:OG1	1:A:195:GLN:OE1	2.33	0.46
1:A:623:GLU:OE1	1:A:627:TYR:OH	2.34	0.45
1:B:171:HIS:HD2	1:B:174:THR:HG23	1.82	0.45
1:A:356:PRO:HB2	1:A:394:LEU:HD11	1.98	0.45
1:B:623:GLU:OE1	1:B:627:TYR:OH	2.34	0.44
1:B:435:ILE:HG22	1:B:497:GLY:CA	2.34	0.44

Continued on next page...



Continued from previous page...

Continued from prec		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)
1:A:435:ILE:HG22	1:A:497:GLY:CA	2.34	0.44
1:A:199:VAL:HG12	1:A:201:VAL:HG23	1.99	0.44
1:A:171:HIS:HD2	1:A:174:THR:HG23	1.82	0.44
1:A:81:GLU:HG3	1:A:86:ASP:HB3	2.00	0.43
1:A:405:GLU:HA	1:A:762:ARG:HH22	1.84	0.43
1:B:81:GLU:HG3	1:B:86:ASP:HB3	2.00	0.43
1:B:127:GLU:HG2	1:B:141:HIS:CD2	2.54	0.43
1:B:405:GLU:HA	1:B:762:ARG:HH22	1.84	0.43
1:B:467:LYS:O	1:B:467:LYS:CG	2.60	0.43
1:A:138:THR:HA	1:A:141:HIS:CE1	2.54	0.43
1:B:499:GLY:HA2	1:B:500:PRO:HD3	1.90	0.42
1:A:467:LYS:O	1:A:467:LYS:CG	2.60	0.42
1:A:638:PHE:HD1	1:A:638:PHE:HA	1.74	0.42
1:A:153:VAL:HG22	1:A:181:LEU:HD21	2.01	0.42
1:B:153:VAL:HG22	1:B:181:LEU:HD21	2.01	0.42
1:B:199:VAL:HG12	1:B:201:VAL:HG23	1.99	0.42
1:B:138:THR:HA	1:B:141:HIS:CE1	2.54	0.42
1:A:127:GLU:HG2	1:A:141:HIS:CD2	2.54	0.41
1:A:191:VAL:HG21	1:A:200:ARG:HH11	1.85	0.41
1:A:688:LEU:HD13	1:A:688:LEU:HA	1.92	0.41
1:A:218:GLY:O	1:A:222:ASN:ND2	2.54	0.41
1:B:191:VAL:HG21	1:B:200:ARG:HH11	1.86	0.41
1:B:283:PHE:HE2	1:A:643:GLU:HG2	1.86	0.41
1:B:350:LEU:HD21	1:B:367:LEU:HG	2.02	0.41
1:B:218:GLY:O	1:B:222:ASN:ND2	2.54	0.40
1:A:350:LEU:HD21	1:A:367:LEU:HG	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Favoured Allowed		Percentiles	
1	A	719/825 (87%)	683 (95%)	36 (5%)	0	100	100
1	В	719/825 (87%)	683 (95%)	36 (5%)	0	100	100
All	All	1438/1650 (87%)	1366 (95%)	72 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	658/744~(88%)	653 (99%)	5 (1%)	81 93		
1	В	658/744~(88%)	653 (99%)	5 (1%)	81 93		
All	All	1316/1488 (88%)	1306 (99%)	10 (1%)	82 93		

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	417	ARG
1	В	591	ASN
1	В	694	ARG
1	В	722	MET
1	В	762	ARG
1	A	417	ARG
1	A	591	ASN
1	A	694	ARG
1	A	722	MET
1	A	762	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	В	107	HIS
1	В	141	HIS
1	В	171	HIS

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
1	В	269	ASN
1	В	591	ASN
1	В	633	ASN
1	A	107	HIS
1	A	141	HIS
1	A	171	HIS
1	A	269	ASN
1	A	591	ASN
1	A	633	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
Moi Ty	туре			Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	С	1	1,2	14,14,15	0.31	0	17,19,21	0.59	0
2	NAG	С	2	2	14,14,15	0.34	0	17,19,21	0.50	0
2	NAG	D	1	1,2	14,14,15	0.24	0	17,19,21	0.57	0
2	NAG	D	2	2	14,14,15	0.35	0	17,19,21	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	С	1	1,2	-	4/6/23/26	0/1/1/1
2	NAG	С	2	2	-	2/6/23/26	0/1/1/1
2	NAG	D	1	1,2	-	4/6/23/26	0/1/1/1
2	NAG	D	2	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	2	NAG	O5-C5-C6-O6
2	С	2	NAG	O5-C5-C6-O6
2	С	2	NAG	C4-C5-C6-O6
2	D	2	NAG	C4-C5-C6-O6
2	С	1	NAG	C4-C5-C6-O6
2	С	1	NAG	O5-C5-C6-O6
2	D	1	NAG	C4-C5-C6-O6
2	С	1	NAG	C8-C7-N2-C2
2	С	1	NAG	O7-C7-N2-C2
2	D	1	NAG	C8-C7-N2-C2
2	D	1	NAG	O7-C7-N2-C2
2	D	1	NAG	O5-C5-C6-O6

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trmo	Chain	Res	Link	Bond lengths				Bond angles		
IVIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
3	EUJ	A	1000	-	47,47,47	1.11	6 (12%)	61,65,65	1.00	2 (3%)	
3	EUJ	В	1000	-	47,47,47	1.11	6 (12%)	61,65,65	1.01	2 (3%)	
4	NAG	A	1003	1	14,14,15	0.40	0	17,19,21	0.69	1 (5%)	



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
MIOI					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	В	1003	1	14,14,15	0.35	0	17,19,21	0.67	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EUJ	A	1000	-	-	15/44/68/68	0/1/1/1
3	EUJ	В	1000	-	-	14/44/68/68	0/1/1/1
4	NAG	A	1003	1	-	2/6/23/26	0/1/1/1
4	NAG	В	1003	1	-	2/6/23/26	0/1/1/1

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	Ideal(Å)
3	В	1000	EUJ	P5-O5	2.62	1.64	1.59
3	A	1000	EUJ	P5-O5	2.61	1.64	1.59
3	В	1000	EUJ	P3-O3	2.59	1.64	1.59
3	A	1000	EUJ	P3-O3	2.50	1.64	1.59
3	В	1000	EUJ	O2C-C2C	-2.36	1.40	1.46
3	A	1000	EUJ	O2C-C2C	-2.32	1.40	1.46
3	В	1000	EUJ	O3C-C1B	2.29	1.40	1.33
3	A	1000	EUJ	O3C-C1B	2.28	1.40	1.33
3	A	1000	EUJ	O3C-C3C	-2.25	1.40	1.45
3	В	1000	EUJ	O3C-C3C	-2.23	1.40	1.45
3	A	1000	EUJ	O2C-C1A	2.03	1.40	1.34
3	В	1000	EUJ	O2C-C1A	2.02	1.40	1.34

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
3	A	1000	EUJ	O2C-C1A-C2A	4.71	121.64	111.50
3	В	1000	EUJ	O2C-C1A-C2A	4.69	121.61	111.50
3	A	1000	EUJ	O3C-C1B-C2B	2.76	120.57	111.91
3	В	1000	EUJ	O3C-C1B-C2B	2.74	120.51	111.91
4	A	1003	NAG	C1-O5-C5	2.22	115.20	112.19
4	В	1003	NAG	C1-O5-C5	2.13	115.08	112.19

There are no chirality outliers.



All (33) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	В	1000	EUJ	C1C-O13-P1-O1
3	В	1000	EUJ	C1C-O13-P1-O11
3	A	1000	EUJ	C1C-O13-P1-O1
3	A	1000	EUJ	C1C-O13-P1-O11
3	В	1000	EUJ	C2A-C1A-O2C-C2C
3	A	1000	EUJ	C2A-C1A-O2C-C2C
3	В	1000	EUJ	O1A-C1A-O2C-C2C
3	A	1000	EUJ	O1A-C1A-O2C-C2C
3	В	1000	EUJ	C2B-C1B-O3C-C3C
3	A	1000	EUJ	C2B-C1B-O3C-C3C
4	В	1003	NAG	C8-C7-N2-C2
4	В	1003	NAG	O7-C7-N2-C2
4	A	1003	NAG	C8-C7-N2-C2
4	A	1003	NAG	O7-C7-N2-C2
3	В	1000	EUJ	O1B-C1B-O3C-C3C
3	A	1000	EUJ	O1B-C1B-O3C-C3C
3	A	1000	EUJ	C1A-C2A-C3A-C4A
3	В	1000	EUJ	C1A-C2A-C3A-C4A
3	A	1000	EUJ	C3B-C4B-C5B-C6B
3	В	1000	EUJ	C3A-C4A-C5A-C6A
3	В	1000	EUJ	C3B-C4B-C5B-C6B
3	A	1000	EUJ	C3A-C4A-C5A-C6A
3	В	1000	EUJ	C5A-C6A-C7A-C8A
3	В	1000	EUJ	C1C-C2C-O2C-C1A
3	A	1000	EUJ	C1C-C2C-O2C-C1A
3	В	1000	EUJ	C1C-C2C-C3C-O3C
3	A	1000	EUJ	C1C-C2C-C3C-O3C
3	A	1000	EUJ	C5A-C6A-C7A-C8A
3	В	1000	EUJ	O2C-C2C-C3C-O3C
3	A	1000	EUJ	O2C-C2C-C3C-O3C
3	A	1000	EUJ	C3-O3-P3-O33
3	A	1000	EUJ	C2A-C3A-C4A-C5A
3	В	1000	EUJ	C2A-C3A-C4A-C5A


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.



Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-7435. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)


6.1.1 Primary map




The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)


6.2.1 Primary map



X Index: 120



Y Index: 120




Z Index: 120



The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)


6.3.1 Primary map



The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views (i)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0446. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.



6.5 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is 46 nm^3 ; this corresponds to an approximate mass of 42 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)



^{*}Reported resolution corresponds to spatial frequency of 0.312 $\rm \AA^{-1}$



8 Fourier-Shell correlation (i)


This section was not generated. No FSC curve or half-maps provided.



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-7435 and PDB model 6C9A. Per-residue inclusion information can be found in section 3 on page 6.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.0446 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0446).



9.4 Atom inclusion (i)



At the recommended contour level, 67% of all backbone atoms, 58% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.0446) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score	
All	0.5794	0.4900	
A	0.5854	0.4920	
В	0.5768	0.4880	
С	0.1786	0.4600	
D	0.2500	0.4750	



