



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 25, 2024 – 03:40 PM EST

PDB ID : 6EC7  
Title : Glutamylation domain, TbtB, from thiomuracin biosynthesis  
Authors : Cogan, D.P.; Nair, S.K.  
Deposited on : 2018-08-07  
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	861	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6480 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lantibiotic dehydratase domain protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	796	6250	3966	1176	1092	16	0	1	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP D6Y502
A	-1	GLY	-	expression tag	UNP D6Y502
A	0	SER	-	expression tag	UNP D6Y502


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	230	Total	O	0	0
			230	230		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lantibiotic dehydratase domain protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	44.82Å 96.41Å 107.01Å 90.00° 98.76° 90.00°	Depositor
Resolution (Å)	105.76 – 2.15 105.76 – 2.15	Depositor EDS
% Data completeness (in resolution range)	100.0 (105.76-2.15) 100.0 (105.76-2.15)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.00 (at 2.14Å)	Xtrriage
Refinement program	REFMAC 5.8.0189	Depositor
R, $R_{free}$	0.214 , 0.277 0.221 , 0.277	Depositor DCC
$R_{free}$ test set	2496 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	40.2	Xtrriage
Anisotropy	0.663	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 46.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6480	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.49% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.74	0/6398	0.91	4/8702 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	218	ASN	N-CA-C	6.88	129.57	111.00
1	A	359	LEU	C-N-CD	-6.66	105.95	120.60
1	A	229	ARG	NE-CZ-NH2	5.13	122.87	120.30
1	A	764	SER	N-CA-C	-5.06	97.33	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6250	0	6339	53	0
2	A	230	0	0	2	0
All	All	6480	0	6339	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:318:PRO:O	1:A:322:ARG:HG3	1.70	0.91
1:A:727:GLU:HA	2:A:901:HOH:O	1.76	0.85
1:A:318:PRO:O	1:A:322:ARG:CG	2.32	0.78
1:A:252:ARG:HD2	2:A:1107:HOH:O	1.92	0.70
1:A:547:GLN:O	1:A:550:THR:HG22	1.97	0.65
1:A:761:ASP:OD1	1:A:764:SER:OG	2.16	0.63
1:A:322:ARG:HD2	1:A:337:ASP:OD1	1.99	0.62
1:A:331:ALA:O	1:A:332:GLU:HB2	2.00	0.62
1:A:297:PRO:O	1:A:301:THR:HG23	2.01	0.60
1:A:128:ARG:HD2	1:A:129:ALA:N	2.17	0.59
1:A:301:THR:HG22	1:A:304:ARG:NH2	2.17	0.59
1:A:23:GLU:HB3	1:A:850:ILE:HG22	1.84	0.57
1:A:362:THR:HG22	1:A:363:GLY:N	2.19	0.56
1:A:384:TYR:CZ	1:A:388:LEU:HD11	2.41	0.56
1:A:267:ARG:NH2	1:A:640:PRO:O	2.39	0.55
1:A:450:ARG:NH1	1:A:704:GLY:O	2.40	0.54
1:A:405:THR:HG23	1:A:405:THR:O	2.08	0.54
1:A:384:TYR:CE2	1:A:388:LEU:HD11	2.47	0.49
1:A:399:TRP:CD2	1:A:400:PRO:HD2	2.48	0.49
1:A:484:ALA:O	1:A:491:ARG:NH2	2.46	0.49
1:A:186:GLN:NE2	1:A:190:ARG:HH11	2.12	0.48
1:A:853:LEU:C	1:A:853:LEU:HD23	2.34	0.48
1:A:439:TRP:CE2	1:A:544:LEU:HD22	2.49	0.48
1:A:159:ARG:NH2	1:A:410:GLU:OE1	2.46	0.48
1:A:25:GLY:N	1:A:208:PHE:O	2.44	0.47
1:A:764:SER:O	1:A:768:ARG:HG3	2.15	0.46
1:A:446:LYS:HD3	1:A:701:ARG:HA	1.96	0.46
1:A:248:HIS:O	1:A:252:ARG:HG3	2.16	0.46
1:A:332:GLU:HG3	1:A:335:GLN:HE21	1.80	0.46
1:A:23:GLU:CB	1:A:850:ILE:HG22	2.46	0.45
1:A:318:PRO:O	1:A:322:ARG:HG2	2.12	0.45
1:A:362:THR:CG2	1:A:363:GLY:N	2.81	0.44
1:A:815:VAL:O	1:A:819:GLN:HG2	2.17	0.44
1:A:405:THR:O	1:A:406:ALA:HB3	2.18	0.43
1:A:64:LEU:HD11	1:A:82:LEU:HD22	1.99	0.43
1:A:128:ARG:CD	1:A:129:ALA:N	2.80	0.43
1:A:359:LEU:HA	1:A:360:PRO:HD3	1.31	0.43
1:A:437:ARG:HD2	1:A:561:CYS:HB2	2.00	0.43
1:A:218:ASN:C	1:A:218:ASN:HD22	2.22	0.43
1:A:609:GLU:HA	1:A:610:PRO:C	2.37	0.43
1:A:725:PRO:O	1:A:726:GLN:CB	2.66	0.42
1:A:218:ASN:C	1:A:218:ASN:ND2	2.73	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:178:GLU:O	1:A:181:ARG:CB	2.68	0.41
1:A:331:ALA:O	1:A:332:GLU:CB	2.68	0.41
1:A:583:GLY:N	1:A:855:GLU:OE1	2.53	0.41
1:A:159:ARG:NH2	1:A:238:VAL:HG21	2.35	0.41
1:A:560:SER:HB3	1:A:581:GLY:O	2.21	0.41
1:A:218:ASN:ND2	1:A:219:GLY:O	2.53	0.41
1:A:841[B]:ARG:NH1	1:A:848:ARG:O	2.54	0.41
1:A:756:ARG:NH2	1:A:762:ASP:O	2.54	0.41
1:A:333:PRO:O	1:A:336:VAL:HG12	2.21	0.41
1:A:204:PRO:HA	1:A:209:THR:OG1	2.21	0.40
1:A:178:GLU:O	1:A:181:ARG:HB3	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	783/861 (91%)	747 (95%)	26 (3%)	10 (1%)	<b>12</b>   <b>6</b>

All (10) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	218	ASN
1	A	331	ALA
1	A	640	PRO
1	A	750	ALA
1	A	205	TYR
1	A	277	ASP
1	A	398	MET
1	A	726	GLN
1	A	497	SER
1	A	182	ARG



### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	643/698 (92%)	614 (96%)	29 (4%)	27 24

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	46	GLN
1	A	94	LEU
1	A	128	ARG
1	A	135	GLU
1	A	174	ARG
1	A	181	ARG
1	A	182	ARG
1	A	186	GLN
1	A	191	LEU
1	A	197	ARG
1	A	203	SER
1	A	218	ASN
1	A	261	LEU
1	A	301	THR
1	A	322	ARG
1	A	332	GLU
1	A	397	MET
1	A	409	HIS
1	A	550	THR
1	A	572	ARG
1	A	609	GLU
1	A	640	PRO
1	A	648	ARG
1	A	673	ARG
1	A	733	ARG
1	A	755	ILE
1	A	774	ASP
1	A	820	ILE
1	A	827	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	186	GLN
1	A	218	ASN
1	A	335	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	796/861 (92%)	0.20	25 (3%) 49 58	26, 45, 80, 128	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	331	ALA	5.7
1	A	405	THR	4.9
1	A	218	ASN	4.7
1	A	319	MET	4.7
1	A	205	TYR	4.0
1	A	497	SER	3.7
1	A	404	THR	3.5
1	A	498	ALA	3.3
1	A	359	LEU	3.2
1	A	323	ASP	3.2
1	A	507	HIS	3.1
1	A	136	LEU	3.1
1	A	360	PRO	3.0
1	A	276	ALA	2.8
1	A	335	GLN	2.6
1	A	750	ALA	2.5
1	A	206	SER	2.5
1	A	630	VAL	2.4
1	A	186	GLN	2.4
1	A	261	LEU	2.3
1	A	377	HIS	2.2
1	A	645	ASP	2.1
1	A	561	CYS	2.1
1	A	325	LEU	2.1
1	A	72	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.