



Full wwPDB X-ray Structure Validation Report

May 23, 2020 – 11:24 am BST

PDB ID : 6EUA
Title : The fibrinogen-like domain of human Angptl3
Authors : Biterova, E.I.; Esmaeeli, M.E.; Alanen, H.I.; Saaranen, M.; Ruddock, L.W.
Deposited on : 2017-10-30
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	227	<p>2% 74% 15% 11%</p>
1	B	227	<p>0% 78% 9% 11%</p>
1	C	227	<p>4% 73% 14% 13%</p>

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 5139 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Angiotensinogen-converting enzyme 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	202	1684	1075	287	315	7	0	2	0
1	B	201	1663	1063	285	308	7	0	0	0
1	C	198	1631	1042	277	305	7	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	234	MET	-	initiating methionine	UNP Q9Y5C1
A	235	HIS	-	expression tag	UNP Q9Y5C1
A	236	HIS	-	expression tag	UNP Q9Y5C1
A	237	HIS	-	expression tag	UNP Q9Y5C1
A	238	HIS	-	expression tag	UNP Q9Y5C1
A	239	HIS	-	expression tag	UNP Q9Y5C1
A	240	HIS	-	expression tag	UNP Q9Y5C1
A	241	MET	-	expression tag	UNP Q9Y5C1
B	234	MET	-	initiating methionine	UNP Q9Y5C1
B	235	HIS	-	expression tag	UNP Q9Y5C1
B	236	HIS	-	expression tag	UNP Q9Y5C1
B	237	HIS	-	expression tag	UNP Q9Y5C1
B	238	HIS	-	expression tag	UNP Q9Y5C1
B	239	HIS	-	expression tag	UNP Q9Y5C1
B	240	HIS	-	expression tag	UNP Q9Y5C1
B	241	MET	-	expression tag	UNP Q9Y5C1
C	234	MET	-	initiating methionine	UNP Q9Y5C1
C	235	HIS	-	expression tag	UNP Q9Y5C1
C	236	HIS	-	expression tag	UNP Q9Y5C1
C	237	HIS	-	expression tag	UNP Q9Y5C1
C	238	HIS	-	expression tag	UNP Q9Y5C1
C	239	HIS	-	expression tag	UNP Q9Y5C1
C	240	HIS	-	expression tag	UNP Q9Y5C1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	241	MET	-	expression tag	UNP Q9Y5C1


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	63	Total O 63 63	0	0
2	B	55	Total O 55 55	0	0
2	C	43	Total O 43 43	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Angiotensin-related protein 3



- Molecule 1: Angiotensin-related protein 3



- Molecule 1: Angiotensin-related protein 3



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	60.05Å 63.65Å 169.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.99 – 2.10 48.99 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.3 (48.99-2.10) 99.3 (48.99-2.10)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.22 (at 2.10Å)	Xtrriage
Refinement program	PHENIX 1.12_2829	Depositor
R, R_{free}	0.194 , 0.254 0.194 , 0.254	Depositor DCC
R_{free} test set	1940 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	35.4	Xtrriage
Anisotropy	0.250	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 45.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5139	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/1739	0.54	0/2359
1	B	0.41	0/1716	0.56	0/2329
1	C	0.34	0/1682	0.51	0/2284
All	All	0.38	0/5137	0.54	0/6972

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1684	0	1565	19	0
1	B	1663	0	1544	15	0
1	C	1631	0	1515	18	0
2	A	63	0	0	1	0
2	B	55	0	0	0	0
2	C	43	0	0	1	0
All	All	5139	0	4624	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:339:LYS:HD3	1:B:341:ASN:HD21	1.50	0.76
1:A:268:GLN:NE2	2:A:501:HOH:O	2.21	0.72
1:C:356:THR:HG22	1:C:359:THR:H	1.60	0.65
1:A:405:HIS:CD2	1:A:406:ASP:H	2.17	0.62
1:A:337:ASP:OD2	1:A:341:ASN:HB2	2.01	0.61
1:A:373:ILE:HG12	1:A:411:ASN:HD22	1.66	0.61
1:B:339:LYS:HD3	1:B:341:ASN:ND2	2.16	0.60
1:A:365:ILE:HG22	1:A:366:THR:HG23	1.83	0.59
1:A:343:HIS:HB3	1:A:369:VAL:HB	1.88	0.56
1:C:396:GLU:HG2	1:C:410:GLU:HG2	1.86	0.56
1:C:452:HIS:ND1	1:C:453:PRO:O	2.38	0.53
1:B:339:LYS:HG2	1:B:339:LYS:O	2.08	0.53
1:A:340:ASP:OD1	1:A:445:LYS:NZ	2.38	0.52
1:B:286:GLN:HG2	1:B:449:MET:SD	2.49	0.51
1:C:435:LYS:HG3	1:C:441:LEU:HA	1.92	0.51
1:C:334:GLU:HB2	1:C:448:LYS:HB2	1.91	0.51
1:C:287:HIS:HD2	2:C:511:HOH:O	1.95	0.50
1:A:335:LEU:HB3	1:A:444:ILE:HD12	1.94	0.49
1:A:371:ASN:ND2	1:A:373:ILE:O	2.41	0.49
1:B:379:LEU:HD23	1:B:405:HIS:HB2	1.95	0.49
1:A:405:HIS:CD2	1:A:406:ASP:N	2.81	0.48
1:C:246:CYS:HA	1:C:249:ILE:HD12	1.95	0.48
1:C:342:LYS:HE3	1:C:342:LYS:HB3	1.61	0.48
1:A:243:PRO:HB3	1:A:248:THR:OG1	2.15	0.47
1:B:349:PHE:HB3	1:B:362:LEU:HD23	1.97	0.47
1:B:441:LEU:HD12	1:B:442:TYR:N	2.31	0.46
1:C:303:LYS:HG2	1:C:318:GLU:HG2	1.98	0.46
1:B:435:LYS:HB2	1:B:441:LEU:HA	1.98	0.46
1:B:452:HIS:HB2	1:B:453:PRO:HD2	1.98	0.45
1:B:294:ASN:ND2	1:B:296:ASN:H	2.15	0.45
1:A:282:TRP:HB3	1:A:450:LEU:HB3	1.98	0.45
1:A:245:GLU:OE1	1:A:247:THR:HB	2.17	0.45
1:A:349:PHE:HB3	1:A:362:LEU:HD23	1.99	0.45
1:C:365:ILE:HG22	1:C:366:THR:HG23	2.00	0.44
1:C:379:LEU:HB2	1:C:403:TRP:HB3	1.99	0.44
1:B:349:PHE:CB	1:B:362:LEU:HD23	2.48	0.44
1:B:362:LEU:HD12	1:B:373:ILE:HG21	2.00	0.44
1:C:353:ASN:O	1:C:356:THR:HB	2.17	0.44
1:C:289:ILE:N	1:C:312:GLU:OE1	2.48	0.43
1:A:437:GLN:HG2	1:A:437:GLN:H	1.51	0.43
1:A:370:PRO:HB2	1:A:436:SER:HB2	2.00	0.43
1:C:333:ILE:HB	1:C:345:ILE:HG23	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:328:ASN:HB2	1:A:350:TYR:CZ	2.55	0.42
1:B:317:LEU:HD13	1:B:358:TYR:CD1	2.55	0.41
1:C:379:LEU:HD23	1:C:405:HIS:HB2	2.01	0.41
1:C:282:TRP:HB3	1:C:450:LEU:HB3	2.03	0.41
1:A:303:LYS:HD3	1:A:304:TYR:CZ	2.56	0.41
1:B:273:TYR:OH	1:C:308:ARG:HD3	2.21	0.41
1:C:369:VAL:HG22	1:C:370:PRO:HD2	2.02	0.40
1:B:343:HIS:CG	1:B:369:VAL:HG21	2.56	0.40
1:A:438:ASN:OD1	1:A:438:ASN:N	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	198/227 (87%)	190 (96%)	8 (4%)	0	100	100
1	B	195/227 (86%)	190 (97%)	5 (3%)	0	100	100
1	C	192/227 (85%)	187 (97%)	5 (3%)	0	100	100
All	All	585/681 (86%)	567 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/203 (90%)	179 (98%)	3 (2%)	62	69
1	B	179/203 (88%)	175 (98%)	4 (2%)	52	57
1	C	176/203 (87%)	172 (98%)	4 (2%)	50	55
All	All	537/609 (88%)	526 (98%)	11 (2%)	53	60

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	245	GLU
1	A	308	ARG
1	A	430	ARG
1	B	265	SER
1	B	280	SER
1	B	286	GLN
1	B	452	HIS
1	C	329	TYR
1	C	342	LYS
1	C	353	ASN
1	C	440	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	271	HIS
1	A	328	ASN
1	A	405	HIS
1	B	294	ASN
1	B	296	ASN
1	B	343	HIS
1	C	266	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	202/227 (88%)	0.18	5 (2%) 57 62	24, 41, 75, 103	0
1	B	201/227 (88%)	0.10	2 (0%) 82 85	23, 38, 69, 119	0
1	C	198/227 (87%)	0.21	9 (4%) 33 38	24, 51, 93, 145	0
All	All	601/681 (88%)	0.17	16 (2%) 54 60	23, 43, 81, 145	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	437	GLN	5.2
1	B	398	TYR	5.1
1	C	438	ASN	4.7
1	B	391	HIS	4.1
1	C	339	LYS	3.6
1	C	437	GLN	3.4
1	C	355	GLU	3.3
1	C	398	TYR	2.9
1	A	370	PRO	2.5
1	C	328	ASN	2.4
1	C	394	CYS	2.3
1	C	393	ASN	2.2
1	A	436	SER	2.2
1	A	241	MET	2.1
1	A	440	ARG	2.0
1	C	326	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.