



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 8, 2023 – 02:36 PM EDT

PDB ID : 6I1P
Title : Respiratory complex I from *Thermus thermophilus* with bound NADH
Authors : Gutierrez-Fernandez, J.; Minhas, G.S.; Sazanov, L.A.
Deposited on : 2018-10-29
Resolution : 3.21 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<https://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : **FAILED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS failed to run properly.

Mol	Chain	Length	Quality of chain
1	1	438	60% 38% .
1	B	438	55% 44% .
2	2	181	71% 25% ..
2	C	181	60% 38% ..
3	3	783	57% 38% ..
3	D	783	59% 37% .
4	4	409	49% 43% . 6%
4	E	409	51% 42% . 6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	5	207	55% 39% • 5%
5	F	207	57% 36% • 5%
6	6	181	43% 46% • 8%
6	G	181	39% 51% • 8%
7	9	182	61% 37% ••
7	O	182	58% 39% ••
8	7	129	71% 26% ••
8	I	129	64% 33% ••
9	W	131	66% 30% ••
9	X	131	74% 23% •
10	A	119	56% 40% ••
10	P	119	61% 35% ••
11	J	176	54% 35% • 9%
11	R	176	55% 35% • 9%
12	K	95	68% 32%
12	S	95	77% 22% •
13	L	606	67% 33%
13	T	606	68% 32%
14	M	469	61% 38% •
14	U	469	65% 33% •
15	N	427	71% 29%
15	V	427	64% 36%
16	H	365	52% 43% ••
16	Q	365	49% 45% ••

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
17	SF4	1	501	-	-	X	-
17	SF4	3	803	-	-	X	-
17	SF4	9	201	-	-	X	-
17	SF4	9	202	-	-	X	-
17	SF4	G	201	-	-	X	-
17	SF4	O	202	-	-	X	-
20	FES	C	201	-	-	X	-
20	FES	D	804	-	-	X	-

2 Entry composition [i](#)

There are 20 unique types of molecules in this entry. The entry contains 74174 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NADH-quinone oxidoreductase subunit 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	1	437	3417	2180	595	624	18	0	0	0
1	B	437	3417	2180	595	624	18	0	0	0

- Molecule 2 is a protein called NADH-quinone oxidoreductase subunit 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	2	178	1406	895	238	265	8	0	0	0
2	C	178	1406	895	238	265	8	0	0	0

- Molecule 3 is a protein called NADH-quinone oxidoreductase subunit 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	3	756	5895	3754	1057	1053	31	0	0	0
3	D	756	5895	3754	1057	1053	31	0	0	0

- Molecule 4 is a protein called NADH-quinone oxidoreductase subunit 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	4	384	3067	1975	522	559	11	0	0	0
4	E	384	3067	1975	522	559	11	0	0	0

- Molecule 5 is a protein called NADH-quinone oxidoreductase subunit 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	5	196	Total 1607	C 1043	N 273	O 288	S 3	0	0	0
5	F	196	Total 1607	C 1043	N 273	O 288	S 3	0	0	0

- Molecule 6 is a protein called NADH-quinone oxidoreductase subunit 6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
6	6	166	Total 1289	C 815	N 235	O 226	S 13	0	0	0
6	G	166	Total 1289	C 815	N 235	O 226	S 13	0	0	0

- Molecule 7 is a protein called NADH-quinone oxidoreductase subunit 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
7	9	180	Total 1388	C 890	N 232	O 255	S 11	0	0	0
7	O	180	Total 1388	C 890	N 232	O 255	S 11	0	0	0

- Molecule 8 is a protein called NADH-quinone oxidoreductase subunit 15.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
8	7	127	Total 1031	C 664	N 183	O 181	S 3	0	0	0
8	I	127	Total 1031	C 664	N 183	O 181	S 3	0	0	0

- Molecule 9 is a protein called NADH-quinone oxidoreductase subunit 16.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
9	W	127	Total 967	C 623	N 165	O 175	S 4	0	0	0
9	X	127	Total 967	C 623	N 165	O 175	S 4	0	0	0

- Molecule 10 is a protein called NADH-quinone oxidoreductase subunit 7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
10	A	117	Total 910	C 624	N 138	O 144	S 4	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
10	P	117	910	624	138	144	4	0	0	0

- Molecule 11 is a protein called NADH-quinone oxidoreductase subunit 10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
11	J	160	1183	806	183	191	3	0	0	0
11	R	160	1183	806	183	191	3	0	0	0

- Molecule 12 is a protein called NADH-quinone oxidoreductase subunit 11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
12	K	95	703	456	118	126	3	0	0	0
12	S	95	703	456	118	126	3	0	0	0

- Molecule 13 is a protein called NADH-quinone oxidoreductase subunit 12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
13	L	605	4604	3089	740	756	19	0	0	0
13	T	605	4604	3089	740	756	19	0	0	0

- Molecule 14 is a protein called NADH-quinone oxidoreductase subunit 13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
14	M	467	3489	2363	546	572	8	0	0	0
14	U	467	3489	2363	546	572	8	0	0	0


- Molecule 15 is a protein called NADH-quinone oxidoreductase subunit 14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
15	N	427	3154	2125	505	518	6	0	0	0
15	V	427	3154	2125	505	518	6	0	0	0

- Molecule 16 is a protein called NADH-quinone oxidoreductase subunit 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
16	H	353	Total 2838	C 1943	N 431	O 457	S 7	0	0	0
16	Q	353	Total 2838	C 1943	N 431	O 457	S 7	0	0	0

- Molecule 17 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
			Total	Fe S			
17	1	1	Total 8	Fe 4	S 4	0	0
17	3	1	Total 8	Fe 4	S 4	0	0
17	3	1	Total 8	Fe 4	S 4	0	0
17	3	1	Total 8	Fe 4	S 4	0	0
17	6	1	Total 8	Fe 4	S 4	0	0
17	9	1	Total 8	Fe 4	S 4	0	0
17	9	1	Total 8	Fe 4	S 4	0	0
17	B	1	Total 8	Fe 4	S 4	0	0
17	D	1	Total 8	Fe 4	S 4	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
17	D	1	Total	Fe	S	0	0
			8	4	4		
17	D	1	Total	Fe	S	0	0
			8	4	4		
17	G	1	Total	Fe	S	0	0
			8	4	4		
17	O	1	Total	Fe	S	0	0
			8	4	4		
17	O	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 18 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: C₁₇H₂₁N₄O₉P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
18	1	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
18	B	1	Total	C	N	O	P	0	0
			31	17	4	9	1		

- Molecule 19 is 1,4-DIHYDRONICOTINAMIDE ADENINE DINUCLEOTIDE (three-letter code: NAI) (formula: C₂₁H₂₉N₇O₁₄P₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
19	1	1	44	21	7	14	2	0	0
19	B	1	44	21	7	14	2	0	0

- Molecule 20 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe₂S₂).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Fe S		
20	2	1	4	2 2	0	0
20	3	1	4	2 2	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
20	C	1	Total 4	Fe 2	S 2	0	0
20	D	1	Total 4	Fe 2	S 2	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS failed to run properly.

- Molecule 1: NADH-quinone oxidoreductase subunit 1




- Molecule 1: NADH-quinone oxidoreductase subunit 1








• Molecule 2: NADH-quinone oxidoreductase subunit 2




• Molecule 2: NADH-quinone oxidoreductase subunit 2




• Molecule 3: NADH-quinone oxidoreductase subunit 3






• Molecule 3: NADH-kinone oxidoreductase subunit 3




• Molecule 4: NADH-kinone oxidoreductase subunit 4





• Molecule 4: NADH-quinone oxidoreductase subunit 4

Chain E: 51% 42% 6%



• Molecule 5: NADH-quinone oxidoreductase subunit 5

Chain 5: 55% 39% 5%




• Molecule 5: NADH-quinone oxidoreductase subunit 5

Chain F: 57% 36% 5%




- Molecule 6: NADH-quinone oxidoreductase subunit 6

Chain 6: 43% 46% 8%




- Molecule 6: NADH-quinone oxidoreductase subunit 6

Chain G: 39% 51% 8%





- Molecule 7: NADH-quinone oxidoreductase subunit 9

Chain 9: 61% 37% 2%



- Molecule 7: NADH-quinone oxidoreductase subunit 9


Chain O: 58% 39% 3%




• Molecule 8: NADH-quinone oxidoreductase subunit 15




• Molecule 8: NADH-quinone oxidoreductase subunit 15





• Molecule 9: NADH-quinone oxidoreductase subunit 16




• Molecule 9: NADH-quinone oxidoreductase subunit 16




• Molecule 10: NADH-quinone oxidoreductase subunit 7




- Molecule 10: NADH-quinone oxidoreductase subunit 7




- Molecule 11: NADH-quinone oxidoreductase subunit 10




- Molecule 11: NADH-quinone oxidoreductase subunit 10



- Molecule 12: NADH-quinone oxidoreductase subunit 11




- Molecule 12: NADH-quinone oxidoreductase subunit 11




- Molecule 13: NADH-quinone oxidoreductase subunit 12






• Molecule 13: NADH-quinone oxidoreductase subunit 12




• Molecule 14: NADH-quinone oxidoreductase subunit 13






• Molecule 14: NADH-quinone oxidoreductase subunit 13

Chain U: 65% 33%




• Molecule 15: NADH-quinone oxidoreductase subunit 14

Chain N: 71% 29%




• Molecule 15: NADH-quinone oxidoreductase subunit 14

Chain V:  64% 36%




• Molecule 16: NADH-quinone oxidoreductase subunit 8


Chain H:  52% 43%



• Molecule 16: NADH-quinone oxidoreductase subunit 8

Chain Q:  49% 45%





4 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	96.11Å 341.45Å 263.89Å 90.00° 100.52° 90.00°	Depositor
Resolution (Å)	58.71 – 3.21	Depositor
% Data completeness (in resolution range)	76.1 (58.71-3.21)	Depositor
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.27 (at 3.19Å)	Xtrriage
Refinement program	PHENIX (1.13rc1_2961: ???)	Depositor
R, R_{free}	0.210 , 0.231	Depositor
Wilson B-factor (Å ²)	73.9	Xtrriage
Anisotropy	0.034	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.35$, $\langle L^2 \rangle = 0.17$	Xtrriage
Estimated twinning fraction	0.357 for h,-k,-h-l	Xtrriage
Reported twinning fraction	0.470 for -H,-K,H+L	Depositor
Outliers	0 of 207039 reflections	Xtrriage
Total number of atoms	74174	wwPDB-VP
Average B, all atoms (Å ²)	74.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: FES, NAI, SF4, FMN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	1	0.34	0/3506	0.51	0/4745
1	B	0.31	0/3506	0.49	0/4745
2	2	0.35	0/1439	0.50	0/1953
2	C	0.32	0/1439	0.50	0/1953
3	3	0.51	2/6035 (0.0%)	0.73	3/8185 (0.0%)
3	D	0.40	0/6035	0.60	0/8185
4	4	0.44	0/3150	0.62	0/4284
4	E	0.34	0/3150	0.51	0/4284
5	5	0.43	0/1656	0.66	1/2246 (0.0%)
5	F	0.37	0/1656	0.57	0/2246
6	6	0.46	0/1319	0.64	0/1786
6	G	0.46	0/1319	0.62	0/1786
7	9	0.53	1/1423 (0.1%)	0.65	0/1933
7	O	0.48	1/1423 (0.1%)	0.65	1/1933 (0.1%)
8	7	0.32	0/1059	0.53	0/1429
8	I	0.36	0/1059	0.57	0/1429
9	W	0.43	0/985	0.62	0/1335
9	X	0.37	0/985	0.57	0/1335
10	A	0.33	0/940	0.52	0/1280
10	P	0.35	0/940	0.51	0/1280
11	J	0.31	0/1206	0.50	0/1649
11	R	0.32	0/1206	0.53	0/1649
12	K	0.31	0/710	0.53	0/962
12	S	0.30	0/710	0.51	0/962
13	L	0.29	0/4741	0.47	0/6460
13	T	0.29	0/4741	0.48	1/6460 (0.0%)
14	M	0.30	0/3591	0.48	0/4896
14	U	0.32	0/3591	0.51	0/4896
15	N	0.31	0/3238	0.47	0/4434
15	V	0.34	0/3238	0.52	0/4434
16	H	0.34	0/2935	0.55	0/4014
16	Q	0.37	0/2935	0.56	0/4014

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
All	All	0.37	4/75866 (0.0%)	0.56	6/103182 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
7	9	0	1
7	O	0	2
10	P	0	1
16	H	0	3
16	Q	0	3
All	All	0	10

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	9	101	CYS	CB-SG	-6.08	1.72	1.82
3	3	263	CYS	CB-SG	-5.98	1.72	1.81
7	O	101	CYS	CB-SG	-5.30	1.73	1.81
3	3	381	LEU	C-N	-5.26	1.22	1.34

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	O	59	CYS	CA-CB-SG	7.71	127.88	114.00
3	3	276	ARG	NE-CZ-NH1	6.14	123.37	120.30
13	T	151	TYR	N-CA-CB	5.82	121.08	110.60
3	3	435	LEU	CB-CG-CD1	-5.49	101.67	111.00
5	5	189	ARG	NE-CZ-NH1	5.35	122.97	120.30

There are no chirality outliers.

5 of 10 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
7	9	20	LYS	Peptide
16	H	217	THR	Peptide
16	H	266	GLY	Peptide
16	H	43	GLN	Peptide
7	O	20	LYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1	3417	0	3389	116	0
1	B	3417	0	3388	152	0
2	2	1406	0	1373	37	0
2	C	1406	0	1373	71	0
3	3	5895	0	5929	232	0
3	D	5895	0	5930	218	0
4	4	3067	0	3049	167	0
4	E	3067	0	3049	162	0
5	5	1607	0	1574	77	0
5	F	1607	0	1574	76	0
6	6	1289	0	1298	84	0
6	G	1289	0	1298	100	0
7	9	1388	0	1383	72	0
7	O	1388	0	1383	69	0
8	7	1031	0	1029	36	0
8	I	1031	0	1029	36	0
9	W	967	0	1010	32	0
9	X	967	0	1010	23	0
10	A	910	0	939	53	0
10	P	910	0	939	53	0
11	J	1183	0	1286	63	0
11	R	1183	0	1286	56	0
12	K	703	0	747	29	0
12	S	703	0	747	17	0
13	L	4604	0	4734	162	0
13	T	4604	0	4734	136	0
14	M	3489	0	3606	131	0
14	U	3489	0	3606	124	0
15	N	3154	0	3343	98	0
15	V	3154	0	3343	113	0
16	H	2838	0	2903	154	0
16	Q	2838	0	2903	175	0
17	1	8	0	0	3	0
17	3	24	0	0	2	0
17	6	8	0	0	0	0
17	9	16	0	0	6	0
17	B	8	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
17	D	24	0	0	1	0
17	G	8	0	0	2	0
17	O	16	0	0	3	0
18	1	31	0	19	8	0
18	B	31	0	19	2	0
19	1	44	0	27	3	0
19	B	44	0	27	4	0
20	2	4	0	0	1	0
20	3	4	0	0	0	0
20	C	4	0	0	2	0
20	D	4	0	0	2	0
All	All	74174	0	75276	2718	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 2718 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:4:190:LEU:O	4:4:194:LEU:HD13	1.54	1.05
7:9:133:LYS:O	7:9:137:LEU:HD13	1.57	1.04
3:D:286:ASN:ND2	3:D:289:TRP:O	1.96	0.98
4:E:47:LEU:HD13	4:E:51:GLU:O	1.62	0.97
3:3:42:ILE:HD12	3:3:42:ILE:O	1.65	0.96

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	1	435/438 (99%)	403 (93%)	32 (7%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	435/438 (99%)	404 (93%)	31 (7%)	0	100	100
2	2	176/181 (97%)	168 (96%)	8 (4%)	0	100	100
2	C	176/181 (97%)	168 (96%)	8 (4%)	0	100	100
3	3	750/783 (96%)	696 (93%)	54 (7%)	0	100	100
3	D	750/783 (96%)	703 (94%)	47 (6%)	0	100	100
4	4	382/409 (93%)	359 (94%)	23 (6%)	0	100	100
4	E	382/409 (93%)	360 (94%)	22 (6%)	0	100	100
5	5	194/207 (94%)	185 (95%)	9 (5%)	0	100	100
5	F	194/207 (94%)	187 (96%)	7 (4%)	0	100	100
6	6	164/181 (91%)	147 (90%)	16 (10%)	1 (1%)	25	64
6	G	164/181 (91%)	145 (88%)	19 (12%)	0	100	100
7	9	178/182 (98%)	172 (97%)	6 (3%)	0	100	100
7	O	178/182 (98%)	171 (96%)	7 (4%)	0	100	100
8	7	125/129 (97%)	116 (93%)	9 (7%)	0	100	100
8	I	125/129 (97%)	115 (92%)	10 (8%)	0	100	100
9	W	125/131 (95%)	121 (97%)	4 (3%)	0	100	100
9	X	125/131 (95%)	121 (97%)	4 (3%)	0	100	100
10	A	115/119 (97%)	106 (92%)	9 (8%)	0	100	100
10	P	115/119 (97%)	106 (92%)	9 (8%)	0	100	100
11	J	158/176 (90%)	147 (93%)	11 (7%)	0	100	100
11	R	158/176 (90%)	146 (92%)	12 (8%)	0	100	100
12	K	93/95 (98%)	89 (96%)	4 (4%)	0	100	100
12	S	93/95 (98%)	89 (96%)	4 (4%)	0	100	100
13	L	603/606 (100%)	569 (94%)	33 (6%)	1 (0%)	47	79
13	T	603/606 (100%)	569 (94%)	33 (6%)	1 (0%)	47	79
14	M	465/469 (99%)	437 (94%)	28 (6%)	0	100	100
14	U	465/469 (99%)	437 (94%)	28 (6%)	0	100	100
15	N	425/427 (100%)	400 (94%)	25 (6%)	0	100	100
15	V	425/427 (100%)	404 (95%)	21 (5%)	0	100	100
16	H	351/365 (96%)	303 (86%)	42 (12%)	6 (2%)	9	42
16	Q	351/365 (96%)	302 (86%)	42 (12%)	7 (2%)	7	38

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	9478/9796 (97%)	8845 (93%)	617 (6%)	16 (0%)	47 79

5 of 16 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
16	H	51	VAL
16	Q	51	VAL
16	H	50	ARG
16	H	218	PRO
16	H	44	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	1	355/356 (100%)	344 (97%)	11 (3%)	40 72
1	B	355/356 (100%)	344 (97%)	11 (3%)	40 72
2	2	150/152 (99%)	144 (96%)	6 (4%)	31 66
2	C	150/152 (99%)	144 (96%)	6 (4%)	31 66
3	3	609/628 (97%)	597 (98%)	12 (2%)	55 80
3	D	609/628 (97%)	599 (98%)	10 (2%)	62 84
4	4	332/355 (94%)	325 (98%)	7 (2%)	53 79
4	E	332/355 (94%)	325 (98%)	7 (2%)	53 79
5	5	167/175 (95%)	166 (99%)	1 (1%)	86 94
5	F	167/175 (95%)	164 (98%)	3 (2%)	59 82
6	6	135/149 (91%)	125 (93%)	10 (7%)	13 46
6	G	135/149 (91%)	125 (93%)	10 (7%)	13 46
7	9	148/150 (99%)	145 (98%)	3 (2%)	55 80
7	O	148/150 (99%)	146 (99%)	2 (1%)	67 86
8	7	104/106 (98%)	103 (99%)	1 (1%)	76 90
8	I	104/106 (98%)	102 (98%)	2 (2%)	57 81

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
9	W	99/101 (98%)	98 (99%)	1 (1%)	76	90
9	X	99/101 (98%)	97 (98%)	2 (2%)	55	80
10	A	90/92 (98%)	86 (96%)	4 (4%)	28	64
10	P	90/92 (98%)	89 (99%)	1 (1%)	73	88
11	J	118/130 (91%)	114 (97%)	4 (3%)	37	70
11	R	118/130 (91%)	115 (98%)	3 (2%)	47	77
12	K	71/71 (100%)	69 (97%)	2 (3%)	43	74
12	S	71/71 (100%)	70 (99%)	1 (1%)	67	86
13	L	453/454 (100%)	446 (98%)	7 (2%)	65	85
13	T	453/454 (100%)	446 (98%)	7 (2%)	65	85
14	M	332/332 (100%)	326 (98%)	6 (2%)	59	82
14	U	332/332 (100%)	323 (97%)	9 (3%)	44	75
15	N	302/302 (100%)	298 (99%)	4 (1%)	69	87
15	V	302/302 (100%)	297 (98%)	5 (2%)	60	83
16	H	293/300 (98%)	286 (98%)	7 (2%)	49	77
16	Q	293/300 (98%)	288 (98%)	5 (2%)	60	83
All	All	7516/7706 (98%)	7346 (98%)	170 (2%)	50	78

5 of 170 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	E	143	LEU
12	S	28	PHE
4	E	262	PHE
6	G	156	LYS
13	T	554	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 57 such sidechains are listed below:

Mol	Chain	Res	Type
16	H	112	GLN
15	V	277	ASN
3	D	436	GLN
14	U	255	GLN
13	T	432	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

22 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
17	SF4	3	801	3	0,12,12	-	-	-		
20	FES	3	804	3	0,4,4	-	-	-		
17	SF4	9	201	7	0,12,12	-	-	-		
17	SF4	3	802	3	0,12,12	-	-	-		
17	SF4	D	803	3	0,12,12	-	-	-		
17	SF4	D	802	3	0,12,12	-	-	-		
17	SF4	B	501	1	0,12,12	-	-	-		
17	SF4	D	801	3	0,12,12	-	-	-		
17	SF4	O	202	7	0,12,12	-	-	-		
19	NAI	1	503	-	42,48,48	0.53	0	47,73,73	0.66	1 (2%)
17	SF4	1	501	1	0,12,12	-	-	-		
19	NAI	B	503	-	42,48,48	0.57	0	47,73,73	0.66	1 (2%)
17	SF4	9	202	7	0,12,12	-	-	-		
17	SF4	O	201	7	0,12,12	-	-	-		
17	SF4	6	201	6	0,12,12	-	-	-		
18	FMN	B	502	-	33,33,33	1.12	2 (6%)	48,50,50	1.27	9 (18%)
20	FES	D	804	3	0,4,4	-	-	-		

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
17	SF4	3	803	3	0,12,12	-	-	-		
17	SF4	G	201	6	0,12,12	-	-	-		
18	FMN	1	502	-	33,33,33	1.11	2 (6%)	48,50,50	1.31	9 (18%)
20	FES	2	201	2	0,4,4	-	-	-		
20	FES	C	201	2	0,4,4	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
17	SF4	3	801	3	-	-	0/6/5/5
20	FES	3	804	3	-	-	0/1/1/1
17	SF4	9	201	7	-	-	0/6/5/5
17	SF4	3	802	3	-	-	0/6/5/5
17	SF4	D	803	3	-	-	0/6/5/5
17	SF4	D	802	3	-	-	0/6/5/5
17	SF4	B	501	1	-	-	0/6/5/5
17	SF4	D	801	3	-	-	0/6/5/5
17	SF4	O	202	7	-	-	0/6/5/5
19	NAI	1	503	-	-	11/25/72/72	0/5/5/5
17	SF4	1	501	1	-	-	0/6/5/5
19	NAI	B	503	-	-	9/25/72/72	0/5/5/5
17	SF4	9	202	7	-	-	0/6/5/5
17	SF4	O	201	7	-	-	0/6/5/5
17	SF4	6	201	6	-	-	0/6/5/5
18	FMN	B	502	-	-	8/18/18/18	0/3/3/3
20	FES	D	804	3	-	-	0/1/1/1
17	SF4	3	803	3	-	-	0/6/5/5
17	SF4	G	201	6	-	-	0/6/5/5
18	FMN	1	502	-	-	8/18/18/18	0/3/3/3
20	FES	2	201	2	-	-	0/1/1/1
20	FES	C	201	2	-	-	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
18	B	502	FMN	C4A-N5	3.84	1.38	1.30
18	1	502	FMN	C4A-N5	3.69	1.37	1.30
18	B	502	FMN	C10-N1	2.69	1.38	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
18	1	502	FMN	C10-N1	2.30	1.37	1.33

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
18	B	502	FMN	C4-N3-C2	-3.04	120.02	125.64
18	B	502	FMN	O4-C4-C4A	-2.90	118.92	126.60
18	1	502	FMN	C4-N3-C2	-2.88	120.32	125.64
18	B	502	FMN	C4A-C4-N3	2.79	120.27	113.19
18	1	502	FMN	O4-C4-C4A	-2.59	119.73	126.60

There are no chirality outliers.

5 of 36 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
18	1	502	FMN	N10-C1'-C2'-O2'
18	1	502	FMN	N10-C1'-C2'-C3'
18	1	502	FMN	C1'-C2'-C3'-O3'
18	1	502	FMN	C1'-C2'-C3'-C4'
18	1	502	FMN	O2'-C2'-C3'-O3'


There are no ring outliers.


16 monomers are involved in 36 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
17	9	201	SF4	4	0
17	B	501	SF4	1	0
17	D	801	SF4	1	0
17	O	202	SF4	2	0
19	1	503	NAI	3	0
17	1	501	SF4	3	0
19	B	503	NAI	4	0
17	9	202	SF4	2	0
17	O	201	SF4	1	0
18	B	502	FMN	2	0
20	D	804	FES	2	0
17	3	803	SF4	2	0
17	G	201	SF4	2	0
18	1	502	FMN	8	0
20	2	201	FES	1	0
20	C	201	FES	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS failed to run properly - this section is therefore empty.

6.4 Ligands [i](#)

EDS failed to run properly - this section is therefore empty.

6.5 Other polymers [i](#)

EDS failed to run properly - this section is therefore empty.