



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 16, 2023 – 01:43 AM JST

PDB ID : 6KDY
Title : Crystal structure of the alpha beta heterodimer of human IDH3 in complex with NAD.
Authors : Sun, P.; Ding, J.
Deposited on : 2019-07-03
Resolution : 3.02 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2399 (3.04-3.00)
Clashscore	141614	2734 (3.04-3.00)
Ramachandran outliers	138981	2640 (3.04-3.00)
Sidechain outliers	138945	2643 (3.04-3.00)
RSRZ outliers	127900	2287 (3.04-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	341	74% 25% .
1	C	341	73% 24% ..
1	E	341	4% 71% 24% . .
1	G	341	2% 72% 26% .
2	B	356	% 69% 25% . .
2	D	356	72% 23% . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	356	 73% 22% . .
2	H	356	 77% 19% . .

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 20156 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Isocitrate dehydrogenase [NAD] subunit alpha, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	335	2472	1554	428	469	21	0	0	0
1	C	335	2406	1514	416	456	20	0	0	0
1	E	327	2322	1461	403	440	18	0	0	0
1	G	335	2407	1513	416	458	20	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP P50213
A	0	SER	-	expression tag	UNP P50213
C	-1	GLY	-	expression tag	UNP P50213
C	0	SER	-	expression tag	UNP P50213
E	-1	GLY	-	expression tag	UNP P50213
E	0	SER	-	expression tag	UNP P50213
G	-1	GLY	-	expression tag	UNP P50213
G	0	SER	-	expression tag	UNP P50213

- Molecule 2 is a protein called Isocitrate dehydrogenase [NAD] subunit beta, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	341	2583	1634	447	480	22	0	0	0
2	D	342	2571	1627	445	477	22	0	0	0
2	F	341	2559	1618	442	478	21	0	0	0
2	H	341	2565	1624	447	472	22	0	0	0

There are 64 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	341	GLU	-	expression tag	UNP O43837
B	342	ILE	-	expression tag	UNP O43837
B	343	CYS	-	expression tag	UNP O43837
B	344	ARG	-	expression tag	UNP O43837
B	345	ARG	-	expression tag	UNP O43837
B	346	VAL	-	expression tag	UNP O43837
B	347	LYS	-	expression tag	UNP O43837
B	348	ASP	-	expression tag	UNP O43837
B	349	LEU	-	expression tag	UNP O43837
B	350	ASP	-	expression tag	UNP O43837
B	351	GLU	-	expression tag	UNP O43837
B	352	ASN	-	expression tag	UNP O43837
B	353	LEU	-	expression tag	UNP O43837
B	354	TYR	-	expression tag	UNP O43837
B	355	PHE	-	expression tag	UNP O43837
B	356	GLN	-	expression tag	UNP O43837
D	341	GLU	-	expression tag	UNP O43837
D	342	ILE	-	expression tag	UNP O43837
D	343	CYS	-	expression tag	UNP O43837
D	344	ARG	-	expression tag	UNP O43837
D	345	ARG	-	expression tag	UNP O43837
D	346	VAL	-	expression tag	UNP O43837
D	347	LYS	-	expression tag	UNP O43837
D	348	ASP	-	expression tag	UNP O43837
D	349	LEU	-	expression tag	UNP O43837
D	350	ASP	-	expression tag	UNP O43837
D	351	GLU	-	expression tag	UNP O43837
D	352	ASN	-	expression tag	UNP O43837
D	353	LEU	-	expression tag	UNP O43837
D	354	TYR	-	expression tag	UNP O43837
D	355	PHE	-	expression tag	UNP O43837
D	356	GLN	-	expression tag	UNP O43837
F	341	GLU	-	expression tag	UNP O43837
F	342	ILE	-	expression tag	UNP O43837
F	343	CYS	-	expression tag	UNP O43837
F	344	ARG	-	expression tag	UNP O43837
F	345	ARG	-	expression tag	UNP O43837
F	346	VAL	-	expression tag	UNP O43837
F	347	LYS	-	expression tag	UNP O43837
F	348	ASP	-	expression tag	UNP O43837
F	349	LEU	-	expression tag	UNP O43837
F	350	ASP	-	expression tag	UNP O43837

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
F	351	GLU	-	expression tag	UNP O43837
F	352	ASN	-	expression tag	UNP O43837
F	353	LEU	-	expression tag	UNP O43837
F	354	TYR	-	expression tag	UNP O43837
F	355	PHE	-	expression tag	UNP O43837
F	356	GLN	-	expression tag	UNP O43837
H	341	GLU	-	expression tag	UNP O43837
H	342	ILE	-	expression tag	UNP O43837
H	343	CYS	-	expression tag	UNP O43837
H	344	ARG	-	expression tag	UNP O43837
H	345	ARG	-	expression tag	UNP O43837
H	346	VAL	-	expression tag	UNP O43837
H	347	LYS	-	expression tag	UNP O43837
H	348	ASP	-	expression tag	UNP O43837
H	349	LEU	-	expression tag	UNP O43837
H	350	ASP	-	expression tag	UNP O43837
H	351	GLU	-	expression tag	UNP O43837
H	352	ASN	-	expression tag	UNP O43837
H	353	LEU	-	expression tag	UNP O43837
H	354	TYR	-	expression tag	UNP O43837
H	355	PHE	-	expression tag	UNP O43837
H	356	GLN	-	expression tag	UNP O43837

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
3	A	1	Total 44	C 21	N 7	O 14	P 2	0	0
3	C	1	Total 44	C 21	N 7	O 14	P 2	0	0
3	E	1	Total 44	C 21	N 7	O 14	P 2	0	0
3	G	1	Total 44	C 21	N 7	O 14	P 2	0	0

- Molecule 4 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: C₈H₁₈O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	B	1	Total 13	C 8	O 5	0	0
4	D	1	Total 13	C 8	O 5	0	0

- Molecule 5 is NONAETHYLENE GLYCOL (three-letter code: 2PE) (formula: C₁₈H₃₈O₁₀).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	C	O	0	0
			28	18	10		

- Molecule 6 is 1-DEOXY-1-THIO-HEPTAETHYLENE GLYCOL (three-letter code: PE7) (formula: $C_{14}H_{30}O_7S$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	F	1	Total	C	O	S	0	0
			22	14	7	1		

- Molecule 7 is HEXAETHYLENE GLYCOL (three-letter code: P6G) (formula: $C_{12}H_{26}O_7$).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	H	1	Total	C O	0	0
			19	12 7		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Isocitrate dehydrogenase [NAD] subunit alpha, mitochondrial

Chain A: 




- Molecule 1: Isocitrate dehydrogenase [NAD] subunit alpha, mitochondrial


Chain C: 





- Molecule 1: Isocitrate dehydrogenase [NAD] subunit alpha, mitochondrial

Chain E: 







- Molecule 1: Isocitrate dehydrogenase [NAD] subunit alpha, mitochondrial



- Molecule 2: Isocitrate dehydrogenase [NAD] subunit beta, mitochondrial




- Molecule 2: Isocitrate dehydrogenase [NAD] subunit beta, mitochondrial





- Molecule 2: Isocitrate dehydrogenase [NAD] subunit beta, mitochondrial

Chain F:  73% 22% ..



- Molecule 2: Isocitrate dehydrogenase [NAD] subunit beta, mitochondrial

Chain H:  77% 19% ..



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	99.46Å 162.98Å 114.41Å 90.00° 100.31° 90.00°	Depositor
Resolution (Å)	48.93 – 3.02 48.93 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.2 (48.93-3.02) 99.2 (48.93-3.00)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.39 (at 3.01Å)	Xtrriage
Refinement program	PHENIX 1.14_3260	Depositor
R, R_{free}	0.197 , 0.246 0.196 , 0.244	Depositor DCC
R_{free} test set	3599 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	69.3	Xtrriage
Anisotropy	0.313	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 48.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	20156	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 2PE, P6G, PE7, NAD, PG4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	0/2515	0.64	1/3411 (0.0%)
1	C	0.42	0/2449	0.56	0/3337
1	E	0.44	0/2363	0.61	2/3226 (0.1%)
1	G	0.43	0/2450	0.57	0/3339
2	B	0.49	0/2630	0.66	0/3559
2	D	0.47	0/2618	0.66	0/3545
2	F	0.49	0/2606	0.65	0/3532
2	H	0.48	0/2611	0.65	0/3533
All	All	0.46	0/20242	0.63	3/27482 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	319	GLY	N-CA-C	6.98	130.55	113.10
1	E	235	LEU	CA-CB-CG	5.42	127.76	115.30
1	E	126	TYR	CB-CA-C	-5.04	100.33	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2472	0	2428	65	0
1	C	2406	0	2292	68	0
1	E	2322	0	2167	66	0
1	G	2407	0	2287	73	0
2	B	2583	0	2534	68	0
2	D	2571	0	2510	68	0
2	F	2559	0	2481	61	0
2	H	2565	0	2522	48	0
3	A	44	0	26	5	0
3	C	44	0	25	2	0
3	E	44	0	24	2	0
3	G	44	0	25	2	0
4	B	13	0	18	0	0
4	D	13	0	18	3	0
5	D	28	0	38	5	0
6	F	22	0	30	1	0
7	H	19	0	26	1	0
All	All	20156	0	19451	476	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 476 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:89:LYS:NZ	1:E:127:SER:CB	2.06	1.19
2:B:284:ASN:ND2	2:B:334:THR:HB	1.67	1.09
2:B:284:ASN:HD21	2:B:334:THR:HB	1.22	0.99
1:E:89:LYS:HZ3	1:E:127:SER:CB	1.72	0.98
1:G:130:GLU:OE1	2:H:191:GLY:N	1.95	0.98

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	333/341 (98%)	309 (93%)	24 (7%)	0	100	100
1	C	333/341 (98%)	311 (93%)	22 (7%)	0	100	100
1	E	323/341 (95%)	301 (93%)	22 (7%)	0	100	100
1	G	333/341 (98%)	315 (95%)	18 (5%)	0	100	100
2	B	339/356 (95%)	321 (95%)	18 (5%)	0	100	100
2	D	340/356 (96%)	317 (93%)	23 (7%)	0	100	100
2	F	339/356 (95%)	318 (94%)	21 (6%)	0	100	100
2	H	339/356 (95%)	318 (94%)	21 (6%)	0	100	100
All	All	2679/2788 (96%)	2510 (94%)	169 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	257/278 (92%)	254 (99%)	3 (1%)	71	89
1	C	239/278 (86%)	234 (98%)	5 (2%)	53	81
1	E	224/278 (81%)	220 (98%)	4 (2%)	59	84
1	G	239/278 (86%)	238 (100%)	1 (0%)	91	97
2	B	269/300 (90%)	258 (96%)	11 (4%)	30	66
2	D	265/300 (88%)	257 (97%)	8 (3%)	41	74
2	F	263/300 (88%)	259 (98%)	4 (2%)	65	86
2	H	265/300 (88%)	259 (98%)	6 (2%)	50	79
All	All	2021/2312 (87%)	1979 (98%)	42 (2%)	53	81

5 of 42 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	178	ARG
1	G	129	ILE
1	E	206	ASP
2	F	194	LEU
2	H	98	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	284	ASN
1	C	139	GLN
1	E	131	HIS
1	G	139	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	2PE	D	401	-	27,27,27	0.56	0	26,26,26	0.61	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	P6G	H	401	-	18,18,18	0.57	0	17,17,17	0.43	0
3	NAD	C	401	-	42,48,48	4.56	17 (40%)	50,73,73	2.09	13 (26%)
4	PG4	D	402	-	12,12,12	0.53	0	11,11,11	0.32	0
3	NAD	G	401	-	42,48,48	4.51	16 (38%)	50,73,73	1.59	8 (16%)
3	NAD	E	401	-	42,48,48	4.55	16 (38%)	50,73,73	1.97	11 (22%)
6	PE7	F	401	-	21,21,21	0.52	0	20,20,20	0.46	0
3	NAD	A	401	-	42,48,48	4.36	14 (33%)	50,73,73	1.76	10 (20%)
4	PG4	B	401	-	12,12,12	0.55	0	11,11,11	0.66	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	2PE	D	401	-	-	14/25/25/25	-
7	P6G	H	401	-	-	8/16/16/16	-
3	NAD	C	401	-	-	16/26/62/62	0/5/5/5
4	PG4	D	402	-	-	8/10/10/10	-
3	NAD	G	401	-	-	11/26/62/62	0/5/5/5
3	NAD	E	401	-	-	14/26/62/62	0/5/5/5
6	PE7	F	401	-	-	12/19/19/19	-
3	NAD	A	401	-	-	10/26/62/62	0/5/5/5
4	PG4	B	401	-	-	2/10/10/10	-

The worst 5 of 63 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	G	401	NAD	C2B-C1B	-16.55	1.28	1.53
3	A	401	NAD	C2B-C1B	-16.32	1.29	1.53
3	E	401	NAD	C2B-C1B	-16.28	1.29	1.53
3	C	401	NAD	C2B-C1B	-16.27	1.29	1.53
3	G	401	NAD	O4B-C1B	15.60	1.62	1.41

The worst 5 of 42 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	401	NAD	C3N-C7N-N7N	7.81	127.12	117.75
3	E	401	NAD	O7N-C7N-C3N	6.36	127.24	119.63

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	401	NAD	O7N-C7N-N7N	-5.95	114.13	122.58
3	A	401	NAD	N3A-C2A-N1A	-5.25	120.48	128.68
3	E	401	NAD	N3A-C2A-N1A	-5.03	120.81	128.68

There are no chirality outliers.

5 of 95 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	401	NAD	C5B-O5B-PA-O2A
3	A	401	NAD	C5B-O5B-PA-O3
3	A	401	NAD	PA-O3-PN-O5D
3	A	401	NAD	C3D-C4D-C5D-O5D
3	A	401	NAD	O4D-C1D-N1N-C2N


There are no ring outliers.


8 monomers are involved in 20 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	401	2PE	5	0
7	H	401	P6G	1	0
3	C	401	NAD	2	0
4	D	402	PG4	3	0
3	G	401	NAD	2	0
3	E	401	NAD	2	0
6	F	401	PE7	1	0
3	A	401	NAD	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	335/341 (98%)	-0.44	1 (0%) 94 83	41, 61, 88, 131	0
1	C	335/341 (98%)	-0.25	1 (0%) 94 83	38, 82, 125, 148	0
1	E	327/341 (95%)	-0.07	12 (3%) 41 17	41, 91, 123, 131	0
1	G	335/341 (98%)	-0.17	7 (2%) 63 34	48, 91, 123, 158	0
2	B	341/356 (95%)	-0.40	2 (0%) 89 72	38, 58, 78, 94	0
2	D	342/356 (96%)	-0.41	1 (0%) 94 83	34, 62, 89, 104	0
2	F	341/356 (95%)	-0.35	0 100 100	38, 63, 91, 100	0
2	H	341/356 (95%)	-0.47	1 (0%) 94 83	38, 61, 89, 105	0
All	All	2697/2788 (96%)	-0.32	25 (0%) 84 62	34, 68, 113, 158	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	300	ILE	3.8
1	E	296	HIS	3.5
1	G	47	PRO	3.5
1	G	55	SER	3.4
1	E	313	SER	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PG4	B	401	13/13	0.79	0.31	58,71,80,82	0
7	P6G	H	401	19/19	0.84	0.18	70,84,92,95	0
5	2PE	D	401	28/28	0.86	0.20	60,77,84,86	0
4	PG4	D	402	13/13	0.87	0.20	76,82,87,95	0
6	PE7	F	401	22/22	0.92	0.18	62,76,91,104	0
3	NAD	G	401	44/44	0.94	0.14	84,90,94,95	0
3	NAD	E	401	44/44	0.95	0.22	74,87,94,99	0
3	NAD	C	401	44/44	0.96	0.14	65,79,86,104	0
3	NAD	A	401	44/44	0.96	0.17	51,58,67,78	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around 2PE D 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around PE7 F 401:**




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)








Electron density around NAD E 401:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around NAD C 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.